

Some Strongly Almost Summable Sequence Spaces

Sunil K. Sharma and Ayhan Esi

ABSTRACT: In the present paper we introduce some strongly almost summable sequence spaces using ideal convergence and Musielak-Orlicz function $\mathcal{M} = (M_k)$ in n -normed spaces. We examine some topological properties of the resulting sequence spaces.

AMS Subject Classification: 40A05, 46A45, 46B70.

Keywords and Phrases: Paranorm space; I-convergence; Λ -convergent; Orlicz function; Musielak-Orlicz function; n -normed spaces.

1. Introduction and preliminaries

Mursaleen and Noman [18] introduced the notion of λ -convergent and λ -bounded sequences as follows:

Let $\lambda = (\lambda_k)_{k=1}^{\infty}$ be a strictly increasing sequence of positive real numbers tending to infinity i.e.

$$0 < \lambda_0 < \lambda_1 < \dots \text{ and } \lambda_k \rightarrow \infty \text{ as } k \rightarrow \infty$$

and said that a sequence $x = (x_k) \in w$ is λ -convergent to the number L , called the λ -limit of x if $\Lambda_m(x) \rightarrow L$ as $m \rightarrow \infty$, where

$$\lambda_m(x) = \frac{1}{\lambda_m} \sum_{k=1}^m (\lambda_k - \lambda_{k-1}) x_k.$$

The sequence $x = (x_k) \in w$ is λ -bounded if $\sup_m |\Lambda_m(x)| < \infty$. It is well known [18] that if $\lim_m x_m = a$ in the ordinary sense of convergence, then

$$\lim_m \left(\frac{1}{\lambda_m} \left(\sum_{k=1}^m (\lambda_k - \lambda_{k-1}) |x_k - a| \right) \right) = 0.$$

This implies that

$$\lim_m |\Lambda_m(x) - a| = \lim_m \left| \frac{1}{\lambda_m} \sum_{k=1}^m (\lambda_k - \lambda_{k-1}) (x_k - a) \right| = 0,$$

which yields that $\lim_m \Lambda_m(x) = a$ and hence $x = (x_k) \in w$ is λ -convergent to a . The concept of 2-normed spaces was initially developed by Gähler [6] in the mid of 1960's, while that of n -normed spaces one can see in Misiak [15]. Since then, many others have studied this concept and obtained various results, see Gunawan ([8, 9]) and Gunawan and Mashadi [10] and references therein. Let $n \in \mathbb{N}$ and X be a linear space over the field \mathbb{K} , where \mathbb{K} is the field of real or complex numbers of dimension d , where $d \geq n \geq 2$. A real valued function $\|\cdot, \dots, \cdot\|$ on X^n satisfying the following four conditions:

1. $\|x_1, x_2, \dots, x_n\| = 0$ if and only if x_1, x_2, \dots, x_n are linearly dependent in X ;
2. $\|x_1, x_2, \dots, x_n\|$ is invariant under permutation;
3. $\|\alpha x_1, x_2, \dots, x_n\| = |\alpha| \|x_1, x_2, \dots, x_n\|$ for any $\alpha \in \mathbb{K}$, and
4. $\|x + x', x_2, \dots, x_n\| \leq \|x, x_2, \dots, x_n\| + \|x', x_2, \dots, x_n\|$

is called a n -norm on X , and the pair $(X, \|\cdot, \dots, \cdot\|)$ is called a n -normed space over the field \mathbb{K} .

For example, we may take $X = \mathbb{R}^n$ being equipped with the Euclidean n -norm $\|x_1, x_2, \dots, x_n\|_E =$ the volume of the n -dimensional paralleliped spanned by the vectors x_1, x_2, \dots, x_n which may be given explicitly by the formula

$$\|x_1, x_2, \dots, x_n\|_E = |\det(x_{ij})|,$$

where $x_i = (x_{i1}, x_{i2}, \dots, x_{in}) \in \mathbb{R}^n$ for each $i = 1, 2, \dots, n$. Let $(X, \|\cdot, \dots, \cdot\|)$ be a n -normed space of dimension $d \geq n \geq 2$ and $\{a_1, a_2, \dots, a_n\}$ be linearly independent set in X . Then the following function $\|\cdot, \dots, \cdot\|_\infty$ on X^{n-1} defined by

$$\|x_1, x_2, \dots, x_{n-1}\|_\infty = \max\{\|x_1, x_2, \dots, x_{n-1}, a_i\| : i = 1, 2, \dots, n\}$$

defines an $(n-1)$ -norm on X with respect to $\{a_1, a_2, \dots, a_n\}$.

A sequence (x_k) in a n -normed space $(X, \|\cdot, \dots, \cdot\|)$ is said to converge to some $L \in X$ if

$$\lim_{k \rightarrow \infty} \|x_k - L, z_1, \dots, z_{n-1}\| = 0 \text{ for every } z_1, \dots, z_{n-1} \in X.$$

A sequence (x_k) in a n -normed space $(X, \|\cdot, \dots, \cdot\|)$ is said to be Cauchy if

$$\lim_{\substack{k \rightarrow \infty \\ p \rightarrow \infty}} \|x_k - x_p, z_1, \dots, z_{n-1}\| = 0 \text{ for every } z_1, \dots, z_{n-1} \in X.$$

If every Cauchy sequence in X converges to some $L \in X$, then X is said to be complete with respect to the n -norm. Any complete n -normed space is said to be n -Banach space.

An Orlicz function $M : [0, \infty) \rightarrow [0, \infty)$ is a continuous, non-decreasing and convex function such that $M(0) = 0$, $M(x) > 0$ for $x > 0$ and $M(x) \rightarrow \infty$ as $x \rightarrow \infty$.

Lindenstrauss and Tzafriri [13] used the idea of Orlicz function to define the following sequence space,

$$\ell_M = \left\{ x \in w : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty \right\},$$

which is called as an Orlicz sequence space. Also ℓ_M is a Banach space with the norm

$$\|x\| = \inf \left\{ \rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\}.$$

Also, it was shown in [13] that every Orlicz sequence space ℓ_M contains a subspace isomorphic to ℓ_p ($p \geq 1$). The Δ_2 -condition is equivalent to $M(Lx) \leq LM(x)$, for all L with $0 < L < 1$. An Orlicz function M can always be represented in the following integral form

$$M(x) = \int_0^x \eta(t) dt,$$

where η is known as the kernel of M , is right differentiable for $t \geq 0$, $\eta(0) = 0$, $\eta(t) > 0$, η is non-decreasing and $\eta(t) \rightarrow \infty$ as $t \rightarrow \infty$.

A sequence $\mathcal{M} = (M_k)$ of Orlicz function is called a Musielak-Orlicz function see ([14, 23]). A sequence $\mathcal{N} = (N_k)$ defined by

$$N_k(v) = \sup\{|v|u - M_k(u) : u \geq 0\}, \quad k = 1, 2, \dots$$

is called the complementary function of the Musielak-Orlicz function \mathcal{M} . For a given Musielak-Orlicz function \mathcal{M} , the Musielak-Orlicz sequence space $t_{\mathcal{M}}$ and its subspace $h_{\mathcal{M}}$ are defined as follows

$$t_{\mathcal{M}} = \left\{ x \in w : I_{\mathcal{M}}(cx) < \infty \text{ for some } c > 0 \right\},$$

$$h_{\mathcal{M}} = \left\{ x \in w : I_{\mathcal{M}}(cx) < \infty \text{ for all } c > 0 \right\},$$

where $I_{\mathcal{M}}$ is a convex modular defined by

$$I_{\mathcal{M}}(x) = \sum_{k=1}^{\infty} M_k(x_k), \quad x = (x_k) \in t_{\mathcal{M}}.$$

We consider $t_{\mathcal{M}}$ equipped with the Luxemburg norm

$$\|x\| = \inf \left\{ k > 0 : I_{\mathcal{M}}\left(\frac{x}{k}\right) \leq 1 \right\}$$

or equipped with the Orlicz norm

$$\|x\|^0 = \inf \left\{ \frac{1}{k} \left(1 + I_{\mathcal{M}}(kx) \right) : k > 0 \right\}.$$

Let X be a linear metric space. A function $p : X \rightarrow \mathbb{R}$ is called paranorm, if

1. $p(x) \geq 0$ for all $x \in X$,
2. $p(-x) = p(x)$ for all $x \in X$,
3. $p(x + y) \leq p(x) + p(y)$ for all $x, y \in X$,
4. if (λ_n) is a sequence of scalars with $\lambda_n \rightarrow \lambda$ as $n \rightarrow \infty$ and (x_n) is a sequence of vectors with $p(x_n - x) \rightarrow 0$ as $n \rightarrow \infty$, then $p(\lambda_n x_n - \lambda x) \rightarrow 0$ as $n \rightarrow \infty$.

A paranorm p for which $p(x) = 0$ implies $x = 0$ is called total paranorm and the pair (X, p) is called a total paranormed space. It is well known that the metric of any linear metric space is given by some total paranorm (see [30], Theorem 10.4.2, p. 183). For more details about sequence spaces (see [16, 17, 19, 20, 21, 22, 24, 25, 26, 27, 29]) and reference therein.

A sequence space E is said to be solid (or normal) if $(x_k) \in E$ implies $(\alpha_k x_k) \in E$ for all sequences of scalars (α_k) with $|\alpha_k| \leq 1$ and for all $k \in \mathbb{N}$.

The notion of ideal convergence was introduced first by P. Kostyrko [11] as a generalization of statistical convergence which was further studied in topological spaces (see [2]). More applications of ideals can be seen in [2, 3].

A linear functional \mathcal{L} on ℓ_∞ is said to be a Banach limit see [1] if it has the properties:

1. $\mathcal{L}(x) \geq 0$ if $x \geq 0$ (i.e. $x_n \geq 0$ for all n),
2. $\mathcal{L}(e) = 1$, where $e = (1, 1, \dots)$,
3. $\mathcal{L}(Dx) = \mathcal{L}(x)$,

where the shift operator D is defined by $(Dx_n) = (x_{n+1})$.

Let \mathfrak{B} be the set of all Banach limits on ℓ_∞ . A sequence x is said to be almost convergent to a number L if $\mathcal{L}(x) = L$ for all $\mathcal{L} \in \mathfrak{B}$. Lorentz [12] has shown that x is almost convergent to L if and only if

$$t_{km} = t_{km}(x) = \frac{x_m + x_{m+1} + \dots + x_{m+k}}{k+1} \rightarrow L \text{ as } k \rightarrow \infty, \text{ uniformly in } m.$$

Recently a lot of activities have started to study sumability, sequence spaces and related topics in these non linear spaces see [4, 28]. In particular Sahiner [28] combined these two concepts and investigated ideal sumability in these spaces and introduced certain sequence spaces using 2-norm.

We continue in this direction and by using Musielak-Orlicz function, generalized sequences and also ideals we introduce I-convergence of generalized sequences with respect to Musielak-Orlicz function in n -normed spaces.

Let $(X, \|\cdot\|)$ be a normed space. Recall that a sequence $(x_n)_{n \in \mathbb{N}}$ of elements of X is called statistically convergent to $x \in X$ if the set $A(\epsilon) = \{n \in \mathbb{N} : \|x_n - x\| \geq \epsilon\}$ has natural density zero for each $\epsilon > 0$.

A family $\mathcal{I} \subset 2^Y$ of subsets of a non empty set Y is said to be an ideal in Y if

1. $\phi \in \mathcal{I}$;
2. $A, B \in \mathcal{I}$ imply $A \cup B \in \mathcal{I}$;
3. $A \in \mathcal{I}, B \subset A$ imply $B \in \mathcal{I}$, while an admissible ideal \mathcal{I} of Y further satisfies $\{x\} \in \mathcal{I}$ for each $x \in Y$ (see [6]).

Given $\mathcal{I} \subset 2^{\mathbb{N}}$ be a non trivial ideal in \mathbb{N} . A sequence $(x_n)_{n \in \mathbb{N}}$ in X is said to be \mathcal{I} -convergent to $x \in X$, if for each $\epsilon > 0$ the set $A(\epsilon) = \{n \in \mathbb{N} : \|x_n - x\| \geq \epsilon\}$ belongs to \mathcal{I} (see [11]).

Let I be an admissible ideal of \mathbb{N} , $\mathcal{M} = (M_k)$ be a Musielak-Orlicz function and $(X, \|\cdot, \dots, \cdot\|)$ be a n -normed space. Let $p = (p_k)$ be a bounded sequence of positive real numbers and $u = (u_k)$ be any sequence of strictly positive real numbers. By $S(n - X)$ we denote the space of all sequences defined over $(X, \|\cdot, \dots, \cdot\|)$. We define the following sequence spaces in this paper:

$$\hat{w}^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|) =$$

$$\left\{ x = (x_k) \in S(n - X) : \forall \epsilon > 0, \left\{ n \in \mathbb{N} : \right.$$

$$\left. \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x)) - L}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq \epsilon \right\} \in I$$

$$\text{for some } \rho > 0, L \in X \text{ and } z_1, \dots, z_{n-1} \in X \left. \right\},$$

$$\hat{w}_0^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|) =$$

$$\left\{ x = (x_k) \in S(n - X) : \forall \epsilon > 0, \left\{ n \in \mathbb{N} : \right.$$

$$\left. \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq \epsilon \right\} \in I$$

$$\text{for some } \rho > 0, \text{ and } z_1, \dots, z_{n-1} \in X \left. \right\},$$

$$\hat{w}_\infty(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|) =$$

$$\left\{ x = (x_k) \in S(n - X) : \exists K > 0 \text{ such that} \right.$$

$$\sup_{n \in \mathbb{N}} \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \leq K$$

$$\text{for some } \rho > 0, \text{ and } z_1, \dots, z_{n-1} \in X \left. \right\},$$

$$\hat{w}_\infty^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|) =$$

$$\left\{ x = (x_k) \in S(n - X) : \exists K > 0 \text{ such that} \right.$$

$$\left\{ n \in \mathbb{N} : \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq K \right\} \in I$$

for some $\rho > 0$, and $z_1, \dots, z_{n-1} \in X$.

The following inequality will be used throughout the paper. If $0 \leq p_k \leq \sup p_k = H$, $D = \max(1, 2^{H-1})$ then

$$|a_k + b_k|^{p_k} \leq D\{|a_k|^{p_k} + |b_k|^{p_k}\}$$

for all k and $a_k, b_k \in \mathbb{C}$. Also $|a|^{p_k} \leq \max(1, |a|^H)$ for all $a \in \mathbb{C}$.

The main aim of this paper is to study some topological properties and inclusion relations between the above defined sequence spaces.

2. Main results

Theorem 2.1. *Let $\mathcal{M} = (M_k)$ be a Musielak-Orlicz function, $p = (p_k)$ be a bounded sequence of positive real numbers and I be an admissible ideal of \mathbb{N} . Then $\hat{w}^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$, $\hat{w}_0^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$, $\hat{w}_\infty(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$ and $\hat{w}_\infty^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$ are linear spaces.*

Proof. Let $x, y \in \hat{w}^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$ and $\alpha, \beta \in \mathbb{C}$. So

$$\left\{ \left\{ n \in \mathbb{N} : \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x) - L)}{\rho_1}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq \epsilon \right\} \in I \text{ for some } \rho_1 > 0, \right.$$

$$\left. L \in X \text{ and } z_1, \dots, z_{n-1} \in X \right\}$$

and

$$\left\{ \left\{ n \in \mathbb{N} : \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(y) - L)}{\rho_2}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq \epsilon \right\} \in I \text{ for some } \rho_2 > 0, \right.$$

$$\left. L \in X \text{ and } z_1, \dots, z_{n-1} \in X \right\}.$$

Since $\|\cdot, \dots, \cdot\|$ is a n -norm, $\mathcal{M} = (M_k)$ be a Musielak-Orlicz function and so by using inequality (1.1), we have

$$\begin{aligned}
 & \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(\alpha x + \beta y) - L)}{|\alpha|\rho_1 + |\beta|\rho_2}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \\
 & \leq D \frac{1}{n} \sum_{k=1}^n \left[\frac{|\alpha|}{(|\alpha|\rho_1 + |\beta|\rho_2)} M_k \left(\left\| \frac{t_{km}(\Lambda_k(x) - L)}{\rho_1}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \\
 & + D \frac{1}{n} \sum_{k=1}^n \left[\frac{|\beta|}{(|\alpha|\rho_1 + |\beta|\rho_2)} M_k \left(\left\| \frac{t_{km}(\Lambda_k(y) - L)}{\rho_2}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \\
 & \leq DF \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x) - L)}{\rho_1}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \\
 & + DF \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(y) - L)}{\rho_2}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k},
 \end{aligned}$$

where $F = \max \left[1, \left(\frac{|\alpha|}{(|\alpha|\rho_1 + |\beta|\rho_2)} \right)^H, \left(\frac{|\beta|}{(|\alpha|\rho_1 + |\beta|\rho_2)} \right)^H \right]$. From the above inequality, we get

$$\begin{aligned}
 & \left\{ n \in \mathbb{N} : \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(\alpha x + \beta y) - L)}{|\alpha|\rho_1 + |\beta|\rho_2}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq \epsilon \right\} \\
 & \subseteq \left\{ n \in \mathbb{N} : DF \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x) - L)}{\rho_1}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq \frac{\epsilon}{2} \right\} \\
 & \cup \left\{ n \in \mathbb{N} : DF \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(y) - L)}{\rho_2}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq \frac{\epsilon}{2} \right\}.
 \end{aligned}$$

Two sets on the right hand side belong to I and this completes the proof. Similarly, we can prove that $\hat{w}_0^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$, $\hat{w}_\infty(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$ and $\hat{w}_\infty^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$ are linear spaces.

Theorem 2.2. *Let $\mathcal{M} = (M_k)$ be a Musielak-Orlicz function, $p = (p_k)$ be a bounded sequence of positive real numbers. For any fixed $n \in \mathbb{N}$, $\hat{w}_\infty(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$ is a paranormed space with the paranorm defined by*

$$\begin{aligned}
 g(x) &= \inf \left\{ \rho^{\frac{p_n}{H}} : \rho > 0 \text{ is such that } \sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \right. \\
 & \left. \leq 1, \forall z_1, \dots, z_{n-1} \in X \right\}.
 \end{aligned}$$

Proof. It is clear that $g(x) = g(-x)$. Since $M_k(0) = 0$, we get $\inf \{ \rho^{\frac{p_n}{H}} \} = 0$ for $x = 0$ therefore, $g(0) = 0$. Let us take $x, y \in \hat{w}_\infty(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$. Let

$$B(x) = \left\{ \rho^{\frac{p_n}{H}} : \rho > 0, \sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \leq 1, \right.$$

$$\begin{aligned} & \forall z_1, \dots, z_{n-1} \in X \}, \\ B(y) = & \left\{ \rho^{\frac{p_n}{H}} : \rho > 0, : \sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(y))}{\rho} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_k} \leq 1, \right. \\ & \left. \forall z_1, \dots, z_{n-1} \in X \right\}. \end{aligned}$$

Let $\rho_1 \in B(x)$ and $\rho_2 \in B(y)$. If $\rho = \rho_1 + \rho_2$, then we have

$$\begin{aligned} & \sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x+y))}{\rho} \right\|, z_1, \dots, z_{n-1} \right) \right] \\ & \leq \frac{\rho_1}{\rho_1 + \rho_2} \sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho_1} \right\|, z_1, \dots, z_{n-1} \right) \right] \\ & \quad + \frac{\rho_2}{\rho_1 + \rho_2} \sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(y))}{\rho_2} \right\|, z_1, \dots, z_{n-1} \right) \right]. \end{aligned}$$

Thus $\sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x+y))}{\rho_1 + \rho_2} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_k} \leq 1$ and

$$\begin{aligned} g(x+y) & \leq \inf \left\{ (\rho_1 + \rho_2)^{\frac{p_n}{H}} : \rho_1 \in B(x), \rho_2 \in B(y) \right\} \\ & \leq \inf \left\{ \rho_1^{\frac{p_n}{H}} : \rho_1 \in B(x) \right\} + \inf \left\{ \rho_2^{\frac{p_n}{H}} : \rho_2 \in B(y) \right\} \\ & = g(x) + g(y). \end{aligned}$$

Let $\sigma^m \rightarrow \sigma$ where $\sigma, \sigma^m \in \mathbb{C}$ and let $g(x^m - x) \rightarrow 0$ as $m \rightarrow \infty$. We have to show that $g(\sigma^m x^m - \sigma_x) \rightarrow 0$ as $m \rightarrow \infty$. Let

$$\begin{aligned} B(x^m) = & \left\{ \rho_m^{\frac{p_n}{H}} : \rho_m > 0, \sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x^m))}{\rho_m} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_k} \leq 1, \right. \\ & \left. \forall z_1, \dots, z_{n-1} \in X \right\}, \\ B(x^m - x) = & \left\{ \rho'_m{}^{\frac{p_n}{H}} : \rho'_m > 0, \sup_k \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x^m - x))}{\rho'_m} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_k} \leq 1, \right. \\ & \left. \forall z_1, \dots, z_{n-1} \in X \right\}. \end{aligned}$$

If $\rho_m \in B(x^m)$ and $\rho'_m \in B(x^m - x)$ then we observe that

$$\begin{aligned}
 & \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\sigma^m \Lambda_k(x^m) - \sigma \Lambda_k(x))}{\rho_m |\sigma^m - \sigma| + \rho'_m |\sigma|} \right\|, z_1, \dots, z_{n-1} \right) \right] \\
 & \leq \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\sigma^m \Lambda_k(x^m) - \sigma \Lambda_k(x^m))}{\rho_m |\sigma^m - \sigma| + \rho'_m |\sigma|} \right\|, z_1, \dots, z_{n-1} \right) \right. \\
 & \quad \left. + \left\| \frac{t_{km}(\sigma \Lambda_k(x^m) - \sigma \Lambda_k(x))}{\rho_m |\sigma^m - \sigma| + \rho'_m |\sigma|} \right\|, z_1, \dots, z_{n-1} \right) \right] \\
 & \leq \frac{|\sigma^m - \sigma| \rho_m}{\rho_m |\sigma^m - \sigma| + \rho'_m |\sigma|} \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x^m))}{\rho_m} \right\|, z_1, \dots, z_{n-1} \right) \right] \\
 & \quad + \frac{|\sigma| \rho'_m}{\rho_m |\sigma^m - \sigma| + \rho'_m |\sigma|} \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x^m) - \Lambda_k(x))}{\rho'_m} \right\|, z_1, \dots, z_{n-1} \right) \right].
 \end{aligned}$$

From the above inequality, it follows that

$$\frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\sigma^m \Lambda_k(x^m) - \sigma \Lambda_k(x))}{\rho_m |\sigma^m - \sigma| + \rho'_m |\sigma|} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_k} \leq 1$$

and consequently,

$$\begin{aligned}
 g(\sigma^m x^m - \sigma x) & \leq \inf \left\{ \left(\rho_m |\sigma^m - \sigma| + \rho'_m |\sigma| \right)^{\frac{p_n}{H}} : \rho_m \in B(x^m), \rho'_m \in B(x^m - x) \right\} \\
 & \leq (|\sigma^m - \sigma|)^{\frac{p_n}{H}} \inf \left\{ \rho^{\frac{p_n}{H}} : \rho \in B(x^m) \right\} \\
 & \quad + (|\sigma|)^{\frac{p_n}{H}} \inf \left\{ (\rho'_m)^{\frac{p_n}{H}} : \rho'_m \in B(x^m - x) \right\} \longrightarrow 0 \text{ as } m \longrightarrow \infty.
 \end{aligned}$$

This completes the proof.

Theorem 2.3. *Let $\mathcal{M}, \mathcal{M}', \mathcal{M}''$ are Musielak-Orlicz functions. Then we have*
 (i) $\hat{w}_0^I(\mathcal{M}', \Lambda, p, \|\cdot, \dots, \cdot\|) \subseteq \hat{w}_0^I(\mathcal{M} \circ \mathcal{M}', \Lambda, p, \|\cdot, \dots, \cdot\|)$ provided (p_k) is such that

$$H_0 = \inf p_k > 0.$$

(ii) $\hat{w}_0^I(\mathcal{M}', \Lambda, p, \|\cdot, \dots, \cdot\|) \cap \hat{w}_0^I(\mathcal{M}'', \Lambda, p, \|\cdot, \dots, \cdot\|) \subseteq \hat{w}_0^I(\mathcal{M}' + \mathcal{M}'', \Lambda, p, \|\cdot, \dots, \cdot\|)$.

Proof. (i) For given $\epsilon > 0$, first choose $\epsilon_0 > 0$ such that $\max\{\epsilon_0^H, \epsilon_0^{H_0}\} < \epsilon$. Since (M_k) is continuous, choose $0 < \delta < 1$ such that $0 < t < \delta$, this implies that $M_k(t) < \epsilon_0$. Let $x \in \hat{w}_0^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$. Now from the definition

$$B(\delta) = \left\{ n \in \mathbb{N} : \frac{1}{n} \sum_{k=1}^n \left[M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_k} \geq \delta^H \right\} \in I.$$

Thus if $n \notin B(\delta)$ then

$$\frac{1}{n} \sum_{k=1}^n \left[M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho} \right\|, z_1, \dots, z_{n-1} \right) \right]^{p_k} < \delta^H$$

$$\begin{aligned}
&\implies \sum_{k=1}^n \left[M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} < n\delta^H \\
&\implies \left[M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} < \delta^H \text{ for all } k, m = 1, 2, 3, \dots \\
&\implies \left[M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right] < \delta \text{ for all } k, m = 1, 2, 3, \dots
\end{aligned}$$

Hence from above and using the continuity of $\mathcal{M} = (M_k)$, we have

$$\left[M_k \left(M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right) \right] < \epsilon_0 \quad \forall k, m = 1, 2, 3, \dots,$$

which consequently implies that

$$\sum_{k=1}^n \left[M_k \left(M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right) \right]^{p_k} < \max\{\epsilon_0^H, \epsilon_0^{H_0}\} < \epsilon.$$

$$\text{Thus } \frac{1}{n} \sum_{k=1}^n \left[M_k \left(M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right) \right]^{p_k} < \epsilon.$$

This shows that

$$\left\{ n \in \mathbb{N} : \frac{1}{n} \sum_{k=1}^n \left[M_k \left(M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right) \right]^{p_k} \geq \epsilon \right\} \subset B(\delta)$$

and so belongs to I . This proves the result.

(ii) Let $(x_k) \in \hat{w}_0^I(\mathcal{M}', \Lambda, p, \|\cdot, \dots, \cdot\|) \cap \hat{w}_0^I(\mathcal{M}'', \Lambda, p, \|\cdot, \dots, \cdot\|)$. Then the fact

$$\frac{1}{n} \left[(M'_k + M''_k) \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \leq$$

$$D \frac{1}{n} \left[M'_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} + D \frac{1}{n} \left[M''_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k}$$

gives the result.

Theorem 2.4. *The sequence spaces $\hat{w}_0^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$ and $\hat{w}_\infty^I(\mathcal{M}', \Lambda, p, \|\cdot, \dots, \cdot\|)$ are solid.*

Proof. Let $x \in \hat{w}_0^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$, let (α_k) be a sequence of scalars such that $|\alpha_k| \leq 1$ for all $k \in \mathbb{N}$. Then we have

$$\left\{ n \in \mathbb{N} : \frac{1}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(\alpha_k x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \right\} \subset$$

$$\left\{ n \in \mathbb{N} : \frac{C}{n} \sum_{k=1}^n \left[M_k \left(\left\| \frac{t_{km}(\Lambda_k(x))}{\rho}, z_1, \dots, z_{n-1} \right\| \right) \right]^{p_k} \geq \epsilon \right\} \in I,$$

where $C = \max\{1, |\alpha_k|^H\}$. Hence $(\alpha_k x) \in \hat{w}_0^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$ for all sequences of scalars α_k with $|\alpha_k| \leq 1$ for all $k \in \mathbb{N}$ whenever $(x_k) \in \hat{w}_0^I(\mathcal{M}, \Lambda, p, \|\cdot, \dots, \cdot\|)$. Similarly, we can prove that $\hat{w}_\infty^I(\mathcal{M}', \Lambda, p, \|\cdot, \dots, \cdot\|)$ is also solid.

References

- [1] S. Banach, *Theorie Operations Lineaires*, Chelsea Publishing Co., New York 1955.
- [2] P. Das, P. Kostyrko, W. Wilczynski and P. Malik, *I and I* convergence of double sequences*, Math. Slovaca 58 (2008) 605-620.
- [3] P. Das, P. Malik, *On the statistical and I-variation of double sequences*, Real Anal. Exchange 33 (2007-2008) 351-364.
- [4] A. Esi, *Some new sequence spaces defined by a sequence of moduli*, Turk. J. Math. 21 (1997) 61-68.
- [5] A. Esi, *Strongly $[V_2, \lambda_2, M, p]$ -summable double sequence spaces defined by Orlicz function*, Int. J. Nonlinear Anal. Appl. 2 (2011) 110-115.
- [6] S. Gähler, *Linear 2-normietre Rume*, Math. Nachr. 28 (1965), 1-43.
- [7] M. Gurdal, S. Pehlivan, *Statistical convergence in 2-normed spaces*, Southeast Asian Bull. Math. 33 (2009) 257-264.
- [8] H. Gunawan, *On n-inner product, n-norms, and the Cauchy-Schwartz inequality*, Sci. Math. Jpn. 5 (2001) 47-54.
- [9] H. Gunawan, *The space of p-summable sequence and its natural n-norm*, Bull. Aust. Math. Soc. 64 (2001) 137-147.
- [10] H. Gunawan, M. Mashadi, *On n-normed spaces*, Int. J. Math. Math. Sci. 27 (2001) 631-639.
- [11] P. Kostyrko, T. Salat and W. Wilczyński, *I-convergence*, Real Anal. Exchange 26 (2000) 669-686.
- [12] G.G. Lorentz, *A contribution to the theory of divergent series*, Acta Math. 80 (1948) 167-190.
- [13] J. Lindenstrauss, L. Tzafriri, *On Orlicz sequence spaces*, Israel J. Math. 10 (1971) 345-355.
- [14] L. Maligranda, *Orlicz Spaces and Interpolation*, Seminars in Mathematics 5, Departamento de Matemática, Universidade Estadual de Campinas, Campinas SP Brasil 1989.

- [15] A. Misiak, *n-inner product spaces*, Math. Nachr. 140 (1989) 299-319.
- [16] M. Mursaleen, *On some new invariant matrix methods of summability*, Quart. J. Math. Oxford 34 (1983) 77-86.
- [17] M. Mursaleen, A. Alotaibi, *On I-convergence in radom 2-normed spaces*, Math. Slovaca 61 (2011) 933-940.
- [18] M. Mursaleen, A.K. Noman, *On some new sequence spaces of non absolute type related to the spaces l_p and l_∞ I*, Filomat 25 (2011) 33-51.
- [19] M. Mursaleen, A.K. Noman, *On some new sequence spaces of non absolute type related to the spaces l_p and l_∞ II*, Math. Commun. 16 (2011) 383-398.
- [20] M. Mursaleen, S.A. Mohiuddine and O.H.H. Edely, *On ideal convergence of double sequences in intuitionistic fuzzy normed spaces*, Comput. Math. Appl. 59 (2010) 603-611.
- [21] M. Mursaleen, S.A. Mohiuddine, *On ideal convergence of double sequences in probabilistic normed spaces*, Math. Reports 64 (2010) 359-371.
- [22] M. Mursaleen, S.A. Mohiuddine, *On ideal convergence in probabilistic normed spaces*, Math. Slovaca 62 (2012) 49-62.
- [23] J. Musielak, *Orlicz Spaces and Modular Spaces*, Lecture Notes in Mathematics 1034, Springer-Verlag Berlin Heidelberg 1983.
- [24] K. Raj, S.K. Sharma, *Some sequence spaces in 2-normed spaces defined by Musielak-Orlicz function*, Acta Univ. Sapientiae Math. 3 (2011) 97-109.
- [25] K. Raj, S.K. Sharma, *Some generalized difference double sequence spaces defined by a sequence of Orlicz-function*, CUBO A Mathematical Journal 14 (2012) 167-190.
- [26] K. Raj, S.K. Sharma, *Some multiplier sequence spaces defined by a Musielak-Orlicz function in n-normed spaces*, New Zealand J. Math. 42 (2012) 45-56.
- [27] W. Raymond, Y. Freese and J. Cho, *Geometry of Linear 2-Normed Spaces*, N. Y. Nova Science Publishers, Huntington 2001.
- [28] A. Sahiner, M. Gurdal, S. Saltan and H. Gunawan, *Ideal convergence in 2-normed spaces*, Taiwanese J. Math. 11 (2007) 1477-1484.
- [29] B.C. Tripathy, B. Hazarika, *Some I-convergent sequence spaces defined by Orlicz functions*, Acta Mathematicae Applicatae Sinica 27 (2011) 149-154.
- [30] A. Wilansky, *Summability Through Functional Analysis*, North-Holland Math. Stud. 85, Elsevier Science Publishers B.V. 1984.

DOI: 10.7862/rf.2017.12

Sunil K. Sharma

email: sunilksharma42@yahoo.co.in

Department of Mathematics
Model Institute of Engineering & Technology
Kot Bhalwal, Jammu-181122, J&K
INDIA

Ayhan Esi

email: aesi23@hotmail.com

Department of Mathematics
Adiyaman University
02040, Adiyaman
TURKEY

Received 14.03.2017

Accepted 26.07.2017