

# A Contribution on Real and Complex Convexity in Several Complex Variables

*Jamel Abidi*

**ABSTRACT:** Let  $f, g : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic functions. Define  $u(z, w) = |w - f(z)|^4 + |w - g(z)|^4$ ,  $v(z, w) = |w - f(z)|^2 + |w - g(z)|^2$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . A comparison between the convexity of  $u$  and  $v$  is obtained under suitable conditions.

Now consider four holomorphic functions  $\varphi_1, \varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$ . We prove that  $F = |\varphi_1 - g_1|^2 + |\varphi_2 - g_2|^2$  is strictly convex on  $\mathbb{C}^n \times \mathbb{C}^m$  if and only if  $n = m = 1$  and  $\varphi_1, \varphi_2, g_1, g_2$  are affine functions with  $(\varphi_1'g_2' - \varphi_2'g_1') \neq 0$ .

Finally, it is shown that the product of four absolute values of pluriharmonic functions is plurisubharmonic if and only if the functions satisfy special conditions as well.

*AMS Subject Classification:* 32A10, 32A60, 32F32, 32U05, 32W50.

*Keywords and Phrases:* Holomorphic, convex, plurisubharmonic functions; Inequalities; Strictly; Maximum principle.

## 1. Introduction

Convex functions recently are studied in complex analysis because they appear in the theory of holomorphic functions, plurisubharmonic (psh) functions, currents, Lelong numbers, extension problems, holomorphic representation theory (see [2], [5], [6], [7], [8], [10], [11], [13], [14], [15], [16], [17] and [19]).

It is worth mentioning that an interesting relation between convex and plurisubharmonic functions has been obtained in [2].

Several papers appeared recently to this topic, let us mention [2], [3], [5], [6], [15], [19] and the monographs [11], [14], [19] and more recently [5].

Let  $n \geq 1$ . We can construct a  $C^\infty$  strictly psh function  $F$  defined on  $\mathbb{C}^n \times \mathbb{C}$ , such that  $F$  is not convex (and not concave) on each Euclidean not empty open ball subset in  $\mathbb{C}^n \times \mathbb{C}$ . For instance,

$$F(z, w) = |w - e^{\bar{z}_1}|^2 + \dots + |w - e^{\bar{z}_n}|^2, \text{ for } z = (z_1, \dots, z_n) \in \mathbb{C}^n, w \in \mathbb{C}.$$

Moreover, for the case of one complex variable, let  $\lambda(z) = 2x^2 - y^2$ ,  $z = (x + iy) \in \mathbb{C}$ ,  $x = \text{Re}(z)$ . Then  $\lambda$  is a  $C^\infty$  strictly sh function on  $\mathbb{C}$ , while  $\lambda$  is not convex (respectively not concave) at each point of  $\mathbb{C}$ .

This proves that the new class of functions, consisting of convex and strictly psh functions, is well defined because we can not compare the two families (convex functions) and (convex and strictly psh functions).

Now thanks to [2], we know the holomorphic representation of each holomorphic function  $f : \mathbb{C}^n \rightarrow \mathbb{C}$  under the suitable condition of the convexity of its modulus.

Let  $\delta \in [1, +\infty[$ . We have the following observation.

Put  $K(z, w) = |w - f(z)|^\delta$  and  $H(z, w) = |w - f(z)|$ , for  $(z, w) \in \mathbb{C}^2$ , where  $f : \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic. Assume that  $K$  is convex on  $\mathbb{C}^2$  and  $\delta > 1$ . Then  $H$  is convex on  $\mathbb{C}^2$  and we have  $H^s$  is convex on  $\mathbb{C}^2$ , for each  $s \in [1, +\infty[$  independently of  $\delta$  and conversely.

Now let  $f_1, f_2 : \mathbb{C} \rightarrow \mathbb{C}$  be two holomorphic functions and  $s \in \mathbb{N} \setminus \{0\}$ . Define  $K_{2s}(z, w) = |w - f_1(z)|^{2s} + |w - f_2(z)|^{2s}$ , for  $(z, w) \in \mathbb{C}^2$ . By theorem 10, we have that  $K_4$  is convex on  $\mathbb{C}^2$  implies that  $K_2$  is convex on  $\mathbb{C}^2$ . But the converse is not true. For instance, let  $f_1(z) = z^4$ ,  $f_2(z) = -z^4$ ,  $z \in \mathbb{C}$ . Then  $K_2$  is convex on  $\mathbb{C}^2$ . But  $K_4$  is not convex on  $\mathbb{C}^2$ . This remark leads to the following problem.

Let  $N \in \mathbb{N} \setminus \{0, 1\}$  and  $F_1, \dots, F_N : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic functions. Define

$$\psi_\delta(z, w) = |w - F_1(z)|^\delta + \dots + |w - F_N(z)|^\delta, \text{ for } (z, w) \in \mathbb{C}^n \times \mathbb{C}.$$

Suppose that  $\psi_\delta$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ .

Firstly, for the study of the convexity of  $\psi_\delta$ , we observe that we study separately the following two cases.

Case 1.  $\delta \in [1, +\infty[ \setminus \{2\}$ .

Case 2.  $\delta = 2$ .

Is it true that  $\delta \in [1, +\infty[ \setminus \{2\}$ , implies that  $F_1, \dots, F_N$  are affine functions?

Recall that for  $\delta = 2$ , there exists several cases where  $\psi_2$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ , but  $F_1, \dots, F_N$  are not affine functions.

Moreover, for  $N = 2$ , by a limiting argument and a specific holomorphic differential equation, we prove that  $\psi_1$  is convex on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $F_1$  and  $F_2$  are affine functions. Indeed,  $\psi_{2k}$  is convex on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $F_1$  and  $F_2$  are affine functions, for  $k \in \mathbb{N} \setminus \{0, 1\}$ .

The paper is organized as follows. In section 2, we shall use an elementary holomorphic differential equation in the proofs of the following two technical questions. Let  $A_1, A_2 \in \mathbb{C}$  and  $n, m \in \mathbb{N} \setminus \{0\}$ . Characterize exactly all the 3 holomorphic functions  $\varphi : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  such that  $u$  is convex (respectively convex and strictly plurisubharmonic) on  $\mathbb{C}^n \times \mathbb{C}^m$ , where

$$u(z, w) = |A_1\varphi(w) - g_1(z)|^2 + |A_2\varphi(w) - g_2(z)|^2, \text{ for } (z, w) \in \mathbb{C}^n \times \mathbb{C}^m.$$

In this case find the expressions of  $\varphi$ ,  $g_1$  and  $g_2$ .

Moreover, find all the three holomorphic functions  $\varphi : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $f_1, f_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  such that  $v$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ , where

$$v(z, w) = |A_1\varphi(w) - \overline{f_1(z)}|^2 + |A_2\varphi(w) - \overline{f_2(z)}|^2, (z, w) \in \mathbb{C}^n \times \mathbb{C}^m.$$

We prove that we have a great differences between the 2 classes of functions defined similar as  $u$  and  $v$ .

Now let  $k_1, k_2 : G \rightarrow \mathbb{C}^t$  be two holomorphic functions. Then the functions  $\|k_1 + \overline{k_2}\|^2$  and  $(\|k_1 + \lambda\|^2 + \|k_2 + \delta\|^2)$  have the same hermitian Levi form on  $G$ , where  $G$  is a domain of  $\mathbb{C}^s$ ,  $\lambda, \delta \in \mathbb{C}^t$  and  $s, t \in \mathbb{N} \setminus \{0\}$ .

For the applications, we can see the proof of theorem 4, corollary 1, theorem 5 and others.

In section 3, we consider the following problems.

Problem 1. Let  $n, m \geq 1$ . Find all the 4 holomorphic functions  $\varphi_1, \varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  such that  $\psi = |\varphi_1 - g_1|^2 + |\varphi_2 - g_2|^2$  is strictly convex on  $\mathbb{C}^n \times \mathbb{C}^m$ .

Problem 2. Characterize all the holomorphic functions  $\varphi_1, \varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  such that  $\psi = |\varphi_1 - g_1|^2 + |\varphi_2 - g_2|^2$  is convex and strictly psh (respectively convex) on  $\mathbb{C}^n \times \mathbb{C}^m$ .

Before stating it, we can study the analysis question. Find all the holomorphic functions  $\varphi_1, \varphi_2, \psi_1, \psi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $f_1, f_2, g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$ , such that  $u_1$  and  $u_2$  are convex and  $u = (u_1 + u_2)$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ . Where  $u_1(z, w) = |\varphi_1(w) - f_1(z)|^2 + |\varphi_2(w) - f_2(z)|^2$ ,  $u_2(z, w) = |\psi_1(w) - g_1(z)|^2 + |\psi_2(w) - g_2(z)|^2$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .

In section 4, we use an algebraic method to mainly focus on properties of the new structure (convex and strictly psh) and their relations with the holomorphic representation theory.

In section 5 we study the product of several absolute values of pluriharmonic (prh) functions and some auxiliary results are proved.

Let  $U$  be a domain of  $\mathbb{R}^d$ , ( $d \geq 2$ ). Put  $sh(U)$  the set of all subharmonic functions on  $U$ . For  $f : U \rightarrow \mathbb{C}$  be a function,  $|f|$  is the modulus of  $f$ . For  $N \geq 1$  and  $h = (h_1, \dots, h_N)$ , where  $h_1, \dots, h_N : U \rightarrow \mathbb{C}$ ,  $\|h\| = (|h_1|^2 + \dots + |h_N|^2)^{\frac{1}{2}}$ .

Let  $g : D \rightarrow \mathbb{C}$  be an analytic function,  $D$  is a domain of  $\mathbb{C}$ . We denote  $\frac{\partial^m g}{\partial z^m}$  the holomorphic derivative of  $g$  of order  $m$ , for all  $m \in \mathbb{N} \setminus \{0\}$ .

If  $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{C}^n$ , and  $z = (z_1, \dots, z_n) \in \mathbb{C}^n$  we write  $\langle z/\xi \rangle = z_1 \overline{\xi_1} + \dots + z_n \overline{\xi_n}$  and  $B(\xi, r) = \{z \in \mathbb{C}^n / \|z - \xi\| < r\}$  for  $r > 0$ , where  $\sqrt{\langle \xi/\xi \rangle} = \|\xi\|$  is the Euclidean norm of  $\xi$ . The Lebesgue measure on  $\mathbb{C}^n$  is denoted by  $m_{2n}$  and  $C^k(U) = \{\varphi : U \rightarrow \mathbb{C} / \varphi \text{ is a function of class } C^k \text{ on } U\}$ ,  $k \in \mathbb{N} \cup \{\infty\} \setminus \{0\}$ .

Let  $D$  be a domain of  $\mathbb{C}^n$ , ( $n \geq 1$ ). An usual psh( $D$ ) and prh( $D$ ) are respectively the classes of plurisubharmonic and pluriharmonic functions on  $D$ . For all  $a \in \mathbb{C}$ ,  $|a|$  is the modulus of  $a$ ,  $\text{Re}(a)$  is the real part of  $a$  and  $D(a, r) = \{z \in \mathbb{C} / |z - a| < r\}$  for  $r > 0$ .

For the study of properties and extension problems of analytic and plurisubharmonic functions we cite the references [1], [6], [7], [8], [9], [10], [12], [13], [15], [16] and [17]. For the study of convex functions in complex convex domains, we cite [5], [11],

[14], [2] and [19].

For the theory of  $n$ - subharmonic functions we cite [18].

## 2. A family of analytic functions and the holomorphic representation theory

We have

**Lemma 1.** *Let  $g = (g_1, \dots, g_N), f = (f_1, \dots, f_N) : D \rightarrow \mathbb{C}^N$  be two holomorphic functions,  $N \geq 1$ ,  $D$  is a domain of  $\mathbb{C}^n$ ,  $n \geq 1$  and  $a, b \in \mathbb{C}^N$ . Then*

*$\|f + \bar{g}\|^2$  and  $(\|f + a\|^2 + \|g + b\|^2)$  have the same hermitian Levi form on  $D$ .*

*On the other hand, let  $u : D \rightarrow \mathbb{R}$  be a function of class  $C^2$ . Define  $u_1 = (u + \|f + \bar{g}\|^2)$ ,  $u_2 = (u + \|f + a\|^2 + \|g + b\|^2)$ .*

*Then  $u_1$  and  $u_2$  are functions of class  $C^2$  on  $D$  and we have the assertion.*

*The function  $u_1$  is strictly psh on  $D$  if and only if  $u_2$  is strictly psh on  $D$ .*

*(Observe that if  $N < n$ , then  $\|g\|^2$  is not strictly psh at each point of  $D$ ).*

**Proof.** We have  $\|f + \bar{g}\|^2 = |f_1 + \bar{g}_1|^2 + \dots + |f_N + \bar{g}_N|^2 = |f_1|^2 + |g_1|^2 + \dots + |f_N|^2 + |g_N|^2 + \sum_{j=1}^N (g_j f_j + \bar{g}_j \bar{f}_j) = \|g\|^2 + \|f\|^2 + \sum_{j=1}^N (g_j f_j + \bar{g}_j \bar{f}_j)$ .

Since  $(g_j f_j + \bar{g}_j \bar{f}_j)$  is prh on  $D$ , then  $\sum_{j=1}^N (g_j f_j + \bar{g}_j \bar{f}_j)$  is prh on  $D$ .

Consequently,  $\|f + \bar{g}\|^2$  and  $(\|f + a\|^2 + \|g + b\|^2)$  have the same hermitian Levi form on  $D$ .

By [4], we have

**Theorem 1.** *Let  $\varphi : \mathbb{C}^m \rightarrow \mathbb{C}$  be a holomorphic nonconstant function,  $m \geq 1$ . Given  $A_1, A_2 \in \mathbb{C} \setminus \{0\}$  and  $n \geq 1$ .*

*The following conditions are equivalent*

*(I) There exists 2 holomorphic functions  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  such that  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}^m$ ,  $u(z, w) = |A_1 \varphi(w) - g_1(z)|^2 + |A_2 \varphi(w) - g_2(z)|^2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ ;*

*(II) There exists  $c \in \mathbb{C}$  such that  $|\varphi + c|^2$  is convex on  $\mathbb{C}^m$ .*

Now in all of this section,  $(A_1, A_2) \in \mathbb{C}^2$ . Let  $\varphi : \mathbb{C}^m \rightarrow \mathbb{C}$  be a holomorphic nonconstant function,  $m \geq 1$ . Let  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be 2 holomorphic functions,  $n \geq 1$ . Define  $u(z, w) = |A_1 \varphi(w) - g_1(z)|^2 + |A_2 \varphi(w) - g_2(z)|^2$ ,  $u_1(z, w) = |A_1 \varphi(w) - \bar{g}_1(z)|^2 + |A_2 \varphi(w) - \bar{g}_2(z)|^2$ ,  $u_2 = u + u_1$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .  $v(z, w) = |A_1 \bar{\varphi}(w) - g_1(z)|^2 + |A_2 \bar{\varphi}(w) - g_2(z)|^2$ ,  $v_1(z, w) = |A_1 \bar{\varphi}(w) - \bar{g}_1(z)|^2 + |A_2 \bar{\varphi}(w) - \bar{g}_2(z)|^2$  and  $v_2 = v + v_1$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ . We have

**Theorem 2.** *Assume that  $(A_1, A_2) \in \mathbb{C}^2 \setminus \{0\}$ . The following conditions are equivalent*

*(I)  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}^m$ ;*

*(II)  $\varphi$  is an affine function on  $\mathbb{C}^m$ , or  $\varphi$  is not affine and there exists  $c \in \mathbb{C}$  such that  $|\varphi + c|^2$  is convex on  $\mathbb{C}^m$  and we have the following cases.*

*Case 1. The function  $\varphi$  is affine on  $\mathbb{C}^m$ .*

Then we have the representation

$$\begin{cases} g_1(z) = A_1(\langle z/\lambda_1 \rangle + \mu_1) + \overline{A_2}\varphi_1(z) \\ g_2(z) = A_2(\langle z/\lambda_1 \rangle + \mu_1) - \overline{A_1}\varphi_1(z) \end{cases}$$

for each  $z \in \mathbb{C}^n$ , where  $\lambda_1 \in \mathbb{C}^n$ ,  $\mu_1 \in \mathbb{C}$ ,  $\varphi_1 : \mathbb{C}^n \rightarrow \mathbb{C}$  is analytic,  $|\varphi_1|^2$  is convex on  $\mathbb{C}^n$ .

Case 2.  $\varphi$  is not affine on  $\mathbb{C}^n$ .

In this case there exists  $c \in \mathbb{C}$  such that  $|\varphi + c|^2$  is convex on  $\mathbb{C}^m$ . Then we have the representation

$$\begin{cases} g_1(z) = A_1c + \overline{A_2}\varphi_1(z) \\ g_2(z) = A_2c - \overline{A_1}\varphi_1(z) \end{cases}$$

for every  $z \in \mathbb{C}^n$ , where  $\varphi_1 : \mathbb{C}^n \rightarrow \mathbb{C}$  is analytic,  $|\varphi_1|^2$  is convex on  $\mathbb{C}^n$ .

We can discuss the cases  $(A_1, A_2 \in \mathbb{C} \setminus \{0\})$ , or  $(A_1 \in \mathbb{C} \setminus \{0\}, A_2 = 0)$ , or  $(A_1 = 0, A_2 \in \mathbb{C} \setminus \{0\})$ .

This theorem motivates the following questions. Find all the holomorphic representation of the analytic functions  $f_1, f_2, f_3 : \mathbb{C}^n \rightarrow \mathbb{C}$ , such that  $\psi$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ .  $\psi(z, w) = |B_1w - f_1(z)|^2 + |B_2w - f_2(z)|^2 + |B_3w - f_3(z)|^2$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ , where  $(B_1, B_2, B_3) \in \mathbb{C}^3 \setminus \{0\}$ .

Indeed, for instance, in harmonic analysis and convex analysis, actually the following question appeared naturally.

Find all the representation of the harmonic functions  $F_1, F_2, F_3 : \mathbb{C} \rightarrow \mathbb{C}$ , such that  $\psi_1$  is convex and strictly 2-sh on  $\mathbb{C}^2$ . Where  $\psi_1(z, w) = |w - F_1(z)|^2 + |w - F_2(z)|^2 + |w - F_3(z)|^2$ ,  $(z, w) \in \mathbb{C}^2$ . (We study here functions on harmonic representation theory). Define  $\psi_0(z, w) = |w - F_1(z)|^2 + |w - F_2(z)|^2$ , for  $(z, w) \in \mathbb{C}^2$ . If we choose  $F_3$  is affine on  $\mathbb{C}$  and  $\psi_0$  is convex and strictly 2-sh on  $\mathbb{C}^2$ , then we have a family of harmonic functions which satisfy the above condition.

The proof of this theorem is obvious and analogous to the proof of the following.

**Theorem 3.** *The following conditions are equivalent*

(I)  $u$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ ;

(II)  $(A_1, A_2) \in \mathbb{C}^2 \setminus \{0\}$ ,  $n = m = 1$ , there exists  $c \in \mathbb{C}$  such that  $|\varphi + c|^2$  is convex on  $\mathbb{C}$  and we have the following cases.

Case 1.  $A_1A_2 \neq 0$ . Then

$$\begin{cases} g_1(z) + A_1c = A_1(az + b) + \overline{A_2}\psi(z) \\ g_2(z) + A_2c = A_2(az + b) - \overline{A_1}\psi(z) \end{cases}$$

for each  $z \in \mathbb{C}$ , where  $a, b \in \mathbb{C}$ ,  $\psi : \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic,  $|\psi|$  is convex with  $|\psi'| > 0$  and  $|\varphi'| > 0$  on  $\mathbb{C}$ .

Case 2.  $A_1 \neq 0$  and  $A_2 = 0$ .

If  $\varphi$  is affine and nonconstant on  $\mathbb{C}$ . Then we have the representation

$$\begin{cases} g_1(z) = A_1(\lambda z + \mu) \\ g_2(z) = -\overline{A_1}\varphi_2(z) \end{cases}$$

for each  $z \in \mathbb{C}$ , where  $\lambda \in \mathbb{C} \setminus \{0\}$ ,  $\mu \in \mathbb{C}$ ,  $\varphi_2 : \mathbb{C} \rightarrow \mathbb{C}$  is analytic,  $|\varphi_2|^2$  is convex and strictly subharmonic (sh) on  $\mathbb{C}$ .

If  $\varphi$  is not affine on  $\mathbb{C}$ . Then we have the representation

$$\begin{cases} g_1(z) = -A_1 c \\ g_2(z) = -\overline{A_1} \varphi_3(z) \end{cases}$$

for every  $z \in \mathbb{C}$ , where  $\varphi_3 : \mathbb{C} \rightarrow \mathbb{C}$  is analytic,  $|\varphi_3|^2$  is convex and strictly subharmonic on  $\mathbb{C}$ . In this situation we have  $\varphi(w) = e^{(aw+b)} - c$ , for each  $w \in \mathbb{C}$ , with  $a \in \mathbb{C} \setminus \{0\}$  and  $b \in \mathbb{C}$ .

Case 3.  $A_1 = 0$  and  $A_2 \neq 0$ . (Obviously analogous to case 2).

**Proof.** (I) implies (II). We choose the following proof which have technical applications in the case when we study the convexity of the function  $F$ ,  $F(z, w) = |w - \psi_1(z)|^{2N} + |w - \psi_2(z)|^{2N}$ ,  $N \in \mathbb{N}$ ,  $N \geq 2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ ,  $\psi_1, \psi_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be two holomorphic functions. In this situation we prove that  $\psi_1$  and  $\psi_2$  have analytic representations using the holomorphic differential equation  $k''(k + \lambda) = \gamma(k')^2$ , where  $k : \mathbb{C} \rightarrow \mathbb{C}$  is a holomorphic function and  $\lambda, \gamma \in \mathbb{C}$ .

If  $(A_1, A_2) = (0, 0)$ , then  $u$  is independent of  $w$ . Thus  $u$  is not strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ . A contradiction.

The case where  $A_1 \neq 0$  and  $A_2 = 0$ .

Since  $u(0, \cdot)$  is strictly psh on  $\mathbb{C}^m$ . Then the function  $|A_1 \varphi - g_1(0)|^2$  is strictly psh on  $\mathbb{C}^m$ . Thus by lemma 1,  $m = 1$ . Since  $u(\cdot, 0)$  is convex on  $\mathbb{C}$ , then  $|\varphi - \frac{g_1(0)}{A_1}|^2$  is convex and strictly sh on  $\mathbb{C}$ . Put  $c = -\frac{g_1(0)}{A_1}$ . Now  $|\varphi + c|^2$  is convex and strictly sh on  $\mathbb{C}$ , therefore, by Abidi [2], we have

$\varphi(w) = aw + b$ , for all  $w \in \mathbb{C}$ , where  $a \in \mathbb{C} \setminus \{0\}$  and  $b \in \mathbb{C}$ , or  
 $\varphi(w) = e^{(a_1 w + b_1)} - c$ , for all  $w \in \mathbb{C}$ , with  $a_1 \in \mathbb{C} \setminus \{0\}$  and  $b_1 \in \mathbb{C}$ .

If  $\varphi(w) = aw + b$ ,  $\forall w \in \mathbb{C}$ .

Then for each fixed  $w_0 \in \mathbb{C}$ , the function  $u(\cdot, w_0)$  is convex on  $\mathbb{C}^n$ .

Therefore,

$$\begin{aligned} & \left| - \sum_{j,k=1}^n \frac{\partial^2 g_1}{\partial z_j \partial z_k}(z) [\overline{A_1} (\overline{aw_0 + b}) - \overline{g_1}(z)] \alpha_j \alpha_k + \sum_{j,k=1}^n \frac{\partial^2 g_2}{\partial z_j \partial z_k}(z) \overline{g_2}(z) \alpha_j \alpha_k \right| \\ & \leq \left| \sum_{j=1}^n \frac{\partial g_1}{\partial z_j}(z) \alpha_j \right|^2 + \left| \sum_{j=1}^n \frac{\partial g_2}{\partial z_j}(z) \alpha_j \right|^2, \end{aligned}$$

for each  $z \in \mathbb{C}^n$ ,  $w_0 \in \mathbb{C}$  and  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{C}^n$ .

Since the right hand side of the above inequality is independent of  $w_0 \in \mathbb{C}$ , it follows that for every fixed  $z \in \mathbb{C}^n$ ,

$$\sum_{j,k=1}^n \frac{\partial^2 g_1}{\partial z_j \partial z_k}(z) \alpha_j \alpha_k = 0, \text{ for all } \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{C}^n.$$

Therefore  $g_1$  is affine on  $\mathbb{C}^n$ .

Put  $g_1(z) = A_1 \langle z/\gamma \rangle + \delta$ , for  $z \in \mathbb{C}^n$ , where  $\gamma \in \mathbb{C}^n$  and  $\delta \in \mathbb{C}$ .

Let  $T : \mathbb{C}^n \times \mathbb{C} \rightarrow \mathbb{C}^n \times \mathbb{C}$ ,  $T(z, w) = (z, w + \frac{g_1(z)}{A_1 a} - \frac{\delta}{a})$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .

Note that  $T$  is a  $\mathbb{C}$  linear bijective transformation on  $\mathbb{C}^n \times \mathbb{C}$ .

Since  $u$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , then  $\psi$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , where  $\psi(z, w) = u \circ T(z, w) = |A_1(aw + b - \delta)|^2 + |g_2(z)|^2$ , for every  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .

But  $\psi$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , then  $|g_2|^2$  is convex and strictly psh on  $\mathbb{C}^n$ . Thus  $n = 1$ .

Put  $g_2(z) = -\overline{A_1}\varphi_2(z)$ , for  $z \in \mathbb{C}$  ( $\varphi_2$  is analytic on  $\mathbb{C}$ ). Thus  $|\varphi_2|^2$  is convex and strictly sh on  $\mathbb{C}$ .

(II) implies (I). Obvious.

Question. Let  $B_1, B_2 \in \mathbb{C} \setminus \{0\}$ . For  $f_1, f_2 : \mathbb{C}^n \rightarrow \mathbb{C}$ , define  $\psi(z, w) = |B_1 w - f_1(z)|^2 + |B_2 w - f_2(z)|^2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . Find all the pluriharmonic (respectively  $n$ -harmonic) functions  $f_1, f_2 : \mathbb{C}^n \rightarrow \mathbb{C}$ , such that  $\psi$  is convex (respectively convex and strictly  $n$ -subharmonic) on  $\mathbb{C}^n \times \mathbb{C}$ .

**Theorem 4.** *The following conditions are equivalent*

(I)  $u_1$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ ;

(II)  $m = 1$ ,  $n \in \{1, 2\}$ ,  $(A_1, A_2) \in \mathbb{C}^2 \setminus \{0\}$ , there exists  $c \in \mathbb{C}$  such that  $|\varphi + c|^2$  is convex on  $\mathbb{C}$  and we have the following cases.

Case 1. For all  $w \in \mathbb{C}$ ,  $\varphi(w) = aw + b$ , where  $a \in \mathbb{C} \setminus \{0\}$  and  $b \in \mathbb{C}$ .

We have the representation

$$\begin{cases} g_1(z) = \overline{A_1}(\langle z/\lambda_1 \rangle + \mu_1) + A_2\varphi_1(z) \\ g_2(z) = \overline{A_2}(\langle z/\lambda_1 \rangle + \mu_1) - A_1\varphi_1(z) \end{cases}$$

for each  $z \in \mathbb{C}^n$ , where  $\lambda_1 \in \mathbb{C}^n$ ,  $\mu_1 \in \mathbb{C}$ ,  $\varphi_1 : \mathbb{C}^n \rightarrow \mathbb{C}$  is analytic,  $|\varphi_1|^2$  is convex on  $\mathbb{C}^n$ , such that

( $n = 1, \lambda_1 \neq 0$ ), or

( $n = 1, \lambda_1 = 0, \frac{\partial \varphi_1}{\partial z}(z) \neq 0$ , for each  $z \in \mathbb{C}$ ), or

$n = 2, (\lambda_1, (\frac{\partial \varphi_1}{\partial z_1}(z), \frac{\partial \varphi_1}{\partial z_2}(z)))$  is a basis of the complex vector space  $\mathbb{C}^2$ ,

for each  $z = (z_1, z_2) \in \mathbb{C}^2$ .

Case 2. For every  $w \in \mathbb{C}$ ,  $\varphi(w) = e^{(aw+b)} - c$ , where  $a \in \mathbb{C} \setminus \{0\}$  and  $b \in \mathbb{C}$ .

Then  $n = 1$  and we have the representation

$$\begin{cases} g_1(z) = -\overline{A_1}\bar{c} + A_2\psi_1(z) \\ g_2(z) = -\overline{A_2}\bar{c} - A_1\psi_1(z) \end{cases}$$

for each  $z \in \mathbb{C}$ , where  $\psi_1 : \mathbb{C} \rightarrow \mathbb{C}$  is analytic,  $|\psi_1|^2$  is convex and strictly sh on  $\mathbb{C}$ .

The proof follows from the above 3 theorems and lemma 1.

We have

**Corollary 1.** *The following conditions are equivalent*

(I)  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}^m$  and  $u_2$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ ;

(II)  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}^m$  and  $u_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ ;

(III)  $(A_1, A_2) \in \mathbb{C}^2 \setminus \{0\}$ ,  $m = 1$ ,  $n \in \{1, 2\}$ , there exists  $c \in \mathbb{C}$  such that  $|\varphi + c|^2$  is convex and strictly sh on  $\mathbb{C}$  and we have the following 2 cases.

Case 1. For all  $w \in \mathbb{C}$   $\varphi(w) = aw + b$ , ( $a \in \mathbb{C} \setminus \{0\}, b \in \mathbb{C}$ ).

Then we have the holomorphic representation

$$\begin{cases} g_1(z) = A_1(\langle z/\lambda_1 \rangle + \mu_1) + \overline{A_2}\varphi_1(z) \\ g_2(z) = A_2(\langle z/\lambda_1 \rangle + \mu_1) - \overline{A_1}\varphi_1(z) \end{cases}$$

for each  $z \in \mathbb{C}^n$ , where  $\lambda_1 \in \mathbb{C}^n$ ,  $\mu_1 \in \mathbb{C}$ ,  $\varphi_1 : \mathbb{C}^n \rightarrow \mathbb{C}$  is analytic,  $|\varphi_1|^2$  is convex on  $\mathbb{C}^n$ , such that

( $n = 1, \lambda_1 \neq 0$ ), or ( $n = 1, \lambda_1 = 0, \frac{\partial \varphi_1}{\partial z}(z) \neq 0$ , for each  $z \in \mathbb{C}$ ), or

( $n = 2$  and  $(\lambda_1, (\frac{\partial \varphi_1}{\partial z_1}(z), \frac{\partial \varphi_1}{\partial z_2}(z)))$  is a basis of the complex vector space  $\mathbb{C}^2$ ,

for every  $z = (z_1, z_2) \in \mathbb{C}^2$ ).

Case 2. For all  $w \in \mathbb{C}$ ,  $\varphi(w) = e^{(aw+b)} - c$ , where  $a \in \mathbb{C} \setminus \{0\}$  and  $b \in \mathbb{C}$ .

Then  $n = 1$  and we have the holomorphic representation

$$\begin{cases} g_1(z) = -A_1c + \overline{A_2}\psi_1(z) \\ g_2(z) = -A_2c - \overline{A_1}\psi_1(z) \end{cases}$$

for every  $z \in \mathbb{C}$ , where  $\psi_1 : \mathbb{C} \rightarrow \mathbb{C}$  is analytic,  $|\psi_1|^2$  is convex and strictly sh on  $\mathbb{C}$ .

**Proof.** (I) implies (III). Note that  $u, u_1$  and  $u_2$  are functions of class  $C^\infty$  on  $\mathbb{C}^n \times \mathbb{C}^m$ . We have

$u_2$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$  if and only if  $u_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ .

Assume that  $(A_1, A_2) = (0, 0)$ . Then  $u_1$  is independent of  $w \in \mathbb{C}^m$  and  $u_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ . A contradiction.

Consequently,  $(A_1, A_2) \in \mathbb{C}^2 \setminus \{0\}$ .

Define  $u_3(z, w) = (|A_1|^2 + |A_2|^2)|\varphi(w)|^2 + |g_1(z)|^2 + |g_2(z)|^2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .

Then  $u_3$  is a function of class  $C^\infty$  on  $\mathbb{C}^n \times \mathbb{C}^m$ . But  $u_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$  if and only if  $u_3$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ .

By lemma 1, we have  $m = 1$  and  $n \leq 2$ .

Now  $u(0, \cdot)$  is convex on  $\mathbb{C}$  and  $u_3(0, \cdot)$  is strictly sh on  $\mathbb{C}$ . In fact  $(|A_1\varphi - g_1(0)|^2 + |A_2\varphi - g_2(0)|^2)$  is convex on  $\mathbb{C}$  and  $((|A_1|^2 + |A_2|^2)|\varphi|^2 + |g_1(0)|^2 + |g_2(0)|^2)$  is strictly sh on  $\mathbb{C}$ . Then there exists  $c \in \mathbb{C}$  such that  $|\varphi + c|^2$  is convex on  $\mathbb{C}$  and  $|\varphi|^2$  is strictly sh on  $\mathbb{C}$ . Which yields  $|\varphi + c|^2$  is convex and strictly sh on  $\mathbb{C}$ .

By Abidi [2], using the holomorphic differential equation  $k''(k+c) = \gamma(k')^2$  ( $k : \mathbb{C} \rightarrow \mathbb{C}$  be a holomorphic function,  $\gamma, c \in \mathbb{C}$ ), we have

$\varphi(w) = aw + b$ , for all  $w \in \mathbb{C}$ , where  $a \in \mathbb{C} \setminus \{0\}$  and  $b \in \mathbb{C}$ , or

$\varphi(w) = e^{(a_1w+b_1)} - c$ , for all  $w \in \mathbb{C}$ , with  $a_1 \in \mathbb{C} \setminus \{0\}$  and  $b_1 \in \mathbb{C}$ .

The rest of the proof is now obvious.

**Theorem 5.** The following conditions are equivalent

(I)  $v$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ ;

(II)  $m = 1$ ,  $n \in \{1, 2\}$ ,  $(A_1, A_2) \in \mathbb{C}^2 \setminus \{0\}$ , there exists  $c \in \mathbb{C}$  such that  $|\varphi + c|^2$  is convex and strictly sh on  $\mathbb{C}$  and we have the following 2 cases.

Case 1. For all  $w \in \mathbb{C}$ ,  $\varphi(w) = aw + b$ , ( $a \in \mathbb{C} \setminus \{0\}, b \in \mathbb{C}$ ).

Then we have the representation

$$\begin{cases} g_1(z) = \overline{A_1}(\langle z/\lambda \rangle + \mu) + A_2\varphi_1(z) \\ g_2(z) = \overline{A_2}(\langle z/\lambda \rangle + \mu) - A_1\varphi_1(z) \end{cases}$$



for each  $z \in \mathbb{C}^n$ , where  $\lambda \in \mathbb{C}^n$ ,  $\mu \in \mathbb{C}$ ,  $\varphi_1 : \mathbb{C}^n \rightarrow \mathbb{C}$  is analytic,  $|\varphi_1|^2$  is convex on  $\mathbb{C}^n$ , such that

( $n = 1, \lambda \neq 0$ ), or ( $n = 1, \lambda = 0, \frac{\partial \varphi_1}{\partial z}(z) \neq 0$ , for every  $z \in \mathbb{C}$ ), or

( $n = 2$ , and  $(\lambda, (\frac{\partial \varphi_1}{\partial z_1}(z), \frac{\partial \varphi_1}{\partial z_2}(z)))$  is a basis of the complex vector space  $\mathbb{C}^2$ ,

for any  $z = (z_1, z_2) \in \mathbb{C}^2$ ).

Case 2. For each  $w \in \mathbb{C}$ ,  $\varphi(w) = e^{(aw+b)} - c$ , ( $a \in \mathbb{C} \setminus \{0\}$  and  $b \in \mathbb{C}$ ).

Then  $n = 1$  and we have the representation

$$\begin{cases} g_1(z) = -\overline{A_1}c + A_2\psi_1(z) \\ g_2(z) = -A_2\overline{c} - A_1\psi_1(z) \end{cases}$$

for every  $z \in \mathbb{C}$ , where  $\psi_1 : \mathbb{C} \rightarrow \mathbb{C}$  is analytic,  $|\psi_1|^2$  is convex and strictly sh on  $\mathbb{C}$ .

Moreover, we can consider the function  $v_2$  for a study. According to lemma 1, we obtain several holomorphic representations of  $g_1$  and  $g_2$  from the assumptions  $v$  and  $v_1$  are convex on  $\mathbb{C}^n \times \mathbb{C}^m$  and  $v_2 = (v + v_1)$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ .

### 3. Some study in the theory of convex and strictly psh functions

#### 3.1. The analysis of strictly convex functions

Put  $u(z, w) = |\varphi_1(w) - g_1(z)|^2 + |\varphi_2(w) - g_2(z)|^2$ ,  $\varphi_1, \varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be four holomorphic functions,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .

Recall that, for two holomorphic functions  $\varphi : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g : \mathbb{C}^n \rightarrow \mathbb{C}$ , if we denote  $\psi(z, w) = |\varphi(w) - g(z)|^2$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .  $\psi$  is not strictly convex at each point of  $\mathbb{C}^n \times \mathbb{C}^m$  (this is the case of one absolute value of a holomorphic function). But, if we consider the sum of two absolute values of holomorphic functions, there exists several cases where  $\psi_1$  is strictly convex on  $\mathbb{C}^2$ . For example

$$\psi_1(z, w) = |f_1(w) - k_1(z)|^2 + |f_2(w) - k_2(z)|^2$$

for  $(z, w) \in \mathbb{C}^2$  and  $f_1(w) = w$ ,  $f_2(w) = 2w + 1$ ,  $k_1(z) = 2z$ ,  $k_2(z) = 0$ .

Before the two above technical remarks, we pose the following question.

Question. Characterize all the holomorphic functions  $\varphi_1, \varphi_2, g_1, g_2$  such that  $u$  is strictly convex on  $\mathbb{C}^n \times \mathbb{C}^m$  (we prove that  $n = m = 1$ ).

**Remark 1.** Let  $F_1(z) = z^2$ ,  $F_2(z) = -z^2$ ,  $F_3(z) = z$ ,  $K_1(w) = K_2(w) = K_3(w) = w$ ,  $(z, w) \in \mathbb{C}^2$ .  $F_1, F_2, F_3, K_1, K_2, K_3$  are holomorphic functions on  $\mathbb{C}$ . Put  $u(z, w) = |K_1(w) - F_1(z)|^2 + |K_2(w) - F_2(z)|^2 + |K_3(w) - F_3(z)|^2$ . Observe that  $u$  is strictly convex on  $\mathbb{C}^2$ , but  $F_1$  and  $F_2$  are not affine functions.

We begin by

**Lemma 2.** Let  $f_1, f_2 : \mathbb{C}^N \rightarrow \mathbb{C}$  be two holomorphic functions,  $N \geq 1$ . Put  $v = |f_1|^2 + |f_2|^2$ . We have

If  $v$  is strictly psh on  $\mathbb{C}^N$ , then  $N \leq 2$ .

Using the holomorphic differential equation  $k''(k+c) = \gamma(k')^2$ , for  $k : \mathbb{C} \rightarrow \mathbb{C}$  be a holomorphic function and  $(\gamma, c) \in \mathbb{C}^2$ , we have

**Lemma 3.** *Let  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  and  $\varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  be three holomorphic functions and  $a \in \mathbb{C}$ .*

*Put  $u(z, w) = |g_1(z) - a|^2 + |\varphi_2(w) - g_2(z)|^2$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .*

*Then  $u$  is strictly convex on  $\mathbb{C}^n \times \mathbb{C}^m$  if and only if  $n = m = 1$ ,  $g_1$  is affine nonconstant,  $g_2$  is affine and  $\varphi_2$  is affine nonconstant on  $\mathbb{C}$ .*

**Proof.** Assume that  $u$  is strictly convex on  $\mathbb{C}^n \times \mathbb{C}^m$ . By lemma 2, it follows that  $n = m = 1$ . We have

$$|\varphi_2''(w)(\overline{\varphi_2}(w) - \overline{g_2}(z))| < |\varphi_2'(w)|^2$$

for each  $w \in \mathbb{C}$  and for every fixed  $z \in \mathbb{C}$ .

Put  $\psi_2(w) = |\varphi_2(w) - g_2(z)|^2$ , for  $w \in \mathbb{C}$ . By Abidi [2], for each fixed  $z \in \mathbb{C}$ , the function  $\psi_2$  is strictly convex in  $\mathbb{C}$ . Then  $\varphi_2$  is affine nonconstant on  $\mathbb{C}$ , (see [2], [3]). Now we have the inequality

$$|g_2''(z)(\overline{g_2}(z) - \overline{\varphi_2}(w)) + g_1''(z)\overline{g_1}(z)| < |g_1'(z)|^2 + |g_2'(z)|^2$$

for each  $(z, w) \in \mathbb{C}^2$ . Therefore the function  $F(w) = \overline{g_2''}(z)\varphi_2(w)$  is holomorphic and bounded on  $\mathbb{C}$ , for every fixed  $z \in \mathbb{C}$ . Therefore  $F$  is constant on  $\mathbb{C}$ , for each fixed  $z \in \mathbb{C}$ .

Since  $\varphi_2$  is affine nonconstant, it follows that  $g_2'' = 0$  on  $\mathbb{C}$ . Then  $g_2$  is affine on  $\mathbb{C}$ .

Now write  $\varphi_2(w) = A_2w + B_2$ ,  $g_2(z) = a_2z + b_2$ ,  $A_2 \in \mathbb{C} \setminus \{0\}$ ,  $B_2, a_2, b_2 \in \mathbb{C}$ . Let  $T(z, w) = (z, w + \frac{a_2}{A_2}z + \frac{b_2}{A_2})$ .

Thus  $T$  is an affine holomorphic transformation and bijective on  $\mathbb{C}^2$ . Then  $u_1 = u \circ T$  is strictly convex on  $\mathbb{C}^2$  and  $u \circ T(z, w) = |g_1(z) - a|^2 + |\varphi_2(w)|^2 = u_1(z, w)$ .

Consequently,  $g_1$  is affine nonconstant on  $\mathbb{C}$ .

The converse is obvious and the proof is complete.

Now let  $\psi_1, \psi_2, f_1, f_2, k : \mathbb{C} \rightarrow \mathbb{C}$  be holomorphic functions and  $\gamma, c \in \mathbb{C}$ . Using the holomorphic differential equation  $k''(k+c) = \gamma(k')^2$  and the two partial differential equations  $\psi_1''(w)\overline{f_1}'(z) + \psi_2''(w)\overline{f_2}'(z) = 0$ ,  $f_1''(z)\overline{\psi_1}'(w) + f_2''(z)\overline{\psi_2}'(w) = 0$  on  $\mathbb{C}^2$ , we prove

**Theorem 6.** *Let  $\varphi_1, \varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be four holomorphic functions. Put  $u(z, w) = |\varphi_1(w) - g_1(z)|^2 + |\varphi_2(w) - g_2(z)|^2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .*

*The following assertions are equivalent*

*(I)  $u$  is strictly convex on  $\mathbb{C}^n \times \mathbb{C}^m$ ;*

*(II)  $n = m = 1$ ,  $g_1, g_2, \varphi_1, \varphi_2$  are affine functions on  $\mathbb{C}$  and satisfying the condition  $(g_1'\varphi_2' - g_2'\varphi_1') \neq 0$ .*

**Proof.** We have  $n = m = 1$ , because  $u$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ . Since  $u$  is strictly convex on  $\mathbb{C}^n \times \mathbb{C}^m$ , then the function  $u(z, \cdot)$  is strictly convex on  $\mathbb{C}$ , for each  $z \in \mathbb{C}$ . Therefore,

$$|\varphi_1''(w)(\overline{\varphi_1}(w) - \overline{g_1}(z)) + \varphi_2''(w)(\overline{\varphi_2}(w) - \overline{g_2}(z))| < |\varphi_1'(w)|^2 + |\varphi_2'(w)|^2$$

for each  $w \in \mathbb{C}^m$  and for every fixed  $z \in \mathbb{C}^n$ . Thus, for every fixed  $w \in \mathbb{C}$ , the holomorphic function on the variable  $z$ , defined by  $F(z) = (g_1(z)\varphi_1''(w) + g_2(z)\overline{\varphi_2''(w)})$  is bounded on  $\mathbb{C}$ .

By Liouville theorem,  $F$  is constant on  $\mathbb{C}$ . Thus  $(g_1'(z)\overline{\varphi_1''(w)} + g_2'(z)\overline{\varphi_2''(w)}) = 0$ , for every  $z, w \in \mathbb{C}$ .

We discuss the cases  $\varphi_1'' \neq 0$  or  $\varphi_2'' \neq 0$  on  $\mathbb{C}$ . (Also we have  $(\varphi_1'(w)\overline{g_1''(z)} + \varphi_2'(w)\overline{g_2''(z)}) = 0$  on  $\mathbb{C}^2$ ).

Assume that  $\varphi_1'' \neq 0$  and  $\varphi_2'' \neq 0$ . Therefore

$$\frac{\varphi_1''(w)}{\varphi_2''(w)} = -\frac{\overline{g_2''(z)}}{g_1''(z)} = R, \quad R \in \mathbb{C}.$$

Thus,  $\varphi_1''(w) = R\varphi_2''(w)$  and  $g_2'(z) = -\overline{R}g_1'(z)$ , for each  $z, w \in \mathbb{C}$ . It follows that  $\varphi_1(w) = R\varphi_2(w) + aw + b$  and  $g_2(z) = -\overline{R}g_1(z) + \lambda$ ,  $a, b, \lambda \in \mathbb{C}$ .

The function  $F_1$  is strictly convex on  $\mathbb{C}^2$ , where

$$F_1(z, w) = |R\varphi_2(w) + aw + b - g_1(z)|^2 + |\varphi_2(w) + \overline{R}g_1(z) - \lambda|^2.$$

This proves  $|g_1 + \xi_1|^2$  is strictly convex on  $\mathbb{C}$ , where  $\xi_1 \in \mathbb{C}$ .

By the holomorphic differential equation  $k''(k + c) = \gamma(k')^2$ , ( $k : \mathbb{C} \rightarrow \mathbb{C}$  be a holomorphic function and  $c, \gamma \in \mathbb{C}$ ), we have  $g_1$  is affine nonconstant on  $\mathbb{C}$ . Therefore,  $|g_1 - \varphi_1|^2 + |\overline{R}g_1 - (\lambda - \varphi_2)|^2$  is strictly convex on  $\mathbb{C}^2$ .

By theorem 2,  $\varphi_1$  and  $\varphi_2$  are affine functions. A contradiction.

Consequently,  $\varphi_1'' = 0$ , or  $\varphi_2'' = 0$  on  $\mathbb{C}$ .

Assume that  $\varphi_1'' \neq 0$  and  $\varphi_2'' = 0$  on  $\mathbb{C}$ . Therefore  $\varphi_1''\overline{g_1} = 0$  on  $\mathbb{C}$ . Thus  $g_1' = 0$  on  $\mathbb{C}$  and then  $g_1$  is constant on  $\mathbb{C}$ . We have  $|\varphi_1 - g_1(0)|^2 + |\varphi_2 - g_2|^2$  is strictly convex on  $\mathbb{C}^2$ . By lemma 3, we have  $\varphi_1$  and  $g_2$  are affine nonconstant,  $\varphi_2$  is affine on  $\mathbb{C}$ .

Therefore  $\varphi_1$  is affine nonconstant on  $\mathbb{C}$ . A contradiction.

Consequently,  $\varphi_1$  and  $\varphi_2$  are affine functions on  $\mathbb{C}$ .

Now since the function  $u(\cdot, w)$  is strictly convex on  $\mathbb{C}$  (for each fixed  $w \in \mathbb{C}$ ), then  $g_1, g_2, \varphi_1$  and  $\varphi_2$  satisfy the partial differential equation  $g_1''\overline{\varphi_1} + g_2''\overline{\varphi_2} = 0$  in  $\mathbb{C}^2$ .

Using the last above partial differential equation, we prove that  $g_1$  and  $g_2$  are affine functions on  $\mathbb{C}$ . Note that if  $\varphi_1$  and  $g_1$  are constant functions, then  $|g_2 - \varphi_2|^2$  is strictly convex on  $\mathbb{C}^2$ . This is impossible.

Therefore, we have

( $\varphi_1$  or  $g_1$  is non constant) and ( $\varphi_2$  or  $g_2$  is non constant).

Analogously,

( $g_1$  or  $g_2$  is non constant) and ( $\varphi_1$  or  $\varphi_2$  is non constant).

Since now  $u$  is strictly convex on  $\mathbb{C}^2$ , then

$$|\varphi_1'(w)\beta - g_1'(z)\alpha|^2 + |\varphi_2'(w)\beta - g_2'(z)\alpha|^2 > 0$$

for each  $(z, w) \in \mathbb{C}^2$  and  $(\alpha, \beta) \in \mathbb{C}^2 \setminus \{0\}$ . Therefore,  $(g_1'\varphi_2' - g_2'\varphi_1') \neq 0$ .

### 3.2. The analysis of convex and strictly psh functions

Let  $\psi_1, \psi_2, f_1, f_2, k : \mathbb{C} \rightarrow \mathbb{C}$  be holomorphic functions and  $\gamma, c \in \mathbb{C}$ . In the sequel, using the holomorphic differential equation  $k''(k+c) = \gamma(k')^2$  and the two partial differential equations  $\psi_1''(w)\overline{f_1'(z)} + \psi_2''(w)\overline{f_2'(z)} = 0$ ,  $f_1''(z)\overline{\psi_1'(w)} + f_2''(z)\overline{\psi_2'(w)} = 0$  on  $\mathbb{C}^2$ , we have

**Theorem 7.** *Let  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  and  $\varphi_1, \varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  be four holomorphic functions. Put  $u(z, w) = |\varphi_1(w) - g_1(z)|^2 + |\varphi_2(w) - g_2(z)|^2$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ . The following conditions are equivalent*

- (I)  $u$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ ;  
 (II)  $n = m = 1$ ,  $\varphi_1''\overline{g_1'} + \varphi_2''\overline{g_2'} = 0$  and  $g_1''\overline{\varphi_1'} + g_2''\overline{\varphi_2'} = 0$  on  $\mathbb{C}^2$ ,

( $\varphi_1$  or  $\varphi_2$  is nonconstant) and ( $g_1$  or  $g_2$  is nonconstant) and we have the following cases.

Case 1. The functions  $\varphi_1$  and  $\varphi_2$  satisfies  $\varphi_1'' \neq 0$  and  $\varphi_2'' \neq 0$ .

Assume that  $g_1' \neq 0$ .

If  $g_1'' = 0$ , then  $g_2'' = 0$  on  $\mathbb{C}$  (therefore  $g_1$  and  $g_2$  are affine functions with  $g_1$  or  $g_2$  is non constant. In this case, by theorem 2 or theorem 3, we can find  $\varphi_1$  and  $\varphi_2$  by their holomorphic expressions).

If  $g_1'' \neq 0$ . Thus  $g_2'' \neq 0$ . Since  $u(z, \cdot)$  is convex on  $\mathbb{C}$  (for  $z$  fixed), then  $\varphi_2 = c\varphi_1 + \xi_0$ ,  $c, \xi_0 \in \mathbb{C}$ .

$u = |\varphi_1 - g_1|^2 + |c\varphi_1 + \xi_0 - g_2|^2$ , on  $\mathbb{C}^2$ .

Assume that  $g_2' \neq 0$ .

We have an analogous situation to the above case.

Case 2. The function  $\varphi_1$  is not affine and the function  $\varphi_2$  is affine on  $\mathbb{C}$ .

Then  $g_1$  is constant on  $\mathbb{C}$ ,  $|\varphi_1 - g_1(0)|^2$  and  $|g_2 - \varphi_2(0)|^2$  are convex functions and  $|\varphi_1'g_2'| > 0$  on  $\mathbb{C}^2$ , or

$g_2$  is affine nonconstant and  $|\varphi_1'g_2'| > 0$  on  $\mathbb{C}^2$ .

We can study also the case  $\varphi_1'' = 0$  and  $\varphi_2'' \neq 0$ .

Case 3. The functions  $\varphi_1$  and  $\varphi_2$  are affine on  $\mathbb{C}$ .

The discussion is similar to cases 1, 2 and theorem 3.

**Proof.** (I) implies (II). By lemma 2, we have  $2 \leq n + m \leq 2$ . Then  $n = m = 1$ . Since  $u$  is convex and of class  $C^2$  on  $\mathbb{C}^2$ , we have the inequality

$$\left| \frac{\partial^2 u}{\partial w^2} \beta^2 + \frac{\partial^2 u}{\partial z^2} \alpha^2 + \frac{\partial^2 u}{\partial z \partial w} \alpha \beta \right| \leq \frac{\partial^2 u}{\partial w \partial \overline{w}} |\beta|^2 + \frac{\partial^2 u}{\partial z \partial \overline{z}} |\alpha|^2 + 2\operatorname{Re}\left(\frac{\partial^2 u}{\partial \overline{z} \partial w} \overline{\alpha} \beta\right)$$

on  $\mathbb{C}^2$ . It follows that

$$|[\varphi_1''(\overline{\varphi_1} - \overline{g_1}) + \varphi_2''(\overline{\varphi_2} - \overline{g_2})]\beta^2 + [g_1''(\overline{g_1} - \overline{\varphi_1}) + g_2''(\overline{g_2} - \overline{\varphi_2})]\alpha^2| \leq |\varphi_1'\beta - g_1'\alpha|^2 + |\varphi_2'\beta - g_2'\alpha|^2$$

for each  $(\alpha, \beta) \in \mathbb{C}^2$ . If  $\alpha = 0$  and  $\beta \neq 0$ , then

$$|\varphi_1''(\overline{\varphi_1} - \overline{g_1}) + \varphi_2''(\overline{\varphi_2} - \overline{g_2})| \leq |\varphi_1'|^2 + |\varphi_2'|^2$$

on  $\mathbb{C}^2$ . Now let  $\psi(z) = g_1(z)\overline{\varphi_1''(w)} + g_2(z)\overline{\varphi_2''(w)} - \varphi_1(w)\overline{\varphi_1''(w)} - \varphi_2(w)\overline{\varphi_2''(w)}$ , for  $z \in \mathbb{C}$ , ( $w$  is fixed on  $\mathbb{C}$ ).  $\psi$  is holomorphic on  $\mathbb{C}$  and  $|\psi(z)| \leq |\varphi_1'(w)|^2 + |\varphi_2'(w)|^2$ , for

every  $z \in \mathbb{C}$ , ( $w$  fixed). Thus  $\psi$  is constant on  $\mathbb{C}$ . Consequently,  $\psi'(z) = 0$ , for each  $z \in \mathbb{C}$ . Therefore

$$g_1'(z)\overline{\varphi_1''(w)} + g_2'(z)\overline{\varphi_2''(w)} = 0$$

for each  $z, w \in \mathbb{C}$ .

Now if  $\alpha \neq 0$  and  $\beta = 0$ . We obtain  $\varphi_1'(w)\overline{g_1''(z)} + \varphi_2'(w)\overline{g_2''(z)} = 0$ , for every  $(z, w) \in \mathbb{C}^2$ .

For the rest of the proof we use theorem 1, theorem2, theorem 3 and the proof of theorem 7.

**Remark 2.** Using the above technical methods, the following three partial differential equations

$$\begin{aligned} k''(k+c) &= \gamma(k')^2, \\ \psi_1''(w)\overline{f_1'(z)} + \psi_2''(w)\overline{f_2'(z)} &= 0 \text{ on } \mathbb{C}^2, \\ f_1''(z)\overline{\psi_1'(w)} + f_2''(z)\overline{\psi_2'(w)} &= 0 \text{ on } \mathbb{C}^2, \end{aligned}$$

where  $(\psi_1, \psi_2, f_1, f_2, k : \mathbb{C} \rightarrow \mathbb{C}$  are holomorphic functions and  $\gamma, c \in \mathbb{C})$ , we can solve the analogous problem when  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}^m$  and  $u = |\varphi_1 - g_1|^2 + |\varphi_2 - g_2|^2$ ;  $\varphi_1, \varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  are four holomorphic functions with the conditions ( $\varphi_1$  or  $\varphi_2$  is nonconstant) and ( $g_1$  or  $g_2$  is nonconstant).

### 3.3. Essential properties in function theory

In the sequel, we give technical tools for the study of the following families of functions consisting of: convex and not strictly psh functions on any not empty Euclidean open ball subset of  $\mathbb{C}^n \times \mathbb{C}$ ; convex and strictly sh functions but not strictly psh on each Euclidean open ball; convex and  $n$ - strictly sh functions but not strictly psh on every open ball,... . We have

**Theorem 8.** Let  $u : D \rightarrow \mathbb{R}$  be a function of class  $C^2$ ,  $D$  is a domain of  $\mathbb{C}^n, n \geq 1$ . The following conditions are equivalent

- (I)  $u$  is not strictly psh on each not empty Euclidean open ball subset of  $D$ ;
- (II)  $u$  is not strictly psh at each point of  $D$ .

**Example.** Let  $v(z, w) = |w^N - g_1(z)|^2 + |w^N - g_2(z)|^2, n, N \in \mathbb{N}, n, N \geq 2, g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be two holomorphic functions.  $v$  is convex and not strictly psh at each point of  $\mathbb{C}^n \times \mathbb{C}$ , if for example  $g_2(z) = -g_1(z)$ , for each  $z \in \mathbb{C}^n$  and  $|g_1|^2$  is convex on  $\mathbb{C}^n$ .

**Remark 3.** (R1). Let  $u_1(z, w) = |w - z|^2, u_2(z, w) = |w - 2z|^2, (z, w) \in \mathbb{C}^2$ .  $u_1$  and  $u_2$  are  $C^\infty$  and not strictly psh functions at each point of  $\mathbb{C}^2$ . But  $u = (u_1 + u_2)$  is strictly psh on  $\mathbb{C}^2$ .

(R2). Put  $v(z) = \|z\|^4, z = (z_1, \dots, z_n) \in \mathbb{C}^n$ .  $v$  is psh on  $\mathbb{C}^n$  and strictly psh on  $\mathbb{C}^n \setminus \{0\}$ . Therefore  $v$  is strictly psh almost everywhere on  $\mathbb{C}^n$ . But  $v$  is not strictly psh on  $\mathbb{C}^n$ .

**Example.** Let  $u = (u_1 + u_2), v = (v_1 + v_2)$ , where

$$u_1(z, w) = |w - f_1(z)|^2 + |w - f_2(z)|^2,$$

$$u_2(z, w) = |w - g_1(z)|^2 + |w - g_2(z)|^2,$$

$$v_1(z, w) = |w - \overline{f_1}(z)|^2 + |w - \overline{f_2}(z)|^2,$$

$$v_2(z, w) = |w - \overline{g_1}(z)|^2 + |w - \overline{g_2}(z)|^2,$$

$$f_1(z) = -f_2(z) = (z - z^2), \quad g_1(z) = -g_2(z) = (z + z^2), \quad \text{for } (z, w) \in \mathbb{C}^2.$$

$f_1, f_2, g_1, g_2$  are holomorphic functions on  $\mathbb{C}$ . We have  $u$  and  $v$  are strictly convex functions on  $\mathbb{C}^2$ . But  $u_1, u_2, v_1, v_2$  are not convex functions on  $\mathbb{C}^2$ .

**Example.** Let  $N \in \mathbb{N}$ ,  $N \geq 2$  and  $A \in \mathbb{R}_+$ ,  $A \geq 2$  such that  $\psi$  is convex on  $\mathbb{C}$ ,  $\psi(z) = A|z|^2 + |z^N - 1|^2$ , for  $z \in \mathbb{C}$ . Put  $u = (u_1 + u_2)$ , where  $u_1(z, w) = |w - g_1(z)|^2 + |w - g_2(z)|^2$ ,  $u_2(z, w) = |w - \overline{g_1}(z)|^2 + |w - \overline{g_2}(z)|^2$ ,  $g_1(z) = Az + (z^N - 1)$ ,  $g_2(z) = Az - (z^N - 1)$ , for  $(z, w) \in \mathbb{C}^2$ .

Note that  $g_1$  and  $g_2$  are holomorphic functions on  $\mathbb{C}$ . We have  $u_1$  is not strictly psh and not convex on  $\mathbb{C}^2$ .  $u_2$  is strictly psh and not convex on  $\mathbb{C}^2$ . But  $u$  is convex and strictly psh on  $\mathbb{C}^2$ .

We have

**Proposition 1.** Let  $g_1, g_2 : \mathbb{C} \rightarrow \mathbb{C}$  be two holomorphic functions. Put  $u(z, w) = |w - g_1(z)|^4 + |w - g_2(z)|^4$ ,  $v(z, w) = |w - \overline{g_1}(z)|^4 + |w - \overline{g_2}(z)|^4$ , for  $(z, w) \in \mathbb{C}^2$ . We have  $u$  is not strictly psh on  $\mathbb{C}^2$ , for each tuple of holomorphic functions  $g_1$  and  $g_2$ . But there exists several cases where  $v$  is strictly psh on  $\mathbb{C}^2$ .

**Proof.**  $u$  and  $v$  are functions of class  $C^\infty$  on  $\mathbb{C}^2$ . The hermitian Levi form of  $u$  is  $L(u)(z, w)(\alpha, \beta) = 4|w - g_1(z)|^2|\beta - g'_1(z)\alpha|^2 + 4|w - g_2(z)|^2|\beta - g'_2(z)\alpha|^2$ , for  $(z, w) \in \mathbb{C}^2$ ,  $(\alpha, \beta) \in \mathbb{C}^2$ .

Let  $z_0 \in \mathbb{C}$ . Put  $w_0 = g_1(z_0)$ . Let  $\beta = g'_2(z_0)\alpha$ , for  $\alpha \in \mathbb{C} \setminus \{0\}$ .

Then  $L(u)(z_0, w_0)(\alpha, g'_2(z_0)\alpha) = 0$  and  $\alpha \neq 0$ .

The hermitian Levi form of  $v$  is

$L(v)(z, w)(\alpha, \beta) = (2|g'_1(z)|^2|w - \overline{g_1}(z)|^2 + 2|g'_2(z)|^2|w - \overline{g_2}(z)|^2)|\alpha|^2 + (2|w - \overline{g_1}(z)|^2 + 2|w - \overline{g_2}(z)|^2)|\beta|^2 + 2|g'_1(z)(w - \overline{g_1}(z))\alpha - (\overline{w} - g_1(z))\beta|^2 + 2|g'_2(z)(w - \overline{g_2}(z))\alpha - (\overline{w} - g_2(z))\beta|^2$ , for  $(z, w), (\alpha, \beta) \in \mathbb{C}^2$ . Now choose  $|g'_1| > 0$ ,  $|g'_2| > 0$  and  $|g_1 - g_2| > 0$  on  $\mathbb{C}$ . Let  $(z, w) \in \mathbb{C}^2$ . We discuss the following three cases  $(\alpha \neq 0, \beta = 0)$ ,  $(\alpha = 0, \beta \neq 0)$  and  $(\alpha \neq 0, \beta \neq 0)$ , we obtain  $L(v)(z, w)(\alpha, \beta) > 0$  if  $(\alpha, \beta) \in \mathbb{C}^2 \setminus \{0\}$ .

Then  $v$  is strictly psh on  $\mathbb{C}^2$ .

Let  $\psi(z, w) = |w - \psi_1(z)|^2 + |w - \psi_2(z)|^2 + |w - \psi_3(z)|^2$ ,

$\varphi(z, w) = |w - \overline{\psi_1}(z)|^2 + |w - \overline{\psi_2}(z)|^2 + |w - \overline{\psi_3}(z)|^2$ , for  $(z, w) \in \mathbb{C}^2$ , where  $\psi_1, \psi_2, \psi_3 : \mathbb{C} \rightarrow \mathbb{C}$  are three holomorphic functions. Recall that if  $\psi$  is strictly psh on  $\mathbb{C}^2$ , then  $\varphi$  is strictly psh on  $\mathbb{C}^2$ . But we have

**Proposition 2.** There exists three holomorphic functions  $g_1, g_2, g_3 : \mathbb{C} \rightarrow \mathbb{C}$  such that if we define  $u(z, w) = |w - g_1(z)|^4 + |w - g_2(z)|^4 + |w - g_3(z)|^4$  and  $v(z, w) = |w - \overline{g_1}(z)|^4 + |w - \overline{g_2}(z)|^4 + |w - \overline{g_3}(z)|^4$ , for  $(z, w) \in \mathbb{C}^2$ . We have  $u$  is convex on  $\mathbb{C}^2$  and strictly psh on a neighborhood of  $(0, i)$ . But  $v$  is not strictly psh at  $(0, i)$ , while  $v$  is convex on  $\mathbb{C}^2$ .

**Example.** Let  $g_1(z) = z - i$ ,  $g_2(z) = 2z - i$ ,  $g_3(z) = 3z - i$ ,  $z \in \mathbb{C}$ .  $g_1, g_2$  and  $g_3$  are holomorphic functions on  $\mathbb{C}$ .

$z_0 = 0, w_0 = i$ . Put  $u(z, w) = |w - g_1(z)|^4 + |w - g_2(z)|^4 + |w - g_3(z)|^4$ ,  
 $v(z, w) = |w - \overline{g_1}(z)|^4 + |w - \overline{g_2}(z)|^4 + |w - \overline{g_3}(z)|^4$ , for  $(z, w) \in \mathbb{C}^2$ .

Then  $u$  and  $v$  are functions of class  $C^\infty$  and convex on  $\mathbb{C}^2$ .

Let  $\psi(z, w) = |w - g_1(z)|^4$ ,  $(z, w) \in \mathbb{C}^2$ .  $\psi$  is a  $C^\infty$  function on  $\mathbb{C}^2$  and the hermitian Levi form of  $\psi$  is

$$L(\psi)(z, w)(\alpha, \beta) = 4|w - g_1(z)|^2|\beta - g_1'(z)\alpha|^2, \quad (\alpha, \beta) \in \mathbb{C}^2.$$

Denote by  $L(u)(z, w)(\alpha, \beta)$  the hermitian Levi form of  $u$  at  $(z, w)$  and  $(\alpha, \beta)$ . Then  $L(u)(z_0, w_0)(\alpha, \beta) = 16|\beta - \alpha|^2 + 16|\beta - 2\alpha|^2 + 16|\beta - 3\alpha|^2 = 0$  implies that  $\alpha = \beta = 0$ . Thus  $L(u)(z_0, w_0)(\alpha, \beta) > 0$ , for each  $(\alpha, \beta) \in \mathbb{C}^2 \setminus \{0\}$ .

Let  $S = \{(\alpha, \beta) \in \mathbb{C}^2 / |\alpha|^2 + |\beta|^2 = 1\}$ . Thus  $\{(z_0, w_0)\} \times S = K$  is a compact on  $\mathbb{C}^2 \times \mathbb{C}^2$ .

The function  $F$ , defined by

$$F(z, w)(\alpha, \beta) = \frac{\partial^2 u}{\partial z \partial \bar{z}}(z, w)|\alpha|^2 + \frac{\partial^2 u}{\partial w \partial \bar{w}}(z, w)|\beta|^2 + 2\text{Re}\left(\frac{\partial^2 u}{\partial \bar{z} \partial w}(z, w)\bar{\alpha}\beta\right),$$

is continuous on  $\mathbb{C}^2 \times \mathbb{C}^2$ .

Since  $F > 0$  on  $K$ , then  $F > 0$  on  $B((z_0, w_0), r) \times S$ , where  $r > 0$ . Therefore  $u$  is strictly psh on a neighborhood of  $(0, i)$  and convex on  $\mathbb{C}^2$ .

The hermitian Levi form of the  $C^\infty$  function  $\theta$  on  $\mathbb{C}^2$  is

$$\begin{aligned} L(\theta)(z, w)(\alpha, \beta) &= 2|g_1'(z)(w - \overline{g_1}(z))\alpha - (\bar{w} - g_1(z))\beta|^2 + 2|g_1'(z)(\bar{w} - g_1(z))\alpha|^2 \\ &+ 2|w - \overline{g_1}(z)|^2|\beta|^2, \end{aligned}$$

for  $(z, w), (\alpha, \beta) \in \mathbb{C}^2$ , where  $\theta(z, w) = |w - \overline{g_1}(z)|^4$ .

Observe that we have  $w_0 - \overline{g_1}(z_0) = w_0 - \overline{g_2}(z_0) = w_0 - \overline{g_3}(z_0) = 0$ . Therefore  $L(v)(z_0, w_0)(\alpha, \beta) = 0$ , for each  $(\alpha, \beta) \in \mathbb{C}^2$ .

We have the following technical remark.

**Remark 4.** Let  $f_1, \dots, f_N : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic functions,  $n, N, k \in \mathbb{N} \setminus \{0\}$ ,  $k \geq 2$ . Put

$$\begin{aligned} u(z, w) &= |w - f_1(z)|^{2k} + \dots + |w - f_N(z)|^{2k}, \\ v(z, w) &= |w - \overline{f_1}(z)|^{2k} + \dots + |w - \overline{f_N}(z)|^{2k}, \\ u_1(z, w) &= |w - f_1(z)|^2 + \dots + |w - f_N(z)|^2, \\ v_1(z, w) &= |w - \overline{f_1}(z)|^2 + \dots + |w - \overline{f_N}(z)|^2, \\ \varphi &= (u + v) \text{ and } \varphi_1 = (u_1 + v_1). \end{aligned}$$

If  $u$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , we can not deduce that  $v$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ .

If  $\varphi$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , we can not conclude that  $u$  (or  $v$ ) is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ .

But we have the technical properties.

(I) If  $u$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , then  $u_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ .

(II)  $v$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$  implies that  $v_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ .

(III) If  $\varphi_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , then  $v_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ .

(IV) If  $\varphi$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , then  $\varphi_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ .

(V)  $(u + u_1)$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ , implies that  $u_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ .

For example for the proof of the above property (I), since

$u(z, w) = |(w - f_1(z))^k|^2 + \dots + |(w - f_N(z))^k|^2$ , then  $u$  is a function of class  $C^\infty$  on  $\mathbb{C}^n \times \mathbb{C}$ . Therefore the hermitian Levi form of  $u$  is

$$\begin{aligned} L(u)(z, w)(\alpha, \beta) &= |w - f_1(z)|^{2k-2} |\beta - \sum_{j=1}^n \frac{\partial f_1}{\partial z_j}(z) \alpha_j|^2 + \dots \\ &+ |w - f_N(z)|^{2k-2} |\beta - \sum_{j=1}^n \frac{\partial f_N}{\partial z_j}(z) \alpha_j|^2 \end{aligned}$$

for  $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ ,  $w \in \mathbb{C}$ ,  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{C}^n$ ,  $\beta \in \mathbb{C}$ .

Now  $u_1$  is a function of class  $C^\infty$  on  $\mathbb{C}^n \times \mathbb{C}$ . The hermitian Levi form of  $u_1$  is

$$L(u_1)(z, w)(\alpha, \beta) = |\beta - \sum_{j=1}^n \frac{\partial f_1}{\partial z_j}(z) \alpha_j|^2 + \dots + |\beta - \sum_{j=1}^n \frac{\partial f_N}{\partial z_j}(z) \alpha_j|^2.$$

Let  $(z, w), (\alpha, \beta) \in \mathbb{C}^n \times \mathbb{C}$ . Observe that  $L(u)(z, w)(\alpha, \beta) > 0$  implies that  $L(u_1)(z, w)(\alpha, \beta) > 0$ , because the absolute value  $|\beta - \sum_{j=1}^n \frac{\partial f_s}{\partial z_j}(z) \alpha_j|^2 \geq 0$ , for each  $s \in \{1, \dots, N\}$ .

The technical properties (II), (III), (IV) and (V) can be proved similarly.

Observe that for  $\psi : \mathbb{C}^n \rightarrow \mathbb{R}_+$ , if  $\psi^2$  is convex on  $\mathbb{C}^n$ , then  $\psi^4$  is convex on  $\mathbb{C}^n$ . The converse, for instance, is in general not true. But in the sequel, using the holomorphic differential equation,  $k''(k+c) = \gamma(k')^2$  ( $k : \mathbb{C} \rightarrow \mathbb{C}$  be holomorphic and  $c, \gamma \in \mathbb{C}$ ), we have

**Theorem 9.** *Let  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be two holomorphic functions. Put  $u(z, w) = |w - g_1(z)|^2 + |w - g_2(z)|^2$ ,  $v(z, w) = |w - g_1(z)|^4 + |w - g_2(z)|^4$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . We have*

(I) *Assume that  $v$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ , then  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ .*

(II) *Suppose that  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ , we can not conclude that  $v$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ .*

**Proof.** (I). Note that  $u$  and  $v$  are functions of class  $C^\infty$  on  $\mathbb{C}^n \times \mathbb{C}$ .

Assume that  $n = 1$ . We have

$$\left| \frac{\partial^2 v}{\partial z^2}(z, w) \alpha^2 + \frac{\partial^2 v}{\partial w^2} \beta^2 + 2 \frac{\partial^2 v}{\partial z \partial w} \alpha \beta \right| \leq L(v)(z, w)(\alpha, \beta)$$

for each  $(z, w), (\alpha, \beta) \in \mathbb{C}^2$ , where

$$L(v)(z, w)(\alpha, \beta) = \frac{\partial^2 v}{\partial z \partial \bar{z}}(z, w) |\alpha|^2 + \frac{\partial^2 v}{\partial w \partial \bar{w}} |\beta|^2 + 2 \operatorname{Re} \left( \frac{\partial^2 v}{\partial z \partial \bar{w}} \alpha \bar{\beta} \right).$$



We obtain the inequality

$$(E): \begin{aligned} & |[-2g_1''(z)w + 2g_1(z)g_1''(z) + 2(g_1'(z))^2](\bar{w}^2 - 2\bar{g}_1(z)\bar{w} + \bar{g}_1^2(z))\alpha^2 + \\ & [-2g_2''(z)w + 2g_2(z)g_2''(z) + 2(g_2'(z))^2](\bar{w}^2 - 2\bar{g}_2(z)\bar{w} + \bar{g}_2^2(z))\alpha^2 + 2(\bar{w} - \bar{g}_1(z))^2\beta^2 + \\ & 2(\bar{w} - \bar{g}_2(z))^2\beta^2 - 2g_1'(z)(\bar{w} - \bar{g}_1(z))^2\alpha\beta - 2g_2'(z)(\bar{w} - \bar{g}_2(z))^2\alpha\beta| \leq \\ & |2w\beta - 2\beta g_1(z) - 2wg_1'(z)\alpha + 2g_1'(z)g_1(z)\alpha|^2 + |2w\beta - 2\beta g_2(z) - 2wg_2'(z)\alpha + 2g_2'(z)g_2(z)\alpha|^2, \end{aligned}$$

for each  $(z, w)$ ,  $(\alpha, \beta) \in \mathbb{C}^2$ .  
If  $\beta = 0$  and  $w \in \mathbb{R}$ , the coefficient of  $w^3$  is equal to 0. Therefore  $(g_1''(z) + g_2''(z)) = 0$ , for every  $z \in \mathbb{C}$ .

Now we divide the left hand side of the inequality (E) by  $|\bar{w}|^2 > 0$  (for  $w \in \mathbb{C} \setminus \{0\}$ ) and the right hand side of (E) by  $|w|^2$  (observe that  $|\bar{w}|^2 = |w|^2$ ), and letting  $|w|$  go to  $(+\infty)$ , we obtain

$$\begin{aligned} & |(4g_1''(z)\bar{g}_1(z) + 4g_2''(z)\bar{g}_2(z) + 2(g_1'(z))^2 + 2(g_2'(z))^2)\alpha^2 + 4\beta^2 - 4(g_1'(z) + g_2'(z))\alpha\beta| \\ & \leq |2\beta - 2g_1'(z)\alpha|^2 + |2\beta - 2g_2'(z)\alpha|^2. \end{aligned}$$

Put  $\beta = g_1'(z)\alpha$ . Then

$$|4g_1''(z)\bar{g}_1(z) + 4g_2''(z)\bar{g}_2(z) + 2(g_1'(z) - g_2'(z))^2| \leq 4|g_1'(z) - g_2'(z)|^2.$$

Thus

$$|g_1''(z)(\bar{g}_1(z) - \bar{g}_2(z))|^2 \leq 6|g_1'(z) - g_2'(z)|^2$$

for each  $z \in \mathbb{C}$ . Now also we prove that

$$|g_2''(z)(\bar{g}_1(z) - \bar{g}_2(z))| \leq 6|g_1'(z) - g_2'(z)|^2$$

for every  $z \in \mathbb{C}$ . Using the triangle inequality, we have then

$$|g_1''(z)(g_1(z) - g_2(z)) - g_2''(z)(g_1(z) - g_2(z))| \leq 12|g_1'(z) - g_2'(z)|^2$$

for each  $z \in \mathbb{C}$ .

Therefore the function  $(g_1 - g_2)$  satisfies

$$|(g_1''(z) - g_2''(z))(g_1(z) - g_2(z))| \leq 12|g_1'(z) - g_2'(z)|^2$$

for every  $z \in \mathbb{C}$ . Therefore the function  $|g_1 - g_2|^2$  is convex on  $\mathbb{C}$ , by Abidi [2], (we can see [3]).

Since  $(g_1 + g_2)$  is affine on  $\mathbb{C}$ , thus  $g_1(z) = (az + b) + \varphi(z)$ ,  $g_2(z) = (az + b) - \varphi(z)$ , for each  $z \in \mathbb{C}$ , where  $\varphi : \mathbb{C} \rightarrow \mathbb{C}$  is a holomorphic function such that  $|\varphi|$  is convex on  $\mathbb{C}$ . Therefore  $u$  is convex on  $\mathbb{C}^2$ .

In the sequel, we can prove that  $g_1$  and  $g_2$  are affine functions on  $\mathbb{C}$  (see proposition 3). Assume that  $n \geq 2$ . Actually by the above case, it is easy to prove that  $g_1$  and  $g_2$  are affine functions on every complex line  $L \subset \mathbb{C}^n$ . Therefore,  $g_1$  and  $g_2$  are affine functions on  $\mathbb{C}^n$ .

(II). Assume that  $n = 1$ . Put  $g_1(z) = z^2$ ,  $g_2(z) = -z^2$ , for  $z \in \mathbb{C}$ . Then

$$u(z, w) = |w - g_1(z)|^2 + |w - g_2(z)|^2 = 2|w|^2 + 2|z|^2, \quad (z, w) \in \mathbb{C}^2.$$

Thus  $u$  is convex on  $\mathbb{C}^2$ . But  $v$  is not convex on  $\mathbb{C}^2$ , because  $v(z, 1) = |1 - z^2|^4 + |1 + z^2|^4 = 2\psi(z)$ , for each  $z \in \mathbb{C}$ . Observe that  $\psi$  is not convex in a neighborhood of  $\frac{1}{2}$ .

**Proposition 3.** *Let  $u(z, w) = |w + \langle z/a \rangle + b + \varphi(z)|^4 + |w + \langle z/a \rangle + b - \varphi(z)|^4$ ,  $a \in \mathbb{C}^n$ ,  $b \in \mathbb{C}$ ,  $\varphi : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic not affine,  $|\varphi|$  is convex on  $\mathbb{C}^n$ .*

*Then the function  $u$  is not convex on  $\mathbb{C}^n \times \mathbb{C}$ .*

**Proof.** Define  $v(z, w) = |w + \varphi(z)|^4 + |w - \varphi(z)|^4$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . Observe that  $v$  is convex on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ .

Suppose that  $n = 1$ . Since  $\varphi$  is not affine and  $|\varphi|$  is convex on  $\mathbb{C}$ , then by Abidi [3], we have the holomorphic representations

$\varphi(z) = (a_1 z + b_1)^k$ , for each  $z \in \mathbb{C}$ , where  $a_1 \in \mathbb{C} \setminus \{0\}$ ,  $b_1 \in \mathbb{C}$ ,  $k \in \mathbb{N}$ ,  $k \geq 2$ , or  
 $\varphi(z) = e^{(a_2 z + b_2)}$ , for every  $z \in \mathbb{C}$ , with  $a_2 \in \mathbb{C} \setminus \{0\}$  and  $b_2 \in \mathbb{C}$ .

Now for the study of the convexity of the function  $v$ , by an affine change of variable, we can assume that  $\varphi(z) = z^k$ , for any  $z \in \mathbb{C}$ , or  $\varphi(z) = e^z$ , for each  $z \in \mathbb{C}$ .

(I) Assume that  $\varphi(z) = z^k$ ,  $k \in \mathbb{N}$ ,  $k \geq 2$ .

If  $k = 2$ . We can see the above proof and we have the function  $F = v(\cdot, 1)$  is not convex on  $\mathbb{C}$ .

Now suppose that  $k \geq 3$ .

Define  $\psi(z) = v(z, 1)$ , for  $z \in \mathbb{C}$ . Then  $\psi$  is a function of class  $C^\infty$  on  $\mathbb{C}$ .

If  $\psi$  is convex on  $\mathbb{C}$ , then

$$\left| \frac{\partial^2 \psi}{\partial z^2}(z) \right| \leq \frac{\partial^2 \psi}{\partial z \partial \bar{z}}(z)$$

for each  $z \in \mathbb{C}$ .

$$\begin{aligned} \frac{\partial^2 \psi}{\partial z^2}(z) &= [2k^2 z^{2k-2} + 2k(k-1)z^{k-2}(1+z^k)](1+\bar{z}^k)^2 \\ &\quad + [2k^2 z^{2k-2} + 2k(k-1)z^{k-2}(z^k-1)](\bar{z}^k-1)^2. \\ \frac{\partial^2 \psi}{\partial z \partial \bar{z}}(z) &= 4k^2 |z^{2k-2}| |1+z^k|^2 + 4k^2 |z^k-1|^2 |z|^{2k-2}. \end{aligned}$$

For  $z_0 = 1$ ,  $\frac{\partial^2 \psi}{\partial z^2}(1) = 4(6k^2 - 4k) \geq 0$  and  $\frac{\partial^2 \psi}{\partial z \partial \bar{z}}(1) = 16k^2$ . Then  $\frac{\partial^2 \psi}{\partial z^2}(1) = \left| \frac{\partial^2 \psi}{\partial z^2}(1) \right| \leq \frac{\partial^2 \psi}{\partial z \partial \bar{z}}(1)$ . Therefore  $6k^2 - 4k \leq 4k^2$  and  $k \geq 3$ . This is a contradiction.

(II) Assume that  $\varphi(z) = e^z$ , for  $z \in \mathbb{C}$ .

Let  $\psi(z) = v(z, 2)$ ,  $z \in \mathbb{C}$ .  $\psi$  is a function of class  $C^\infty$  on  $\mathbb{C}$ .

$$\begin{aligned} \frac{\partial^2 \psi}{\partial z^2}(z) &= 2(2e^z + 2e^{2z})(2 + e^{\bar{z}})^2 + 2(2e^{2z} - 2e^z)(e^{\bar{z}} - 2)^2. \\ \frac{\partial^2 \psi}{\partial z \partial \bar{z}}(z) &= 4e^{(z+\bar{z})}(e^z + 2)(e^{\bar{z}} + 2) + 4e^{(z+\bar{z})}(e^z - 2)(e^{\bar{z}} - 2). \end{aligned}$$

$\frac{\partial^2 \psi}{\partial z^2}(0) = 72$  and  $\frac{\partial^2 \psi}{\partial z \partial \bar{z}}(0) = 40$ . Therefore  $\left| \frac{\partial^2 \psi}{\partial z^2}(0) \right| > \frac{\partial^2 \psi}{\partial z \partial \bar{z}}(0)$ . Then  $\psi$  is not convex on  $\mathbb{C}$ . Consequently,  $v$  is not convex on  $\mathbb{C}^2$ .

Comparing the preceding theorem and proposition 3, we observe that the exponent 2 is

special in our considerations. For instance, let  $u_k(z, w) = |w - f_1(z)|^{2k} + |w - f_2(z)|^{2k}$ ,  $k \in \mathbb{N} \setminus \{0\}$ ,  $f_1, f_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be two holomorphic functions and  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . We can prove that  $u_k$  is convex on  $\mathbb{C}^n \times \mathbb{C}$  implies that  $u_1$  is convex on  $\mathbb{C}^n \times \mathbb{C}$  if  $(k \geq 2)$ , but the converse is not true.

Let  $v_\delta(z, w) = |A_1 w - f_1(z)|^\delta + |A_2 w - f_2(z)|^\delta$ ,  $\delta \in [1, +\infty[$  and  $(A_1, A_2) \in \mathbb{C}^2 \setminus \{0\}$ . Observe that the study of the convexity of the function  $v_\delta$  is based on two additional cases.

Moreover, observe that by the above technical proof, we have

**Theorem 10.** *Let  $f_1, f_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be two holomorphic functions. Define  $u(z, w) = |w - f_1(z)|^4 + |w - f_2(z)|^4$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . We have  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $f_1$  and  $f_2$  are affine functions on  $\mathbb{C}^n$ .*

**Proof.** We can see the proof of theorem 9 and proposition 3. □

**Remark 5.** Let  $f_1(z) = z^N$ ,  $f_2(z) = -z^N$ ,  $f_3(z) = iz^N$  and  $f_4(z) = -iz^N$ ,  $N \in \mathbb{N} \setminus \{0, 1\}$ , for  $z \in \mathbb{C}$ .

Put  $u(z, w) = |w - f_1(z)|^4 + |w - f_2(z)|^4 + |w - f_3(z)|^4 + |w - f_4(z)|^4$ ,  $(z, w) \in \mathbb{C}^2$ .  $u$  is convex on  $\mathbb{C}^2$ , because  $u(z, w) = c(|w|^2 + |z^N|^2)^2$ , where  $c \in \mathbb{R}$ ,  $c > 0$ . But  $f_1, f_2, f_3$  and  $f_4$  are not affine functions.

We have the following.

**Question 1.** Let  $F_1, F_2, F_3 : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic functions. Put  $\psi_1(z) = (|F_1(z)|^4 + |F_2(z)|^4)$ ,  $\psi_2(z) = (|F_1(z)|^4 + |F_2(z)|^4 + |F_3(z)|^4)$ ,  $z \in \mathbb{C}^n$ .

(I) Is it true that  $\psi_1$  is convex on  $\mathbb{C}^n$  implies that  $F_1$  and  $F_2$  are affine functions on  $\mathbb{C}^n$ ?

(II) Assume that  $\psi_2$  is convex on  $\mathbb{C}^n$ . Is it true that  $F_1, F_2$  and  $F_3$  are affine functions on  $\mathbb{C}^n$ ?

The number of holomorphic functions is it fundamental in the above two situations?

We have

**Proposition 4.** *Let  $k \in \mathbb{N} \setminus \{0, 1\}$  and  $\varphi : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic. Define  $v(z, w) = |w + \langle z/a \rangle + b + \varphi(z)|^{2k} + |w + \langle z/a \rangle + b - \varphi(z)|^{2k}$ ,  $a \in \mathbb{C}^n, b \in \mathbb{C}$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . Assume that  $\varphi$  is not affine and  $|\varphi|$  is convex on  $\mathbb{C}^n$ . Then  $v$  is not convex on  $\mathbb{C}^n \times \mathbb{C}$ .*

**Proof.** Obviously follows from the proof of proposition 3. Observe that, using the holomorphic differential equation cited above, we have the additional result. □

**Theorem 11.** *Let  $g_1, g_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be two holomorphic functions and  $k \in \mathbb{N} \setminus \{0, 1\}$ . Put  $u(z, w) = |w - g_1(z)|^{2k} + |w - g_2(z)|^{2k}$  and  $v(z, w) = |w - g_1(z)|^2 + |w - g_2(z)|^2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .*

(I) Assume that  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ . Then  $v$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ .

(II) Suppose that  $v$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ . We can not conclude that  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ . But we have

(III)  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $g_1$  and  $g_2$  are affine functions.

**Extension of the results.** Let  $\psi_\delta = |w - f_1(z)|^\delta + |w - f_2(z)|^\delta$ ,  $\delta \in [1, +\infty[$ ,  $f_1, f_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be two holomorphic functions and  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . We observe without any assumption on  $\delta \in [1, +\infty[$ , for instance, for the study of the convexity

of the function  $\psi_\delta$ , the proof is organized in two separately cases.

Case 1.  $\delta = 2$ . (In this case, we obtain several solutions not affine functions).

Case 2.  $\delta \in [1, +\infty[\setminus\{2\}$ .

In general we have the following two remarks (R1) and (R2).

(R1). Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a function. Put  $\varphi_\delta(z, w) = |w - f(z)|^\delta$ ,  $\delta \in [1, +\infty[$  and  $(z, w) \in \mathbb{C}^2$ . We have  $\varphi_\delta$  is convex on  $\mathbb{C}^2$  if and only if  $f$  is affine (and in particular  $f$  is a function of class  $C^\infty$  on  $\mathbb{C}$ ).

(Let  $N \in \mathbb{N} \setminus \{0\}$ ,  $2N \geq \delta$  and put  $G(z, w) = |w - f(z)|^{2N}$ ,  $(z, w) \in \mathbb{C}^2$ . Suppose that  $\varphi_\delta$  is convex on  $\mathbb{C}^2$ . Consequently,  $G$  is psh on  $\mathbb{C}^2$ . By Abidi [1], it follows that  $f$  is harmonic on  $\mathbb{C}$ . Now let  $T : \mathbb{C} \rightarrow \mathbb{C}$  be an  $\mathbb{R}$ -linear bijective transformation. Consider  $M(z, w) = (T(z), w)$ , for  $(z, w) \in \mathbb{C}^2$ . Note that  $M$  is  $\mathbb{R}$ -linear and a bijective transformation on  $\mathbb{C}^2$ . Therefore  $G \circ M$  is convex on  $\mathbb{C}^2$  and consequently,  $G \circ M$  is psh on  $\mathbb{C}^2$ . Since  $G \circ M(z, w) = |w - f \circ T(z)|$ , for  $(z, w) \in \mathbb{C}^2$ . Then  $f \circ T$  is harmonic on  $\mathbb{C}$ , for any  $\mathbb{R}$ -linear transformation  $T$ . Then  $f$  is affine on  $\mathbb{C}$ ).

But if we define  $F_\delta(z, w) = |w - f_1(z)|^\delta + |w - g_1(z)|^\delta$ , where

$$f_1(z) = \begin{cases} 1 & \text{if } \operatorname{Re}(z) \geq 0 \\ -1 & \text{if } \operatorname{Re}(z) < 0 \end{cases}$$

and

$$g_1(z) = \begin{cases} -1 & \text{if } \operatorname{Re}(z) \geq 0 \\ 1 & \text{if } \operatorname{Re}(z) < 0 \end{cases}$$

for  $(z, w) \in \mathbb{C}^2$ . Then we have

$F_\delta(z, w) = |w - 1|^\delta + |w + 1|^\delta$  and consequently, the function  $F_\delta$  is convex on  $\mathbb{C}^2$ , for each  $\delta \geq 1$ . But  $f_1$  and  $g_1$  are noncontinuous functions at any point of  $\mathbb{C}$ . Moreover, we have

(R2). There exists two continuous functions  $f, g : \mathbb{C} \rightarrow \mathbb{C}$ , with  $K_\delta(z, w) = |w - f(z)|^\delta + |w - g(z)|^\delta$ ,  $(z, w) \in \mathbb{C}^2$ ,  $K_\delta$  is convex on  $\mathbb{C}^2$  (for each  $\delta \geq 1$ ), but  $f$  and  $g$  are not functions of class  $C^\infty$  on  $\mathbb{C}$ .

Example. Let  $f(z) = |x|$ ,  $g(z) = -|x|$ ,  $z = (x + iy) \in \mathbb{C}$ ,  $x = \operatorname{Re}(z)$ .

**Question 2.** Let  $\psi_1, \dots, \psi_N : \mathbb{C}^n \rightarrow \mathbb{C}$  be analytic functions,  $N, k \in \mathbb{N}$ ,  $k \geq 2$ . Define

$$\psi(z, w) = |w - \psi_1(z)|^{2k} + \dots + |w - \psi_N(z)|^{2k}, \quad (z, w) \in \mathbb{C}^n \times \mathbb{C}.$$

Assume that  $N \leq 2k - 1$  and  $\psi$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ . Characterize  $\psi_1, \dots, \psi_N$  by their analytic expressions.

**Question 3.** Let  $\varphi_1, \varphi_2, \varphi_3, \varphi_4 : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $g_1, g_2, g_3, g_4 : \mathbb{C}^n \rightarrow \mathbb{C}$  be 8 holomorphic functions. Put  $u = (u_1 + u_2)$ , where  $u_1(z, w) = |\varphi_1(w) - g_1(z)|^4 + |\varphi_2(w) - g_2(z)|^4$ ,  $u_2(z, w) = |\varphi_3(w) - g_3(z)|^4 + |\varphi_4(w) - g_4(z)|^4$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .

Characterize  $\varphi_1, \varphi_2, \varphi_3, \varphi_4, g_1, g_2, g_3, g_4$  by their expressions such that  $u_1$  and  $u_2$  are convex functions on  $\mathbb{C}^n \times \mathbb{C}$  and  $u$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ .

In the sequel, for instance, observe that there exists a great differences between the exponent 2 and the exponent 4 (or  $2k, k \in \mathbb{N} \setminus \{0, 1\}$ ) in real convexity.

We have

**Lemma 4.** (I) There exists  $\psi_1, \psi_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  two holomorphic functions such that  $|\psi_1|^2$  and  $|\psi_2|^2$  are not convex functions, while  $u = (|\psi_1|^2 + |\psi_2|^2)$  is convex on  $\mathbb{C}^n$ , but  $v = (|\psi_1|^4 + |\psi_2|^4)$  is not convex on  $\mathbb{C}^n$  (respectively  $(|\psi_1|^{2k} + |\psi_2|^{2k})$  is not convex on  $\mathbb{C}^n$  for each  $k \in \mathbb{N} \setminus \{0, 1\}$ ).

(II) There exists  $\varphi_1, \varphi_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  holomorphic functions, with  $|\varphi_1|^2$  is convex and  $|\varphi_2|^2$  is not convex on  $\mathbb{C}^n$ ,  $(|\varphi_1|^2 + |\varphi_2|^2)$  is convex on  $\mathbb{C}^n$ , but  $(|\varphi_1|^{2k} + |\varphi_2|^{2k})$  is not convex on  $\mathbb{C}^n$ , for each  $k \in \mathbb{N} \setminus \{0, 1\}$ .

(Example.  $\varphi_1(z) = 2z$ ,  $\varphi_2(z) = z^2 - 1$ ,  $z \in \mathbb{C}$ ).

We introduce this lemma because it yields the following questions.

**Question 4.** Let  $f_1, f_2 : \mathbb{C}^n \rightarrow \mathbb{C}$  be analytic functions and  $\delta \in [1, +\infty[$ . Put  $u = (|f_1|^\delta + |f_2|^\delta)$ . Suppose that  $u$  is convex on  $\mathbb{C}^n$  and  $\delta \neq 2$ . Is it true that  $|f_1|$  and  $|f_2|$  are convex functions on  $\mathbb{C}^n$ ?

**Question 5.** Let  $n, m \in \mathbb{N} \setminus \{0\}$ . Find all the holomorphic functions  $f_1, f_2 : \mathbb{C}^n \rightarrow \mathbb{C}$ ,  $\varphi_1, \varphi_2 : \mathbb{C}^m \rightarrow \mathbb{C}$ , such that  $\psi$  is convex on  $\mathbb{C}^n \times \mathbb{C}^m$ , where  $\psi(z, w) = |\varphi_1(w) - f_1(z)|^\delta + |\varphi_2(w) - f_2(z)|^\delta$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .

## 4. Some study of a particular case and algebraic method

**Theorem 12.** Let  $A_1, A_2, A_3, A_4, A_5 \in \mathbb{C} \setminus \{0\}$ . Consider  $g_1, g_2, g_3, g_4, g_5 : \mathbb{C}^n \rightarrow \mathbb{C}$  be five holomorphic functions. Define  $u_1(z, w) = |A_1w - g_1(z)|^2 + |A_2w - g_2(z)|^2$ ,  $v_1(z, w) = |A_3w - g_3(z)|^2 + |A_4w - g_4(z)|^2$ ,  $u(z, w) = u_1(z, w) + v_1(z, w) + |A_5w - g_5(z)|^2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .

The following conditions are equivalent

(I)  $u_1$  and  $v_1$  are convex functions on  $\mathbb{C}^n \times \mathbb{C}$  and  $u$  is (convex and strictly psh) on  $\mathbb{C}^n \times \mathbb{C}$ ;

(II)  $n \in \{1, 2, 3, 4\}$  and we have

$$\begin{cases} g_1(z) = A_1(\langle z/a \rangle + b) + \overline{A_2}\varphi(z) \\ g_2(z) = A_2(\langle z/a \rangle + b) - \overline{A_1}\varphi(z) \end{cases}$$

$$\begin{cases} g_3(z) = A_3(\langle z/c \rangle + d) + \overline{A_4}\psi(z) \\ g_4(z) = A_4(\langle z/c \rangle + d) - \overline{A_3}\psi(z) \end{cases}$$

and  $g_5(z) = (\langle z/\lambda \rangle + \mu)$ , (for all  $z \in \mathbb{C}^n$ , where  $a, c, \lambda \in \mathbb{C}^n$ ,  $b, d, \mu \in \mathbb{C}$ ,  $\varphi, \psi : \mathbb{C}^n \rightarrow \mathbb{C}$  are 2 holomorphic functions,  $|\varphi|$  and  $|\psi|$  are convex functions on  $\mathbb{C}^n$ ) with the following 4 cases.

(1)  $n = 4$ . We have  $(a - c, a - \lambda, (\frac{\overline{\partial\varphi}}{\partial z_1}(z), \frac{\overline{\partial\varphi}}{\partial z_2}(z), \frac{\overline{\partial\varphi}}{\partial z_3}(z), \frac{\overline{\partial\varphi}}{\partial z_4}(z)),$

$(\frac{\overline{\partial\psi}}{\partial z_1}(z), \frac{\overline{\partial\psi}}{\partial z_2}(z), \frac{\overline{\partial\psi}}{\partial z_3}(z), \frac{\overline{\partial\psi}}{\partial z_4}(z)))$  is a basis of the complex vector space  $\mathbb{C}^4$ , for all  $z \in \mathbb{C}^4$ .

(2)  $n = 3$ . Then we have for all  $z \in \mathbb{C}^3$ ,  $z = (z_1, z_2, z_3)$ ,

$(a - c, a - \lambda, (\frac{\overline{\partial\varphi}}{\partial z_1}(z), \frac{\overline{\partial\varphi}}{\partial z_2}(z), \frac{\overline{\partial\varphi}}{\partial z_3}(z)))$ , or  $(a - c, a - \lambda, (\frac{\overline{\partial\psi}}{\partial z_1}(z), \frac{\overline{\partial\psi}}{\partial z_2}(z), \frac{\overline{\partial\psi}}{\partial z_3}(z)))$ , or

$(a - c, (\frac{\partial \varphi}{\partial z_1}(z), \frac{\partial \varphi}{\partial z_2}(z), \frac{\partial \varphi}{\partial z_3}(z)), (\frac{\partial \psi}{\partial z_1}(z), \frac{\partial \psi}{\partial z_2}(z), \frac{\partial \psi}{\partial z_3}(z))),$  or  
 $(a - \lambda, (\frac{\partial \varphi}{\partial z_1}(z), \frac{\partial \varphi}{\partial z_2}(z), \frac{\partial \varphi}{\partial z_3}(z)), (\frac{\partial \psi}{\partial z_1}(z), \frac{\partial \psi}{\partial z_2}(z), \frac{\partial \psi}{\partial z_3}(z)))$  is a basis of the complex vector space  $\mathbb{C}^3$ .

(3)  $n = 2$ . Then for each  $z = (z_1, z_2) \in \mathbb{C}^2$ , the quantity  $(a - c, a - \lambda)$ , or  $(a - c, (\frac{\partial \varphi}{\partial z_1}(z), \frac{\partial \varphi}{\partial z_2}(z)))$ , or  $(a - c, (\frac{\partial \psi}{\partial z_1}(z), \frac{\partial \psi}{\partial z_2}(z)))$ , or  $(a - \lambda, (\frac{\partial \varphi}{\partial z_1}(z), \frac{\partial \varphi}{\partial z_2}(z)))$ , or  $(a - \lambda, (\frac{\partial \psi}{\partial z_1}(z), \frac{\partial \psi}{\partial z_2}(z)))$ , or  $((\frac{\partial \varphi}{\partial z_1}(z), \frac{\partial \varphi}{\partial z_2}(z)), (\frac{\partial \psi}{\partial z_1}(z), \frac{\partial \psi}{\partial z_2}(z)))$  is a basis of the complex vector space  $\mathbb{C}^2$ .

(4)  $n = 1$ . Then we have for all  $z \in \mathbb{C}$ ,  $(a - c) \neq 0$ , or  $(a - \lambda) \neq 0$ , or  $(\frac{\partial \varphi}{\partial z}(z) \neq 0)$ , or  $(\frac{\partial \psi}{\partial z}(z) \neq 0)$ .

**Proof.** (I) implies (II). Since  $u_1$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ , then

$$\begin{cases} g_1(z) = A_1(\langle z/a \rangle + b) + \overline{A_2}\varphi(z) \\ g_2(z) = A_2(\langle z/a \rangle + b) - \overline{A_1}\varphi(z) \end{cases}$$

(for each  $z \in \mathbb{C}^n$ , where  $a \in \mathbb{C}^n$ ,  $b \in \mathbb{C}$ ,  $\varphi : \mathbb{C}^n \rightarrow \mathbb{C}$  is analytic,  $|\varphi|$  is convex on  $\mathbb{C}^n$ ).  $u_2$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ , then

$$\begin{cases} g_3(z) = A_3(\langle z/c \rangle + d) + \overline{A_4}\psi(z) \\ g_4(z) = A_4(\langle z/c \rangle + d) - \overline{A_3}\psi(z) \end{cases}$$

(for every  $z \in \mathbb{C}^n$ , with  $c \in \mathbb{C}^n$ ,  $d \in \mathbb{C}$ ,  $\psi : \mathbb{C}^n \rightarrow \mathbb{C}$  is analytic and  $|\psi|$  is convex on  $\mathbb{C}^n$ ).

Note that  $u$  is a function of class  $C^\infty$  on  $\mathbb{C}^n \times \mathbb{C}$ . Now since  $u$  is convex on  $\mathbb{C}^n \times \mathbb{C}$ , then if we put  $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ ,  $w = z_{n+1}$ ,  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{C}^n$ ,  $\beta = \alpha_{n+1} \in \mathbb{C}$ , we have

$$\left| \sum_{j,k=1}^{n+1} \frac{\partial^2 u}{\partial z_j \partial z_k}(z) \alpha_j \alpha_k \right| \leq \sum_{j,k=1}^{n+1} \frac{\partial^2 u}{\partial z_j \partial \bar{z}_k}(z) \alpha_j \bar{\alpha}_k, \quad \forall (z, w) \in \mathbb{C}^n \times \mathbb{C}, \quad \forall (\alpha, \beta) \in \mathbb{C}^n \times \mathbb{C}.$$

It follows that  $g_5$  is an affine function on  $\mathbb{C}^n$ . Therefore  $g_5(z) = (\langle z/\lambda \rangle + \mu)$ , for each  $z \in \mathbb{C}^n$ , where  $\lambda \in \mathbb{C}^n$  and  $\mu \in \mathbb{C}$ . Then

$$\begin{aligned} u(z, w) &= (|A_1|^2 + |A_2|^2)(|w - \langle z/a \rangle - b|^2 + |\varphi(z)|^2) \\ &+ (|A_3|^2 + |A_4|^2)(|w - \langle z/c \rangle - d|^2 + |\psi(z)|^2) \\ &+ |A_5 w - \langle z/\lambda \rangle - \mu|^2, \quad (z, w) \in \mathbb{C}^n \times \mathbb{C}. \end{aligned}$$

Define

$$\begin{aligned} v(z, w) &= |w - \langle z/a \rangle - b|^2 + |\varphi(z)|^2 + |w - \langle z/c \rangle - d|^2 \\ &+ |\psi(z)|^2 + |A_5 w - \langle z/\lambda \rangle - \mu|^2, \quad (z, w) \in \mathbb{C}^n \times \mathbb{C}. \end{aligned}$$

Then  $v$  is a function of class  $C^\infty$  on  $\mathbb{C}^n \times \mathbb{C}$  and we have ( $u$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $v$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ ). Therefore by lemma 1,  $n + 1 \leq 5$ . Consequently,  $n \in \{1, 2, 3, 4\}$ .

Now let  $T(z, w) = (z, w + \langle z/a \rangle)$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .  $T$  is a  $\mathbb{C}$ -linear bijective transformation on  $\mathbb{C}^n \times \mathbb{C}$ .

Let  $v_2(z, w) = v \circ T(z, w) = |w - b|^2 + |\varphi(z)|^2 + |w + \langle z/a - c \rangle - d|^2 + |\psi(z)|^2 + |A_5 w + \langle z/a - \lambda \rangle - \mu|^2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .

Therefore  $v_2$  is a function of class  $C^\infty$  on  $\mathbb{C}^n \times \mathbb{C}$ . We have  $v$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $v_2$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ . The Levi hermitian form of  $v_2$  is

$$L(v_2)(z, w)(\alpha, \beta) = |\beta|^2 + \left| \sum_{j=1}^n \frac{\partial \varphi}{\partial z_j}(z) \alpha_j \right|^2 + |\beta + \langle \alpha/a - c \rangle|^2 +$$

$$\left| \sum_{j=1}^n \frac{\partial \psi}{\partial z_j}(z) \alpha_j \right|^2 + |A_5 \beta + \langle \alpha/a - \lambda \rangle|^2, (z, w) = ((z_1, \dots, z_n), w) \in \mathbb{C}^n \times \mathbb{C}, (\alpha, \beta) = ((\alpha_1, \dots, \alpha_n), \beta) \in \mathbb{C}^n \times \mathbb{C}.$$

Now  $L(v_2)(z, w)(\alpha, \beta) = 0$  if and only if  $\beta = 0$  and

$$\left| \sum_{j=1}^n \frac{\partial \varphi}{\partial z_j}(z) \alpha_j \right|^2 + |\beta + \langle \alpha/a - c \rangle|^2 + \left| \sum_{j=1}^n \frac{\partial \psi}{\partial z_j}(z) \alpha_j \right|^2 + |A_5 \beta + \langle \alpha/a - \lambda \rangle|^2 = 0,$$

$(z, w), (\alpha, \beta) \in \mathbb{C}^n \times \mathbb{C}$ . It follows that, if we define  $u_2(z) = |\varphi(z)|^2 + |\langle z/a - c \rangle - d|^2 + |\psi(z)|^2 + |\langle z/a - \lambda \rangle - \mu|^2$ , for  $z \in \mathbb{C}^n$ , then  $u_2$  is a function of class  $C^\infty$  on  $\mathbb{C}^n$ .

Now Observe that  $v_2$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $u_2$  is strictly psh on  $\mathbb{C}^n$ .

Case 1.  $n = 4$ . In this case observe that  $u$  is strictly psh on  $\mathbb{C}^4 \times \mathbb{C}$  if and only if the quantity

$$(a - c, a - \lambda, \left( \frac{\partial \varphi}{\partial z_1}(z), \frac{\partial \varphi}{\partial z_2}(z), \frac{\partial \varphi}{\partial z_3}(z), \frac{\partial \varphi}{\partial z_4}(z) \right), \left( \frac{\partial \psi}{\partial z_1}(z), \frac{\partial \psi}{\partial z_2}(z), \frac{\partial \psi}{\partial z_3}(z), \frac{\partial \psi}{\partial z_4}(z) \right))$$

is a basis of the complex vector space  $\mathbb{C}^4$ , for all  $z = (z_1, z_2, z_3, z_4) \in \mathbb{C}^4$ .

Case 2.  $n = 3$ . The Levi hermitian form of  $u_2$  is

$$L(u_2)(z)(\alpha) = \left| \sum_{j=1}^3 \frac{\partial \varphi}{\partial z_j}(z) \alpha_j \right|^2 + |\langle \alpha/a - c \rangle|^2 + \left| \sum_{j=1}^3 \frac{\partial \psi}{\partial z_j}(z) \alpha_j \right|^2 +$$

$|\langle \alpha/a - \lambda \rangle|^2$ , for  $z = (z_1, z_2, z_3) \in \mathbb{C}^3$ ,  $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{C}^3$ .

$L(u_2)(z)(\alpha) = 0$  if and only if

$$\begin{cases} \langle \alpha/a - c \rangle = 0, \\ \langle \alpha/a - \lambda \rangle = 0, \\ \sum_{j=1}^3 \frac{\partial \varphi}{\partial z_j}(z) \alpha_j = 0, \text{ and} \\ \sum_{j=1}^3 \frac{\partial \psi}{\partial z_j}(z) \alpha_j = 0. \end{cases}$$

Therefore  $u_2$  is strictly psh on  $\mathbb{C}^3$  if and only if for all  $z = (z_1, z_2, z_3) \in \mathbb{C}^3$ , we can choose a basis (of the complex vector space  $\mathbb{C}^3$ ) consisting of 3 vectors from the set of vectors

$\{a - c, a - \lambda, (\frac{\partial \varphi}{\partial z_1}(z), \frac{\partial \varphi}{\partial z_2}(z), \frac{\partial \varphi}{\partial z_3}(z)), (\frac{\partial \psi}{\partial z_1}(z), \frac{\partial \psi}{\partial z_2}(z), \frac{\partial \psi}{\partial z_3}(z))\}$ .

Case 3.  $n = 2$ . In this case  $u_2$  is strictly psh on  $\mathbb{C}^2$  if and only if for all  $z = (z_1, z_2) \in \mathbb{C}^2$ , we can choose a basis (consisting by 2 vectors basis of the complex vector space  $\mathbb{C}^2$ ) from the set  $\{a - c, a - \lambda, (\frac{\partial \varphi}{\partial z_1}(z), \frac{\partial \varphi}{\partial z_2}(z)), (\frac{\partial \psi}{\partial z_1}(z), \frac{\partial \psi}{\partial z_2}(z))\}$ .

Case 4.  $n = 1$ .  $u_2$  is strictly sh on  $\mathbb{C}$  if and only if for all  $z \in \mathbb{C}$ , we have  $(a - c) \neq 0$ , or  $(a - \lambda) \neq 0$ , or  $(\frac{\partial \varphi}{\partial z}(z) \neq 0)$ , or  $(\frac{\partial \psi}{\partial z}(z) \neq 0)$ .

The proof is now finished.

Moreover, we have

**Question 6.** Let  $n, m, N \in \mathbb{N} \setminus \{0\}$  and  $(A_1, B_1), \dots, (A_N, B_N) \in \mathbb{C}^2 \setminus \{(0, 0)\}$ . Find all the holomorphic functions  $g_1, f_1, \dots, g_N, f_N : \mathbb{C}^n \rightarrow \mathbb{C}$  and all the holomorphic (respectively prh) nonconstant functions  $k_1, \dots, k_N : \mathbb{C}^m \rightarrow \mathbb{C}$  such that  $u_1, \dots, u_N$  are convex and  $u = (u_1 + \dots + u_N)$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}^m$ , where  $u_j(z, w) = |A_j k_j(w) - f_j(z)|^2 + |B_j k_j(w) - g_j(z)|^2$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ ,  $1 \leq j \leq N$ . In general we prove that this question have applications in the theory of (partial differential equations and (convex and strictly psh functions) in several variables), and therefore for the resolution of certain holomorphic partial differential equations in complex analysis. Because, in the sequel, we have a relation between partial differential equations and the subject (convex and strictly psh functions) in complex analysis and geometry.

**Example.** Find all the holomorphic functions  $f, g : \mathbb{C} \rightarrow \mathbb{C}$ , such that

- (a)  $|f^2 + f|$  and  $|g^2 - g|$  are convex functions on  $\mathbb{C}$ , and
- (b)  $\psi$  is strictly psh on  $\mathbb{C}^2$ , where  $\psi(z_1, z_2) = |f^2(z_1) + f(z_1)|^2 + |g^2(z_2) - g(z_2)|^2$ , for  $(z_1, z_2) \in \mathbb{C}^2$ .

In this case we solve the holomorphic differential equation

$$(f^2 + f)''(f^2 + f) = \gamma(2ff' + f')^2, \text{ where } \gamma \in \{\frac{s-1}{s}, 1 / s \in \mathbb{N} \setminus \{0\}\}, \dots$$

**Example.** Let  $N \geq 2$ . Find all the holomorphic functions  $f_1, \dots, f_N : \mathbb{C}^n \rightarrow \mathbb{C}$ , such that  $v$  is convex and strictly psh on  $\mathbb{C}^n \times \mathbb{C}$ . We can see the problem  $v$  is convex and  $v_1$  is strictly psh on  $\mathbb{C}^n \times \mathbb{C}$  (in this case we apply lemma 1). Where

$$\begin{aligned} v(z, w) &= |w - f_1(z)|^2 + \dots + |w - f_N(z)|^2, \\ v_1(z, w) &= |w - \overline{f_1}(z)|^2 + \dots + |w - \overline{f_N}(z)|^2, \end{aligned}$$

for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . In this situation, we solve several holomorphic partial differential equations which characterize the complex structure strictly psh. Finally, we choose the solution which gives the convexity of  $v$  (or conversely).

**Question 7.** Let  $n, m, k \in \mathbb{N} \setminus \{0\}$ . Find all the holomorphic functions  $g_1, g_2, g_3, g_4 : \mathbb{C}^n \rightarrow \mathbb{C}$  and all the holomorphic functions  $\varphi_1, \varphi_2, \varphi_3, \varphi_4 : \mathbb{C}^m \rightarrow \mathbb{C}$  such that  $v_1$  and  $v_2$  are convex and  $v = (v_1 + v_2)$  is strictly convex on  $\mathbb{C}^n \times \mathbb{C}^m$ , where

$$\begin{aligned} v_1(z, w) &= |\varphi_1(w) - g_1(z)|^{2k} + |\varphi_2(w) - g_2(z)|^{2k}, \\ v_2(z, w) &= |\varphi_3(w) - g_3(z)|^{2k} + |\varphi_4(w) - g_4(z)|^{2k}, \end{aligned}$$

for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .



## 5. The product of several psh functions and applications

The main objective of this section is to study the behaviour of the product of several absolute values of prh functions. Note that it is well known that the product of many psh functions is not in general psh.

**Example.** Let  $v_1(z, w) = |w - \bar{z}||w - 2\bar{z}|$ , for  $(z, w) \in \mathbb{C}^2$ . Then  $v_1$  is not psh on  $\mathbb{C}^2$ . In the sequel, let  $D$  be a domain of  $\mathbb{C}^n$ ,  $n \in \mathbb{N} \setminus \{0\}$ ,  $N \in \mathbb{N} \setminus \{0, 1\}$  and  $\varphi_1, \dots, \varphi_N : D \rightarrow \mathbb{C}$  be holomorphic functions. Define  $u(z, w) = \prod_{1 \leq j \leq N} |w - \overline{\varphi_j(z)}|$ , for  $(z, w) \in D \times \mathbb{C}$ .

Find conditions should satisfy  $N, \varphi_1, \dots, \varphi_N$  so that  $u$  is psh on  $D \times \mathbb{C}$ .

Now let  $f_0, \dots, f_{N-1} : D \rightarrow \mathbb{C}$  be holomorphic functions. Put  $v(z, w) = |w^N + \overline{f_{N-1}(z)}w^{N-1} + \dots + \overline{f_1(z)}w + \overline{f_0(z)}|$ ,  $(z, w) \in D \times \mathbb{C}$ . Characterize  $N, f_{N-1}, \dots, f_1, f_0$  such that  $v$  is psh on  $D \times \mathbb{C}$ .

**Proposition 5.** Let  $f, g : D \rightarrow \mathbb{C}$  be two functions,  $D$  is a domain of  $\mathbb{C}^n$ ,  $n \geq 1$ . Put  $u(z, w) = |w^2 + f(z)w + g(z)|$ ,  $(z, w) \in D \times \mathbb{C}$ . Assume that  $f$  is continuous and  $g$  of class  $C^2$  on  $D$ . Then  $u$  is psh on  $D \times \mathbb{C}$  if and only if we have one assertion of the following conditions.

- (I)  $f$  is holomorphic on  $D$  and  $g$  is prh on  $D$ .
- (II)  $f$  is prh and not holomorphic and  $g = \frac{f^2}{4}$  on  $D$ .

**Proof.** Put  $v = u^2$ . Assume that  $u$  is psh on  $D \times \mathbb{C}$ . Then  $v$  is psh on  $D \times \mathbb{C}$ . By Abidi [2],  $f$  is pluriharmonic (prh) on  $D$ . Thus  $v$  is a function of class  $C^2$  on  $D \times \mathbb{C}$ . Without loss of generality we assume that  $n = 1$ . Let  $(z, w) \in D \times \mathbb{C}$ .

$$\frac{\partial^2 v}{\partial w \partial \bar{w}}(z, w) = |2w + f(z)|^2.$$

$$\frac{\partial^2 v}{\partial \bar{z} \partial w}(z, w) = \frac{\partial f}{\partial \bar{z}}(z)((\bar{w})^2 + \bar{w}\bar{f}(z) + \bar{g}(z)) + (2w + f(z))(\bar{w}\frac{\partial \bar{f}}{\partial \bar{z}}(z) + \frac{\partial \bar{g}}{\partial \bar{z}}(z)).$$

We have

$$\left| \frac{\partial^2 v}{\partial \bar{z} \partial w}(z, w) \right|^2 \leq \frac{\partial^2 v}{\partial w \partial \bar{w}}(z, w) \frac{\partial^2 v}{\partial z \partial \bar{z}}(z, w)$$

for each  $(z, w) \in \mathbb{C}^2$ . Now observe that if  $w = -\frac{f(z)}{2}$ , then  $\frac{\partial^2 v}{\partial \bar{w} \partial w}(z, -\frac{f(z)}{2}) = 0$ .

It follows that  $\frac{\partial^2 v}{\partial \bar{z} \partial w}(z, -\frac{f(z)}{2}) = 0 = \frac{\partial f}{\partial \bar{z}}(z)(\bar{g}(z) - \frac{\bar{f}^2(z)}{4})$ , for each  $z \in D$ . Now since  $f$  is real analytic on  $D$ , then  $\frac{\partial f}{\partial \bar{z}}(z) = 0$ , for every  $z \in D$ , or there exists  $z_0 \in D$ , such that  $\frac{\partial f}{\partial \bar{z}}(z_0) \neq 0$ .

Case 1. For each  $z \in D$ ,  $\frac{\partial f}{\partial \bar{z}}(z) = 0$ .

Then  $f$  is holomorphic on  $D$ . Since  $u(z, w) = |(w + \frac{f(z)}{2})^2 - \frac{f^2(z)}{4} + g(z)|$ , for  $(z, w) \in D \times \mathbb{C}$ , we consider  $T(z, w) = (z, w - \frac{f(z)}{2})$ , for  $(z, w) \in D \times \mathbb{C}$ .  $T$  is a biholomorphism on  $D \times \mathbb{C}$ . Therefore  $u \circ T$  is psh on  $D \times \mathbb{C}$ .  $u \circ T(z, w) = |w^2 - \frac{f^2(z)}{4} + g(z)|$ ,  $(z, w) \in D \times \mathbb{C}$ . By Abidi [1], the function  $(\frac{f^2}{4} - g)$  is harmonic on  $D$ . Consequently,  $g$  is harmonic on  $D$ .

Case 2. There exists  $z_0 \in D$  such that  $\frac{\partial f}{\partial \bar{z}}(z_0) \neq 0$ .

We consider  $E = \{\xi \in D / \frac{\partial f}{\partial \bar{\xi}}(\xi) = 0\}$ . Since  $\frac{\partial f}{\partial \bar{\xi}}$  is antianalytic on  $D$ , then  $E$  is an analytic closed subset on  $D$ . Therefore,  $D \setminus E$  is a domain dense on  $D$ . Now since the function  $(\frac{f^2}{4} - g)$  is continuous on  $D$  and  $(\frac{f^2}{4} - g) = 0$  on  $D \setminus E$ , then  $(\frac{f^2}{4} - g) = 0$  on  $D$ .

Let us mention that, if  $n \geq 2$  and  $f = (f_1 + \overline{f_2})$  is not holomorphic on an open polydisc  $P = P_1 \times \dots \times P_n \subset D$ , where  $f_1, f_2 : P \rightarrow \mathbb{C}$  are holomorphic functions,  $P_1, \dots, P_n$  are discs on  $\mathbb{C}$ . Since  $f_2$  is nonconstant on  $P$ , we assume that  $|\frac{\partial f_2}{\partial z_1}| > 0$  on  $P$ .

Thus  $|\frac{\partial^2 v}{\partial \bar{z}_1 \partial w}|^2 \leq \frac{\partial^2 v}{\partial \bar{z}_1 \partial z_1} \frac{\partial^2 v}{\partial \bar{w} \partial w}$  on  $P$ . Since  $\frac{\partial^2 v}{\partial \bar{w} \partial w}(z, -\overline{\frac{f(z)}{2}}) = 0$ , then  $\frac{\partial^2 v}{\partial \bar{z}_1 \partial w}(z, -\overline{\frac{f(z)}{2}}) = 0$ , for each  $z \in P$ . We obtain  $\frac{\partial f_2}{\partial z_1} [\frac{f^2}{4} - g] = 0$  on  $P$ . Consequently,  $g = \frac{f^2}{4}$  on  $P$ .

Now since  $f$  is not holomorphic on each not empty open polydisc subset of  $D$ , it follows that  $g = \frac{f^2}{4}$  on  $D$ . The proof is now complete.  $\square$

Now we have

**Theorem 13.** Let  $f, g, k : \mathbb{C}^n \rightarrow \mathbb{C}$ ,  $n \geq 1$ .

Define  $u(z, w) = |w^3 + w^2 f(z) + w g(z) + k(z)|$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .

Assume that  $f$  is continuous on  $\mathbb{C}^n$  and  $g$  and  $k$  are functions of class  $C^2$  on  $\mathbb{C}^n$ .

Then  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$  if and only if we have the following two cases.

Case 1.  $f$  and  $g$  are holomorphic functions and  $k$  is prh on  $\mathbb{C}^n$ .

Case 2.  $f$  is prh and not holomorphic on  $\mathbb{C}^n$ .

Put  $q(w) = 3w^2 + 2wf(z) + g(z)$ , for each  $w \in \mathbb{C}$  and every fixed  $z$  on  $\mathbb{C}^n$ .

$q$  have an only one zero on  $\mathbb{C}$ , for each  $z$  fixed on  $\mathbb{C}^n$ , (therefore  $g(z) = \frac{f^2(z)}{3}$  and  $k(z) = \frac{f^3(z)}{27}$ ).

**Proof.** Put  $v = u^2$ . Assume that  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$ . Then  $v$  is psh on  $\mathbb{C}^n \times \mathbb{C}$ . We can prove that  $f$  is prh on  $\mathbb{C}^n$ , using Abidi [2]. Therefore  $v$  is a function of class  $C^2$  on  $\mathbb{C}^n \times \mathbb{C}$ .

Case 1. The function  $f$  is holomorphic on  $\mathbb{C}^n$ .

$w^3 + w^2 f(z) + w g(z) + k(z) = (w + \frac{f(z)}{3})^3 + w(g(z) - \frac{f^2(z)}{3}) - \frac{f^3(z)}{27} + k(z)$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .

Since psh functions are invariant by any change by holomorphic functions, we can replace  $(w + \frac{f(z)}{3})$  by  $w$ , we obtain

$w^3 + (w - \frac{f(z)}{3})(g(z) - \frac{f^2(z)}{3}) - \frac{f^3(z)}{27} + k(z) = w^3 + w(g(z) - \frac{f^2(z)}{3}) + k_1(z)$ ,  $k_1$  is a function of class  $C^2$  on  $\mathbb{C}^n$ .

Now using the proof described in [2], we can prove that  $g$  is prh on  $\mathbb{C}^n$ . Suppose that  $g$  is holomorphic on  $\mathbb{C}^n$ . We can prove that  $k$  is prh on  $\mathbb{C}^n$ . Therefore  $u = |h|$ , where  $h : \mathbb{C}^n \times \mathbb{C} \rightarrow \mathbb{C}$  is prh. Then  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$ . Suppose that  $g$  is not holomorphic on  $\mathbb{C}^n$ . Assume that  $n = 1$ . We have

$$\begin{aligned} \frac{\partial^2 v}{\partial w \partial \bar{w}}(z, w) &= |3w^2 + 2wf(z) + g(z)|^2. \\ \frac{\partial^2 v}{\partial w \partial \bar{z}}(z, w) &= (2w \frac{\partial f}{\partial \bar{z}}(z) + \frac{\partial g}{\partial \bar{z}}(z))(w^3 + f(z)w^2 + g(z)w + k(z)) \\ &+ (3w^2 + 2wf(z) + g(z))((\bar{w})^2 \frac{\partial \bar{f}}{\partial \bar{z}}(z) + \bar{w} \frac{\partial \bar{g}}{\partial \bar{z}}(z) + \frac{\partial \bar{k}}{\partial \bar{z}}(z)). \end{aligned}$$

Since  $v$  is psh then we have the inequality

$$(E) : \left| \frac{\partial^2 v}{\partial w \partial \bar{z}}(z, w) \right|^2 \leq \frac{\partial^2 v}{\partial w \partial \bar{w}}(z, w) \frac{\partial^2 v}{\partial z \partial \bar{z}}(z, w)$$

for each  $(z, w) \in \mathbb{C}^2$ .

Since  $g$  is not holomorphic on  $\mathbb{C}$ , then there exists  $z_0 \in \mathbb{C}$ , such that  $|\frac{\partial g}{\partial \bar{z}}| > 0$  on a neighborhood of  $z_0$ .

Let  $q_1(w) = w^3 + w^2 f(z) + wg(z) + k(z)$  and  $q_2(w) = 3w^2 + 2wf(z) + g(z)$ , for  $(z, w) \in \mathbb{C}^2$ .

Note that  $q_1$  and  $q_2$  are holomorphic polynomials in the variable  $w \in \mathbb{C}$ , for each fixed  $z \in \mathbb{C}$ . Also  $q_1' = q_2$ . The holomorphic polynomial  $q_2$  has two zeros denoted  $w_1$  and  $w_2 \in \mathbb{C}$ .

Assume that  $w_1 \neq w_2$ . Then  $w_1$  and  $w_2$  are distinct zeros of the polynomial  $q_1$  by the inequality (E). Since  $q_1' = q_2$  then  $w_1$  and  $w_2$  are two distinct zeros of order 2 of  $q_1$ . A contradiction because  $\deg(q_1) = 3$ . Therefore  $w_1 = w_2$  is a zero of  $q_2$  of order 2. Thus  $w_1$  is a zero of  $q_1$  of order 3. Then we have  $q_1(w) = (w - w_1)^3$ , for every  $w \in \mathbb{C}$ . Consequently,  $f = -3w_1$  and then  $q_1(w) = (w + \frac{f(z)}{3})^3$ , for each  $z$  in a neighborhood of  $z_0$ . Then  $g(z) = \frac{f^2(z)}{3}$  and therefore  $g$  is holomorphic in a neighborhood of  $z_0$ . A contradiction. This step is impossible.

Case 2. The function  $f$  is not holomorphic on  $\mathbb{C}^n$ .

Assume that  $n = 1$ . Therefore  $\frac{\partial f}{\partial \bar{z}} \neq 0$ . Put  $q_1(w) = w^3 + w^2 f(z) + wg(z) + k(z)$ ,  $q_2(w) = 3w^2 + 2wf(z) + g(z)$ ,  $q_3(w) = 2w \frac{\partial f}{\partial \bar{z}}(z) + \frac{\partial g}{\partial \bar{z}}(z)$ , for  $(z, w) \in \mathbb{C}^2$ . Note that  $q_1$ ,  $q_2$  and  $q_3$  are holomorphic polynomials in the variable  $w \in \mathbb{C}$ , for every fixed  $z \in \mathbb{C}$ . We have  $q_1' = q_2$ . Let  $z_0 \in \mathbb{C}$ , such that  $\frac{\partial f}{\partial \bar{z}}(z) \neq 0$ , for every  $z \in V_0$ , where  $V_0$  is an Euclidean open disc in  $\mathbb{C}$ ,  $z_0 \in V_0$ . Now  $q_2$  have two zeros  $w_0(z)$  and  $w_1(z) \in \mathbb{C}$ . Suppose that  $w_0(z) = w_1(z)$ . From the inequality (E),  $w_0$  is a zero of  $q_1$ . Since  $q_2' = q_1$ , then  $w_0$  is a zero of  $q_1$  of order 3. Therefore  $q_1(w) = (w - w_0)^3$ . If for every  $z \in V_0$ ,  $w_0(z) = w_0 = w_1(z) = w_1$ , then  $q_1(w) = (w - w_0(z))^3 = (w + \frac{f(z)}{3})^3$ , in  $V_0 \times \mathbb{C}$ . Then  $g = \frac{f^2}{3}$  and  $k = \frac{f^3}{27}$ . If there exists  $z_1 \in V_0$  such that  $w_0(z_1) = a \neq w_1(z_1) = b$ . The condition  $a$  and  $b$  are zeros of  $q_1$  is impossible because  $\deg(q_1) = 3$ . By the inequality (E), for example we have  $b$  is a zero of  $q_1$  of order 2 and  $a$  is a zero of  $q_3$ .

Let  $w_2$  the second zero of  $q_1$  of order 1. Then we have the following relations between the zeros and the coefficients of the polynomial  $q_3$ ,  $a + b = -2\frac{f}{3}$ ,  $ab = 3g$  and  $2b + w_2 = -f$ . Thus we have the equalities  $3a + 3b = 4b + 2w_2$ ,  $3a = b + 2w_2$  and  $b^2 + 2bw_2 = g = \frac{1}{3}ab$ .

If  $b \neq 0$  on a neighborhood of  $z_1$ , then  $b + 2w_2 = \frac{1}{3}a = 3a$ . Consequently,  $a = 0$ .

Therefore  $g = 0$  and  $k \neq 0$ . We have then  $b = -2\frac{f}{3}$ ,  $w_2 = \frac{f}{3}$ . Thus  $u$  defined by,  $u(z, w) = |w + 2\frac{f}{3}(z)|^2|w - \frac{f}{3}(z)|$ , is psh on  $\mathbb{C}^2$ . Put  $u_1(z, w) = |w + 2f(z)|^2|w - f(z)|$ , for  $(z, w) \in \mathbb{C}^2$ . Then  $u_1$  is psh on  $\mathbb{C}^2$ . But it is obvious (by theorem 14 below), that  $f$  is holomorphic on  $\mathbb{C}$ . A contradiction. Consequently,  $b = 0$  on a neighborhood of  $z_1$ . Thus  $w_2 = -f$  and  $a = 0$  (because  $g = 0$  on a neighborhood of  $z_1$ ). Therefore  $q_2(w) = 3w^2$  and observe that  $f = 0$ . A contradiction. Therefore, the assumption  $a \neq b$  is impossible. It follows that  $w_0(z) = w_1(z)$ , for each  $z \in V_0$ . Therefore  $q_1(w) = (w - w_0)^3$ , for each  $w \in \mathbb{C}$ . We obtain  $g = \frac{f^2}{3}$  and  $k = \frac{f^3}{27}$ .

Assume now that  $n \geq 2$ . Obviously we consider in this situation an analogous proof of the above theorem as well. The proof is now complete.

Recall that for each  $f : D \rightarrow \mathbb{C}$ ,  $\psi$  is psh on  $D \times \mathbb{C}$  if and only if  $f$  is pluriharmonic (prh) on  $D$ , where  $\psi(z, w) = |w - f(z)|^N$ ,  $N \in \mathbb{N} \setminus \{0\}$ ,  $D$  is a domain of  $\mathbb{C}^n$  and  $(z, w) \in D \times \mathbb{C}$ . Now we prove that there exists a similar characterization of holomorphic functions. We have

**Theorem 14.** *Let  $f : \mathbb{C}^n \rightarrow \mathbb{C}$  be continuous. Put  $u(z, w) = |w + 2f(z)|^2|w - f(z)|$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . Then  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $f$  is holomorphic on  $\mathbb{C}^n$ .*

**Proof.** Assume that  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$ . Since  $u(z, w) = |w^3 + 3f(z)w^2 - 4f^3(z)|$ , for each  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . Then  $f$  is prh on  $\mathbb{C}^n$ , (see [2], page 336). In particular,  $f$  is a function of class  $C^\infty$  on  $\mathbb{C}^n$ . If  $f$  is holomorphic on  $\mathbb{C}^n$ , then  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$ . Assume that  $f$  is not holomorphic on  $\mathbb{C}^n$ . Then  $f$  is nonconstant. Without loss of generality we suppose that  $n = 1$  in all of the rest of the proof.

Case 1. The function  $g = \bar{f}$  is holomorphic on  $\mathbb{C}$ .

Put  $v = u^2$ . Then  $v(z, w) = |w^3 + 3\bar{g}(z)w^2 - 4\bar{g}^3(z)|^2$ ,  $(z, w) \in \mathbb{C}^2$ . Note that  $v$  is a function of class  $C^\infty$  on  $\mathbb{C}^2$ . We have

$$\begin{aligned} \frac{\partial^2 v}{\partial \bar{z} \partial w}(z, w) &= 6 \frac{\partial \bar{g}}{\partial \bar{z}}(z) w \overline{[w^3 + 3\bar{g}(z)w^2 - 4\bar{g}^3(z)]}, \\ \frac{\partial^2 v}{\partial \bar{w} \partial w}(z, w) &= |3w^2 + 6\bar{g}(z)w|^2, \\ \frac{\partial^2 v}{\partial \bar{z} \partial z}(z, w) &= |3 \frac{\partial \bar{g}}{\partial \bar{z}}(z) w^2 - 12(\bar{g})^2(z) \frac{\partial \bar{g}}{\partial \bar{z}}(z)|^2. \end{aligned}$$

Suppose that  $\frac{\partial \bar{g}}{\partial \bar{z}} = 0$  on  $\mathbb{C}$ . Then  $g$  is constant on  $\mathbb{C}$ . It follows that  $f$  is constant on  $\mathbb{C}$ . A contradiction. Consequently,  $\frac{\partial \bar{g}}{\partial \bar{z}} \neq 0$ . Since  $\frac{\partial^2 v}{\partial \bar{z} \partial z}(z, 2\bar{g}(z)) = 0$  and  $|\frac{\partial^2 v}{\partial \bar{z} \partial w}(z, w)|^2 \leq \frac{\partial^2 v}{\partial \bar{z} \partial z}(z, w) \frac{\partial^2 v}{\partial \bar{w} \partial w}(z, w)$ , for each  $(z, w) \in \mathbb{C}^2$ , then  $\frac{\partial^2 v}{\partial \bar{z} \partial w}(z, 2\bar{g}(z)) = 0$ , for any  $z \in \mathbb{C}$ . Thus  $\bar{g}(z)[16(\bar{g}^3)(z)] = 0$ , for all  $z \in \mathbb{C}$ . It follows that  $g = 0$  on  $\mathbb{C}$ . A contradiction. Therefore this case is impossible.

Case 2. The function  $g = \bar{f}$  is not holomorphic on  $\mathbb{C}$ .

Let  $v = u^2$ . Then  $v$  is a function of class  $C^\infty$  and psh on  $\mathbb{C}^2$ . Let  $g_1, g_2 : \mathbb{C} \rightarrow \mathbb{C}$  be two harmonic functions and  $(z, w) \in \mathbb{C}^2$ . Define  $F(z, w) = (w - g_1(z))^2(\bar{w} - \bar{g}_1(z))^2(w - g_2(z))(\bar{w} - \bar{g}_2(z))$ . Note that  $F$  is a  $C^\infty$  function on  $\mathbb{C}^2$ . Assume that  $F$  is psh on  $\mathbb{C}^2$ . We have

$$\begin{aligned} \frac{\partial^2 F}{\partial \bar{w} \partial w}(z, w) &= |2(w - g_2(z)) + (w - g_1(z))|^2 |w - g_1(z)|^2. \\ \frac{\partial^2 F}{\partial \bar{z} \partial w}(z, w) &= -2 \frac{\partial g_1}{\partial \bar{z}}(z) (\bar{w} - \bar{g}_1(z))^2 (w - g_2(z)) (\bar{w} - \bar{g}_2(z)) - 4 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) (w - g_1(z)) (\bar{w} - \bar{g}_1(z)) (w - g_2(z)) (\bar{w} - \bar{g}_2(z)) - 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) (w - g_1(z)) (\bar{w} - \bar{g}_1(z))^2 (w - g_2(z)) - 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) (w - g_1(z)) (\bar{w} - \bar{g}_1(z))^2 (w - g_2(z)) - 2 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) (w - g_1(z)) (\bar{w} - \bar{g}_2(z)) (\bar{w} - \bar{g}_1(z))^2 - 2 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) (\bar{w} - \bar{g}_1(z)) (\bar{w} - \bar{g}_2(z)) (w - g_1(z))^2. \end{aligned}$$

$$\begin{aligned} \frac{\partial^2 F}{\partial \bar{z} \partial z}(z, w) &= 2 \frac{\partial g_1}{\partial \bar{z}}(z) \frac{\partial g_1}{\partial z}(z) (\bar{w} - \bar{g}_1(z))^2 (w - g_2(z)) (\bar{w} - \bar{g}_2(z)) + 4 \frac{\partial g_1}{\partial \bar{z}}(z) \frac{\partial \bar{g}_1}{\partial z}(z) (\bar{w} - \bar{g}_1(z)) (w - g_1(z)) (w - g_2(z)) (\bar{w} - \bar{g}_2(z)) + 2 \frac{\partial g_1}{\partial \bar{z}}(z) \frac{\partial g_2}{\partial z}(z) (w - g_1(z)) (\bar{w} - \bar{g}_1(z))^2 (\bar{w} - \bar{g}_2(z)) + 2 \frac{\partial g_1}{\partial \bar{z}}(z) \frac{\partial \bar{g}_2}{\partial z}(z) (w - g_1(z)) (\bar{w} - \bar{g}_1(z))^2 (w - g_2(z)) + 2 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) \frac{\partial \bar{g}_1}{\partial z}(z) (w - g_1(z))^2 (w - g_2(z)) (\bar{w} - \bar{g}_2(z)) + 4 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) \frac{\partial g_1}{\partial z}(z) (w - g_1(z)) (\bar{w} - \bar{g}_1(z)) (w - g_2(z)) (\bar{w} - \bar{g}_2(z)) + 2 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) \frac{\partial \bar{g}_2}{\partial z}(z) (\bar{w} - \bar{g}_1(z)) (w - g_1(z))^2 (w - g_2(z)) + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial g_1}{\partial z}(z) (w - g_1(z)) (\bar{w} - \bar{g}_1(z))^2 (\bar{w} - \bar{g}_2(z)) + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial \bar{g}_1}{\partial z}(z) (\bar{w} - \bar{g}_1(z)) (\bar{w} - \bar{g}_2(z)) (w - g_1(z))^2 + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial \bar{g}_2}{\partial z}(z) (w - g_1(z))^2 (\bar{w} - \bar{g}_1(z))^2 + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial \bar{g}_1}{\partial z}(z) (\bar{w} - \bar{g}_1(z)) (w - g_1(z))^2 (w - g_2(z)). \end{aligned}$$

Let  $\eta > 0$ . Observe that if we replace  $g_1$  and  $g_2$  respectively by  $\eta g_1$  and  $\eta g_2$ , the new function  $F_1$ , defined by  $F_1(z, w) = |w - \eta g_1(z)|^4 |w - \eta g_2(z)|^2$  for  $(z, w) \in \mathbb{C}^2$ , is also of class  $C^\infty$  and psh on  $\mathbb{C}^2$ .

Therefore if we divide by  $\eta^2$  and letting  $\eta$  go to 0, then

$$\lim_{\eta \rightarrow 0^+} \frac{1}{\eta^2} \left| \frac{\partial^2 F_1}{\partial \bar{z} \partial w}(z, w) \right|^2 \leq \lim_{\eta \rightarrow 0^+} \left[ \frac{1}{\eta^2} \frac{\partial^2 F_1}{\partial \bar{w} \partial w}(z, w) \frac{\partial^2 F_1}{\partial \bar{z} \partial z}(z, w) \right].$$

Let  $N \in \mathbb{N} \setminus \{0\}$ . Write  $f = f_1 + \bar{f}_2$ , where  $f_1$  and  $f_2$  are holomorphic functions on  $\mathbb{C}$ . Consider  $T(z, w) = (z, w + N f_1(z))$ ,  $(z, w) \in \mathbb{C}^2$ .  $T$  is a biholomorphism of  $\mathbb{C}^2$ . Therefore  $u \circ T$  is a function of class  $C^\infty$  and psh on  $\mathbb{C}^2$ .

$$u \circ T(z, w) = |w + (N + 2)f_1(z)2\bar{f}_2(z)|^2 |w + (N - 1)f_1(z) - \bar{f}_2(z)|.$$

Define  $g_1 = -(N + 2)f_1 + 2\bar{f}_2$  and  $g_2 = -(N - 1)f_1 + \bar{f}_2$  on  $\mathbb{C}$ .

$g_1$  and  $g_2$  are harmonic functions on  $\mathbb{C}$ .

Thus for  $w_0 = 1$  and using the above inequality and letting  $\eta$  go to 0

(we replace  $g_1$  and  $g_2$  respectively by  $\eta g_1$  and  $\eta g_2$ ).

We obtain

$$\begin{aligned} & \left| 2 \frac{\partial g_1}{\partial \bar{z}}(z) + 4 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) + 2 \frac{\partial g_2}{\partial \bar{z}}(z) + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) + \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) + 2 \frac{\partial g_1}{\partial \bar{z}}(z) + 2 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) \right|^2 \leq \\ & 9 \left[ 2 \frac{\partial g_1}{\partial \bar{z}}(z) \frac{\partial g_1}{\partial z}(z) + 4 \frac{\partial g_1}{\partial \bar{z}}(z) \frac{\partial \bar{g}_1}{\partial z}(z) + 2 \frac{\partial g_1}{\partial \bar{z}}(z) \frac{\partial g_2}{\partial z}(z) + 2 \frac{\partial g_1}{\partial \bar{z}}(z) \frac{\partial \bar{g}_2}{\partial z}(z) + 2 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) \frac{\partial \bar{g}_1}{\partial z}(z) + \right. \\ & 4 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) \frac{\partial g_1}{\partial z}(z) + 2 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) \frac{\partial g_2}{\partial z}(z) + 2 \frac{\partial \bar{g}_1}{\partial \bar{z}}(z) \frac{\partial \bar{g}_2}{\partial z}(z) + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial g_1}{\partial z}(z) + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial \bar{g}_1}{\partial z}(z) + \\ & \left. \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial \bar{g}_2}{\partial z}(z) + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial g_1}{\partial z}(z) + \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial g_2}{\partial z}(z) + 2 \frac{\partial \bar{g}_2}{\partial \bar{z}}(z) \frac{\partial \bar{g}_1}{\partial z}(z) \right]. \end{aligned}$$

Then  $|4 \frac{\partial \bar{f}_2}{\partial \bar{z}}(z) + 4(N + 2) \frac{\partial \bar{f}_1}{\partial \bar{z}}(z) + 2 \frac{\partial \bar{f}_2}{\partial \bar{z}}(z) + 3(N - 1) \frac{\partial \bar{f}_1}{\partial \bar{z}}(z) + 4 \frac{\partial \bar{f}_2}{\partial \bar{z}}(z) + 2(N + 2) \frac{\partial \bar{f}_1}{\partial \bar{z}}(z)|^2 \leq$

$$9[4(N + 2)^2 \left| \frac{\partial f_1}{\partial z}(z) \right|^2 + 2(N + 2)(N - 1) \left| \frac{\partial f_1}{\partial z}(z) \right|^2 + 2(N - 1)(N + 2) \left| \frac{\partial f_1}{\partial z}(z) \right|^2 + A(N, z)],$$

where  $A(N, z)$  is a function defined on  $\mathbb{N} \times \mathbb{C}$  and satisfy  $\lim_{N \rightarrow +\infty} \frac{1}{N^2} A(N, z) = 0$ , for each  $z$  fixed on  $\mathbb{C}$ .

We divide the last above inequality by  $N^2$  and letting  $N$  go to  $+\infty$ . We obtain  $9 \times 9 \left| \frac{\partial f_1}{\partial z}(z) \right|^2 \leq 9 \times 8 \left| \frac{\partial f_1}{\partial z}(z) \right|^2$ , for all  $z \in \mathbb{C}$ . Thus  $\frac{\partial f_1}{\partial z}(z) = 0$ , for each  $z \in \mathbb{C}$ .

Consequently,  $f_1$  is constant on  $\mathbb{C}$ . Write  $f_1 = c$ ,  $c \in \mathbb{C}$ . Therefore  $f = c + \overline{f_2}$  on  $\mathbb{C}$ . It follows that  $g = \overline{f}$  is holomorphic on  $\mathbb{C}$ . A contradiction.

Consequently, this case is impossible. Therefore the above hypothesis is false and  $f$  is holomorphic on  $\mathbb{C}$ . The converse is obvious.

**Remark 6.** To compare the above theorem and some results of [2], observe that we can not write  $(w + 2f)(w - f)$  on the form  $p(w - f)$ , where  $p$  is a holomorphic polynomial on  $\mathbb{C}$  and  $f : \mathbb{C} \rightarrow \mathbb{C}$ ,  $f \neq 0$ . But if  $q$  is the following holomorphic polynomial on  $\mathbb{C}^2$ , defined by  $q(\xi, w) = (w + 2\xi)(w - \xi)$ , for  $(\xi, w) \in \mathbb{C}^2$ , we can write  $(w + 2f)(w - f) = q(f, w)$ . Denote  $\psi(\xi, w) = |q(\xi, w)|$ . Then  $\psi$  is not psh on  $\mathbb{C}^2$ . (We say in this case that  $|q|$  characterize holomorphic functions).

**Proposition 6.** Let  $f_1, g_1, f_2, g_2, f_3, g_3 : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic functions. Define  $u(z, w) = |w - f_1(z) - \overline{g_1}(z)| |w - f_2(z) - \overline{g_2}(z)| |w - f_3(z) - \overline{g_3}(z)|$ , for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . The following conditions are equivalent

(I)  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$ ;

(II) We have only case 1, or case 2.

Case 1.  $(g_1 + g_2 + g_3)$  is constant,

$(f_1 + \overline{g_1})(f_2 + \overline{g_2}) + (f_1 + \overline{g_1})(f_3 + \overline{g_3}) + (f_2 + \overline{g_2})(f_3 + \overline{g_3})$  is holomorphic on  $\mathbb{C}^n$  and  $(f_1 + \overline{g_1})(f_2 + \overline{g_2})(f_3 + \overline{g_3})$  is prh on  $\mathbb{C}^n$ .

Case 2.  $(g_1 + g_2 + g_3)$  is non constant and

$(f_1 + \overline{g_1})(f_2 + \overline{g_2}) + (f_1 + \overline{g_1})(f_3 + \overline{g_3}) + (f_2 + \overline{g_2})(f_3 + \overline{g_3}) = \frac{1}{3}(f_1 + f_2 + f_3 + \overline{g_1} + \overline{g_2} + \overline{g_3})^2$  on  $\mathbb{C}^n$ .

**Proof.** Obvious by the preceding theorem. In general, we have the following problems.

**Problem 1.** Let  $n, N \in \mathbb{N}$ ,  $N \geq 2$ ,  $D$  is a domain of  $\mathbb{C}^n$ . Find all the analytic functions  $g_1, \dots, g_N : D \rightarrow \mathbb{C}$ , such that  $u$  is psh on  $D \times \mathbb{C}$ . Here  $u(z, w) = |w - \overline{g_1}(z)| \dots |w - \overline{g_N}(z)|$ , for  $(z, w) \in D \times \mathbb{C}$ .

**Problem 2.** Let  $v(z, w) = |f_1(z) - \overline{g_1}(w)| \dots |f_N(z) - \overline{g_N}(w)|$ ,  $f_1, \dots, f_N : \mathbb{C}^n \rightarrow \mathbb{C}$  and  $g_1, \dots, g_N : \mathbb{C}^m \rightarrow \mathbb{C}$  be  $2N$  holomorphic functions,  $N \geq 2$  and  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ . Find all the conditions described by  $f_1, \dots, f_N, g_1, \dots, g_N$  such that  $v$  is convex on  $\mathbb{C}^n \times \mathbb{C}^m$ .

**Problem 3.** Put  $v = |g_1 - \varphi_1| \dots |g_N - \varphi_N|$ , where  $g_1, \dots, g_N : \mathbb{C}^m \rightarrow \mathbb{C}$  and  $\varphi_1, \dots, \varphi_N : \mathbb{C}^n \rightarrow \mathbb{C}$  be  $2N$  prh functions. Establish all the conditions satisfying by  $g_1, \dots, g_N, \varphi_1, \dots, \varphi_N$  such that  $v$  is psh on  $\mathbb{C}^n \times \mathbb{C}^m$ .

**Problem 4.** Let  $a_1, \dots, a_N \in \mathbb{C}^m$ ,  $\varphi_1, \dots, \varphi_N : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic functions,  $N \geq 2$ . Put  $v(z, w) = | \langle w/a_1 \rangle - \overline{\varphi_1}(z) | \dots | \langle w/a_N \rangle - \overline{\varphi_N}(z) |$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ . Characterize  $a_1, \dots, a_N, \varphi_1, \dots, \varphi_N$ , such that  $v$  is psh on  $\mathbb{C}^n \times \mathbb{C}^m$ .

**Remark 7.** Let  $v_N(z, w) = |w - \overline{\varphi_1}(z)| \dots |w - \overline{\varphi_N}(z)|$ ,  $\varphi_1, \dots, \varphi_N : \mathbb{C}^n \rightarrow \mathbb{C}$  be holomorphic functions,  $N \geq 2$ ,  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ .

Consider the problem  $(E_N) : v_N$  is psh on  $\mathbb{C}^n \times \mathbb{C}$ .

A technical key for the study of the problem  $(E_N)$  is a consequence of the classical cases  $(E_2)$ ,  $(E_3)$  and  $(E_4)$  which are proved. Note that if  $|w - \overline{\varphi_1}| |w - \overline{\varphi_2}|$  is psh then for each holomorphic function  $\varphi_3$ , the new function  $|w - \overline{\varphi_1}| |w - \overline{\varphi_2}| |w - \overline{\varphi_3}|$  is not psh on  $\mathbb{C}^n \times \mathbb{C}$  if  $\varphi_1$  and  $\varphi_2$  are nonconstant functions and  $\varphi_3 \neq \varphi_1$ , or  $\varphi_3 \neq \varphi_2$ .

The converse. Let  $u(z, w) = |w - \overline{\varphi_1}(z)| |w - \overline{\varphi_2}(z)| |w - \overline{\varphi_3}(z)|$ . Suppose that  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$  and  $\varphi_j$  is non constant,  $1 \leq j \leq 3$ . Then  $|w - \overline{\varphi_1}| |w - \overline{\varphi_2}|$ ,  $|w - \overline{\varphi_1}| |w - \overline{\varphi_3}|$  and  $|w - \overline{\varphi_2}| |w - \overline{\varphi_3}|$  are not psh if  $(\varphi_1 + \varphi_2 + \varphi_3)$  is constant,  $\varphi_1 \neq \varphi_2$ ,  $\varphi_1 \neq \varphi_3$  and  $\varphi_2 \neq \varphi_3$ .

Recall that  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$  if and only if  $(\varphi_1 + \varphi_2 + \varphi_3)$  is constant and  $(\varphi_1 \varphi_2 + \varphi_1 \varphi_3 + \varphi_2 \varphi_3)$  is constant, or  $(\varphi_1 + \varphi_2 + \varphi_3)$  is nonconstant and  $\varphi_1 = \varphi_2 = \varphi_3$  on  $\mathbb{C}^n$ .

**Remark 8.** Consider the functions  $g_1(z) = z$ ,  $g_2(z) = -z$ ,  $g_3(z) = iz$ ,  $g_4(z) = -iz$ , for  $z \in \mathbb{C}$ .  $g_1, g_2, g_3$  and  $g_4$  are holomorphic functions on  $\mathbb{C}$ . Let  $(z, w) \in \mathbb{C}^2$ .

$$v(z, w) = |w - \overline{g_1}(z)| |w - \overline{g_2}(z)| |w - \overline{g_3}(z)| |w - \overline{g_4}(z)| = |w^2 - (\overline{z})^2| |w^2 + (\overline{z})^2| = |w^4 - (\overline{z})^4|.$$

$v = |h|$ , where  $h : \mathbb{C}^2 \rightarrow \mathbb{C}$  is prh. Then  $v$  is psh on  $\mathbb{C}^2$ . But  $v_1, v_2, v_3$  and  $v_4$  are not psh functions on  $\mathbb{C}^2$ , where

$$v_1(z, w) = |w - \overline{g_1}(z)| |w - \overline{g_2}(z)| |w - \overline{g_3}(z)|,$$

$$v_2(z, w) = |w - \overline{g_1}(z)| |w - \overline{g_2}(z)| |w - \overline{g_4}(z)|,$$

$$v_3(z, w) = |w - \overline{g_1}(z)| |w - \overline{g_3}(z)| |w - \overline{g_4}(z)|,$$

$$v_4(z, w) = |w - \overline{g_2}(z)| |w - \overline{g_3}(z)| |w - \overline{g_4}(z)|.$$

Note that a precise study of the plurisubharmonicity of the two functions  $\psi_1$  and  $\psi_2$  extends some interesting and sharp results in the framework of a slightly different direction. We can study the complex nature of the function  $\psi_3(z, w) = |w - \overline{g_1}(z)| \dots |w - \overline{g_N}(z)|$ , where  $(N = 2^k, \text{ or } N = 3 \times 2^k, k \in \mathbb{N}, k \geq 2)$ ,  $g_1, \dots, g_N : \mathbb{C}^n \rightarrow \mathbb{C}$  are holomorphic functions,  $\psi_1(z, w) = \prod_{1 \leq j \leq 4} |w - \overline{\varphi_j}(z)|$ ,  $\psi_2(z, w) = \prod_{1 \leq j \leq 8} |w - \overline{\varphi_j}(z)|$

and  $\varphi_j : \mathbb{C}^n \rightarrow \mathbb{C}$  is a holomorphic function,  $1 \leq j \leq 8$ .

In the sequel, the next result gives the exact characterization according to algebraic methods in the theory of holomorphic polynomials and related topics. We have

**Theorem 15.** Let  $\varphi_1, \varphi_2, \varphi_3, \varphi_4 : D \rightarrow \mathbb{C}$  be four holomorphic functions,  $D$  is a domain of  $\mathbb{C}$ . Put  $u(z, w) = |w - \overline{\varphi_1}(z)| |w - \overline{\varphi_2}(z)| |w - \overline{\varphi_3}(z)| |w - \overline{\varphi_4}(z)|$ ,  $(z, w) \in D \times \mathbb{C}$ . Let  $v = u^2$ . The following conditions are equivalent

(I)  $u$  is psh on  $D \times \mathbb{C}$ ;

(II) We have the following cases.

Case 1.  $\frac{\partial^2 v}{\partial \overline{z} \partial w} = 0$  on  $D \times \mathbb{C}$ .

Case 2.  $\frac{\partial^2 v}{\partial \overline{z} \partial w} \neq 0$  on  $D \times \mathbb{C}$  and we have the following two conditions.

Step 1.  $(\sum_{j=1}^4 \varphi_j)$  is nonconstant and  $\varphi_1 = \varphi_2 = \varphi_3 = \varphi_4$  on  $D$ .

Step 2.  $(\sum_{j=1}^4 \varphi_j)$  is constant on  $D$  and we have the following assertion.

There exists  $j_1, j_2, j_3, j_4$ , satisfying  $j_1 < j_2, j_3 < j_4, \{j_1, j_2, j_3, j_4\} = \{1, 2, 3, 4\}$ ,  $\varphi_{j_1} = \varphi_{j_2}, \varphi_{j_3} = \varphi_{j_4}$  and the function  $\varphi_{j_1} \varphi_{j_3}$  is nonconstant on  $D$ .

**Proof.** (I) implies (II). Let  $(z, w) \in D \times \mathbb{C}$ . We have

$$v(z, w) = |w^4 - \overline{s_1}(z)w^3 + \overline{s_2}(z)w^2 - \overline{s_3}(z)w + \overline{s_4}(z)|^2.$$

$$s_1 = \sum_{j=1}^4 \varphi_j, \quad s_2 = \sum_{1 \leq j < k \leq 4} \varphi_j \varphi_k, \quad s_3 = \sum_{1 \leq j < k < s \leq 4} \varphi_j \varphi_k \varphi_s, \quad s_4 = \varphi_1 \varphi_2 \varphi_3 \varphi_4.$$

$s_1, s_2, s_3$  and  $s_4$  are holomorphic functions on  $D$ .

$v$  is a function of class  $C^\infty$  and psh on  $D \times \mathbb{C}$ .

$$\frac{\partial^2 v}{\partial \bar{w} \partial w}(z, w) = |4w^3 - 3\overline{s_1}(z)w^2 + 2\overline{s_2}(z)w - \overline{s_3}(z)|^2 \geq 0.$$

$$\frac{\partial^2 v}{\partial \bar{z} \partial z}(z, w) = |-\overline{s'_1}(z)w^3 + \overline{s'_2}(z)w^2 - \overline{s'_3}(z)w + \overline{s'_4}(z)|^2 \geq 0 \text{ and } \frac{\partial^2 v}{\partial \bar{z} \partial w}(z, w) = (-3\overline{s'_1}(z)w^2 + 2\overline{s'_2}(z)w - \overline{s'_3}(z))[\overline{w^4 - s_1(z)w^3 + s_2(z)w^2 - s_3(z)w + s_4(z)}].$$

Since  $v$  is psh on  $D \times \mathbb{C}$ , then we have the inequality

$$(E) : \left| \frac{\partial^2 v}{\partial \bar{z} \partial w}(z, w) \right|^2 \leq \frac{\partial^2 v}{\partial \bar{z} \partial z}(z, w) \frac{\partial^2 v}{\partial \bar{w} \partial w}(z, w)$$

for each  $(z, w) \in D \times \mathbb{C}$ .

Put

$$q_1(w) = (w - \overline{\varphi_1}(z))(w - \overline{\varphi_2}(z))(w - \overline{\varphi_3}(z))(w - \overline{\varphi_4}(z)),$$

$$q_2(w) = -3\overline{s'_1}(z)w^2 + 2\overline{s'_2}(z)w - \overline{s'_3}(z),$$

$$q_3(w) = 4w^3 - 3\overline{s_1}(z)w^2 + 2\overline{s_2}(z)w - \overline{s_3}(z),$$

$$q_4(w) = -\overline{s'_1}(z)w^3 + \overline{s'_2}(z)w^2 - \overline{s'_3}(z)w + \overline{s'_4}(z).$$

$q_1, q_2, q_3$  and  $q_4$  are holomorphic polynomials on  $\mathbb{C}$ , for each fixed  $z$  on  $D$ .

We have  $q'_1 = q_3$  and  $q'_4 = q_2$ . By the inequality (E) we have then  $|q_1 q_2| \leq |q_3 q_4|$  on  $\mathbb{C}$ .

Case 1.  $q_2(w) = 0$ , for every  $w \in \mathbb{C}$  and for any  $z \in D$ .

Then  $s_1, s_2$  and  $s_3$  are constant functions on  $D$ . Therefore  $\frac{\partial^2 v}{\partial \bar{z} \partial w} = 0$  on  $D \times \mathbb{C}$ . Thus we have

$$u(z, w) = |w^4 + c_1 w^3 + c_2 w^2 + c_3 w + \overline{\varphi_1}(z) \overline{\varphi_2}(z) \overline{\varphi_3}(z) \overline{\varphi_4}(z)|$$

where  $c_1, c_2, c_3 \in \mathbb{C}$ . Therefore  $u = |h|$ , when  $h$  is a pluriharmonic (prh) function on  $D \times \mathbb{C}$ . Consequently,  $u$  is psh on  $D \times \mathbb{C}$ .

Case 2.  $q_2 \neq 0$  on  $\mathbb{C}$ .

Now fix  $z \in D$ , such that  $[-3\overline{s'_1}(z)w^2 + 2\overline{s'_2}(z)w - \overline{s'_3}(z)] \neq 0$ . Since  $q_1 q_2 \neq 0$  and the inequality (E), there exists  $c \in \mathbb{C} \setminus \{0\}$  such that  $q_3 q_4 = c q_1 q_2$  on  $\mathbb{C}$ .

Step 1.  $s'_1 \neq 0$  on  $\mathbb{C}$ .

Then  $c = \frac{4}{3}$ . We have  $A = \{\overline{\varphi_1}(z), \overline{\varphi_2}(z), \overline{\varphi_3}(z), \overline{\varphi_4}(z)\}$  is the set of all zeros of the analytic polynomial  $q_1$ . Assume that the cardinality of  $A$  is equal to 4. Observe that because of the property of the order of multiplicity of zeros of a polynomial and the



relation  $q_3q_4 = \frac{4}{3}q_1q_2$ , we have  $\overline{\varphi_1}(z), \overline{\varphi_2}(z), \overline{\varphi_3}(z), \overline{\varphi_4}(z)$  are distinct zeros of  $q_4$ . Therefore  $\deg(q_4) \geq 4$ . A contradiction. Consequently, the cardinal of the subset  $A$  is less than or equal 3.

Without loss of generality, we assume that  $\varphi_1 = \varphi_2$ . Assume that  $\varphi_1 \neq \varphi_3$ . We have  $\overline{\varphi_1}(z)$  is a zero of  $q_3$  and  $q_4$ . Note that  $\{\overline{\varphi_1}(z), \overline{\varphi_3}(z), \overline{\varphi_4}(z)\}$  is exactly the set of zeros of the holomorphic polynomial  $q_4$ . Let  $\overline{w_1}$  and  $\overline{w_2}$  the two zeros of the polynomial  $q_2$ . Indeed, for instance, using possible relations between all the coefficients of a holomorphic polynomial and its zeros, we have then

$$\begin{aligned} \overline{w_1} + \overline{w_2} &= \frac{2}{3} \frac{\overline{s_2}(z)}{\overline{s_1}(z)} = \frac{2}{3} (\overline{\varphi_1}(z) + \overline{\varphi_3}(z) + \overline{\varphi_4}(z)). \\ \overline{w_1 w_2} &= \frac{\overline{s_3}(z)}{3\overline{s_1}'(z)} = \frac{1}{3} (\overline{\varphi_1}(z)\overline{\varphi_3}(z) + \overline{\varphi_1}(z)\overline{\varphi_4}(z) + \overline{\varphi_3}(z)\overline{\varphi_4}(z)). \end{aligned}$$

Assume that  $\overline{\varphi_1}(z), \overline{\varphi_3}(z)$  and  $\overline{\varphi_4}(z)$  are zeros of  $q_4$  of order 1. Then  $\overline{w_1}$  and  $\overline{w_2}$  are not zeros of  $q_4$ .

$\overline{w_1}$  and  $\overline{w_2}$  are zeros of  $q_3$ .

Therefore  $\overline{\varphi_1}(z) + \overline{w_1} + \overline{w_2} = \frac{3}{4}\overline{s_1}(z) = \frac{3}{4}(2\overline{\varphi_1}(z) + \overline{\varphi_3}(z) + \overline{\varphi_4}(z))$ . Then  $w_1 + w_2 = \frac{2}{3}(\varphi_1(z) + \varphi_3(z) + \varphi_4(z))$ .

$w_1 + w_2 = \frac{1}{2}\varphi_1(z) + \frac{3}{4}(\varphi_3(z) + \varphi_4(z))$ . Thus  $\varphi_3(z) + \varphi_4(z) = 2\varphi_1(z)$  and then

$$\begin{aligned} w_1 + w_2 &= 2\varphi_1(z). \\ \overline{w_1 w_2} &= \frac{\overline{s_3}'(z)}{3\overline{s_1}'(z)} = \frac{1}{3} (\overline{\varphi_1}(z)\overline{\varphi_3}(z) + \overline{\varphi_1}(z)\overline{\varphi_4}(z) + \overline{\varphi_3}(z)\overline{\varphi_4}(z)) \\ &= \frac{1}{3} [2(\overline{\varphi_1}(z))^2 + \overline{\varphi_3}(z)\overline{\varphi_4}(z)]. \end{aligned}$$

We have also

$$\begin{aligned} w_1\varphi_1(z) + w_2\varphi_1(z) + w_1w_2 &= \frac{2s_2(z)}{4} = \frac{s_2(z)}{2} = \\ \frac{1}{2}(\varphi_1^2(z) + 2\varphi_1(z)\varphi_3(z) + 2\varphi_1(z)\varphi_4(z) + \varphi_3(z)\varphi_4(z)) &= \frac{1}{2}(5\varphi_1^2(z) + \varphi_3(z)\varphi_4(z)) \\ &= (w_1 + w_2)\varphi_1 + w_1w_2. \end{aligned}$$

Therefore,

$$\begin{aligned} 2\varphi_1^2(z) + w_1w_2 &= \frac{1}{2}(5\varphi_1^2(z) + \varphi_3(z)\varphi_4(z)). \\ w_1w_2 &= \frac{1}{2}\varphi_1^2(z) + \frac{1}{2}\varphi_3(z)\varphi_4(z) = \frac{1}{3}(2\varphi_1^2(z) + \varphi_3(z)\varphi_4(z)). \end{aligned}$$

Thus  $3\varphi_1^2(z) + 3\varphi_3(z)\varphi_4(z) = 4\varphi_1^2(z) + 2\varphi_3(z)\varphi_4(z)$ . Then  $\varphi_3(z)\varphi_4(z) = \varphi_1^2(z)$ . Since  $\varphi_3 + \varphi_4(z) = 2\varphi_1(z)$ . Thus  $\varphi_3(z) = \varphi_4(z) = \varphi_1(z)$ . A contradiction.

Assume now that  $\overline{\varphi_1}(z), \overline{\varphi_3}(z)$  and  $\overline{\varphi_4}(z)$  are not zeros of  $q_4$  of order 1. Recall that  $\{\overline{\varphi_1}(z), \overline{\varphi_3}(z), \overline{\varphi_4}(z)\}$  is exactly the set of zeros of  $q_4$ . Since  $\varphi_1(z) \neq \varphi_3(z)$ , then  $\varphi_1(z) = \varphi_4(z)$ .

Because if  $\varphi_1(z) \neq \varphi_4(z)$ , then  $\varphi_3(z) = \varphi_4(z)$ . Since  $\varphi_1(z) = \varphi_2(z) \neq \varphi_3(z)$ , then  $u(\xi, w) = |w - \overline{\varphi_1}(\xi)|^2 |w - \overline{\varphi_3}(\xi)|^2$ , for each  $(\xi, w) \in G = D(z, R) \times \mathbb{C}$ , where  $R > 0$  satisfying  $D(z, R) \subset D$ . Since  $u$  is a function of class  $C^\infty$  and psh on the domain  $G$ , we can prove that we have the condition  $\varphi_1 = \varphi_3$  on  $D(z, R)$ , or  $(\varphi_1 + \varphi_3) = \frac{s_1}{2}$  is constant on  $D(z, R)$ .

Now since  $s_1$  is holomorphic nonconstant on  $D$ , then  $s_1$  is nonconstant on the open Euclidean disc  $D(z, R)$ . It follows that  $\varphi_1 = \varphi_3$  on  $D(z, R)$ . A contradiction, because  $\varphi_1(z) \neq \varphi_3(z)$ . Consequently,  $\varphi_1(z) = \varphi_4(z)$ . Therefore  $\overline{\varphi_1}(z)$  is a zero of  $q_2$  of order 1. Assume that  $\varphi_1(z) = w_1$ . We have  $\varphi_1(z) = \varphi_2(z) = \varphi_4(z)$ ,  $\varphi_1(z) \neq \varphi_3(z)$ . It follows that  $\overline{\varphi_1}(z)$  is a zero of  $q_3$  of order 2.

$\overline{\varphi_3}(z)$  is not a zero of  $q_3$ .

Now we use the classical relations between all the coefficients of a polynomial and its zeros, we have  $\varphi_1(z) + w_2 = \frac{2s'_2(z)}{3s'_1(z)} = \frac{2}{3}(2\varphi_1(z) + \varphi_3(z))$ . Also  $2\varphi_1(z) + w_2 = \frac{3}{4}(3\varphi_1(z) + \varphi_3(z))$  and  $\varphi_1(z) + w_2 = \frac{2}{3}(2\varphi_1(z) + \varphi_3(z))$ . Then  $\varphi_1(z) + \frac{2}{3}(2\varphi_1(z) + \varphi_3(z)) = \frac{3}{4}(3\varphi_1(z) + \varphi_3(z))$ . Thus,  $12\varphi_1(z) + 8(2\varphi_1(z) + \varphi_3(z)) = 9(3\varphi_1(z) + \varphi_3(z))$ . Consequently,  $\varphi_1(z) = \varphi_3(z)$ . A contradiction. It follows that the assumption  $\varphi_1(z) \neq \varphi_3(z)$  is impossible. Consequently,  $\varphi_1(z) = \varphi_2(z) = \varphi_3(z)$ .

Now assume that  $\varphi_4(z) \neq \varphi_1(z)$ . Let  $\overline{w_0}$  the zero of  $q_2$ ,  $w_0 \neq \varphi_1(z)$ . Note that  $\overline{\varphi_1}(z)$  is a zero of the polynomial  $q_2$  because  $\overline{\varphi_1}(z)$  is a zero of  $q_4$  of order 2.

$\overline{\varphi_1}(z)$  is a zero of  $q_1$  of order 3.

Therefore  $\overline{\varphi_1}(z)$  is a zero of  $q_3$  of order 2. Consequently,  $\overline{w_0}$  is a zero of  $q_3$  of order 1.

We have  $w_0 + \varphi_1(z) = \frac{2s'_2(z)}{3s'_1(z)} = \frac{2}{3}(2\varphi_1(z) + \varphi_4(z))$ . Also  $2\varphi_1(z) + w_0 = \frac{3}{4}s_1(z) = \frac{3}{4}(3\varphi_1(z) + \varphi_4(z))$ . Therefore, we have  $\varphi_1(z) + \frac{2}{3}(2\varphi_1(z) + \varphi_4(z)) = \frac{3}{4}(3\varphi_1(z) + \varphi_4(z))$ . Thus  $\varphi_1(z) = \varphi_4(z)$ . A contradiction. Consequently, the assumption  $\varphi_1(z) \neq \varphi_4(z)$  is impossible. We conclude that  $\varphi_1 = \varphi_2 = \varphi_3 = \varphi_4$  on  $D$ .

Step 2.  $s_1$  is constant on  $D$ .

Let  $(z, w) \in D \times \mathbb{C}$ , such that  $\frac{\partial^2 v}{\partial \bar{z} \partial w}(z, w) \neq 0$ . Assume that  $s'_2(z) \neq 0$ . We have  $q_1 q_2 = c q_3 q_4$ , where  $c = \frac{1}{2}$ . Let  $\overline{w_0} = \frac{\overline{s'_2(z)}}{2s'_2(z)}$  be the only zero of  $q_2$ . Note that  $\{\overline{\varphi_1}(z), \overline{\varphi_2}(z), \overline{\varphi_3}(z), \overline{\varphi_4}(z)\}$  is the set of zeros of the holomorphic polynomial  $q_1$  on  $\mathbb{C}$ . If for example  $\overline{\varphi_1}(z)$  is a zero of  $q_1$  of order 1. Then  $\overline{\varphi_1}(z)$  is not a zero of  $q_3 = q'_1$ . Since now  $q_1 q_2 = \frac{1}{2} q_3 q_4$ , then  $\overline{\varphi_1}(z)$  is a zero of  $q_4$ .

Now if  $\overline{\varphi_1}(z)$  is a zero of  $q_1$  of order 2. Then  $\overline{\varphi_1}(z)$  is a zero of  $q_3 = q'_1$  of order 1. By the fundamental relation  $q_1 q_2 = \frac{1}{2} q_3 q_4$ , we obtain  $\overline{\varphi_1}(z)$  is a zero of  $q_4$ . We conclude that the set of zeros of  $q_4$  is  $\{\overline{\varphi_1}(z), \overline{\varphi_2}(z), \overline{\varphi_3}(z), \overline{\varphi_4}(z)\}$ .

Since now  $\deg(q_4) = 2$  (because  $s'_2(z) \neq 0$ ), then there exists  $j_1, j_2, j_3, j_4$ ,  $\{j_1, j_2, j_3, j_4\} = \{1, 2, 3, 4\}$ , such that  $\varphi_{j_1} = \varphi_{j_2} = \varphi_{j_3} \neq \varphi_{j_4}$  on  $D$ , or  $\varphi_{j_1} = \varphi_{j_2}$  and  $\varphi_{j_3} = \varphi_{j_4}$  on  $D$ .

Suppose that we have  $\varphi_1 = \varphi_2 = \varphi_3 \neq \varphi_4$ . Then  $\overline{\varphi_1}(z)$  is a zero of  $q_1$  of order 3.  $\overline{\varphi_1}(z)$  is a zero of  $q_3$  of order 2.

$\overline{w_0}$  is a zero of  $q_3$  of order 1.

$\overline{\varphi_4}(z)$  is not a zero of  $q_3$ .

We have

$$2\varphi_1(z) + w_0 = \frac{3}{4}s_1(z) = \frac{3}{4}(3\varphi_1(z) + \varphi_4(z)).$$

$$w_0 = \frac{s'_3(z)}{2s'_2(z)} = \frac{1}{2}(\varphi_1(z) + \varphi_4(z)).$$

Thus  $2\varphi_1(z) + \frac{1}{2}(\varphi_1(z) + \varphi_4(z)) = \frac{3}{4}(3\varphi_1(z) + \varphi_4(z))$ . Therefore  $\frac{5}{2}\varphi_1(z) + \frac{1}{2}\varphi_4(z) = \frac{9}{4}\varphi_1(z) + \frac{3}{4}\varphi_4(z)$ . Then  $\varphi_1(z) = \varphi_4(z)$ . A contradiction. Consequently, the above assumption is impossible. It follows that  $\varphi_{j_1} = \varphi_{j_2}$  and  $\varphi_{j_3} = \varphi_{j_4}$  on  $D$ , (for example). We suppose without loss of generality that  $j_1 < j_2$  and  $j_3 < j_4$ . Then  $u(z, w) = |w - \overline{\varphi_{j_1}}(z)|^2 |w - \overline{\varphi_{j_3}}(z)|^2$ , for  $(z, w) \in D \times \mathbb{C}$ .

Actually, we observe that  $\varphi_{j_1} = \varphi_{j_3}$  on  $D$ , or  $(\varphi_{j_1} + \varphi_{j_3})$  is constant on  $D$ . Suppose that  $\varphi_{j_1} = \varphi_{j_3}$  on  $D$ . Then  $\varphi_1 = \varphi_2 = \varphi_3 = \varphi_4$  on  $D$ . Since  $s'_1 = 0$  on  $D$ , then  $\varphi_1$  is constant on  $D$ . Thus  $\frac{\partial^2 v}{\partial \bar{z} \partial w} = 0$  on  $D \times \mathbb{C}$ . A contradiction. Consequently,  $(\varphi_{j_1} + \varphi_{j_3})$  is constant on  $D$  and observe that the product  $\varphi_{j_1} \varphi_{j_3}$  is nonconstant on  $D$ .

Assume that  $s'_2 = 0$  on  $D$ . Then  $s'_3 \neq 0$  on  $D$ , because  $\frac{\partial^2 v}{\partial \bar{z} \partial w} \neq 0$  on  $D \times \mathbb{C}$ . The set of zeros of  $q_4$  is  $\{\overline{\varphi_1}(z), \overline{\varphi_2}(z), \overline{\varphi_3}(z), \overline{\varphi_4}(z)\}$ . Since  $\deg(q_4) = 1$ , then  $\varphi_1 = \varphi_2 = \varphi_3 = \varphi_4$  on  $D$ .

$s'_1 = 0$  on  $D$  implies that  $\varphi_1$  is constant on  $D$ . Therefore,  $\frac{\partial^2 v}{\partial \bar{z} \partial w} = 0$  on  $D \times \mathbb{C}$ . It follows that this case is impossible.

(II) implies (I). Obvious.

**Remark 9 .** Let  $F : \mathbb{C}^2 \rightarrow \mathbb{C}$  be holomorphic,  $a_1, a_2, a_3, a_4 \in \mathbb{C}^4$  and  $b_1, b_2, b_3, b_4 \in \mathbb{C}$ . Define

$$v(z, w) = \left| [w - \overline{\langle F(z)/a_1 \rangle} - b_1][w - \overline{\langle F(z)/a_2 \rangle} - b_2] \cdot [w - \overline{\langle F(z)/a_3 \rangle} - b_3][w - \overline{\langle F(z)/a_4 \rangle} - b_4] \right|,$$

for  $(z, w) \in \mathbb{C}^2 \times \mathbb{C}$ . We can characterize all the conditions on  $a_1, a_2, a_3, a_4, b_1, b_2, b_3, b_4$ , which ensure technical hypothesis for the plurisubharmonicity of  $v$ . Indeed, we have the following of various behaviour.

**Theorem 16.** Let  $\varphi_1, \varphi_2, \varphi_3, \varphi_4 : D \rightarrow \mathbb{C}$  be holomorphic functions,  $D$  is a domain of  $\mathbb{C}^n$ ,  $n \geq 1$ .

Put  $u(z, w) = |w - \overline{\varphi_1}(z)| |w - \overline{\varphi_2}(z)| |w - \overline{\varphi_3}(z)| |w - \overline{\varphi_4}(z)|$ ,  $(z, w) \in D \times \mathbb{C}$ .

Let  $v = u^2$ ,  $s_1 = \sum_{j=1}^4 \varphi_j$ ,  $s_2 = \sum_{1 \leq j < k \leq 4} \varphi_j \varphi_k$ ,  $s_3 = \sum_{1 \leq j < k < s \leq 4} \varphi_j \varphi_k \varphi_s$ ,  $s_4 =$

$\varphi_1 \varphi_2 \varphi_3 \varphi_4$ , ( $s_1, s_2, s_3, s_4$  are holomorphic functions on  $D$ ).

The following assertions are equivalent

(I)  $u$  (respectively  $v$ ) is psh on  $D \times \mathbb{C}$ ;

(II) We have the following three cases.

Case 1.  $s_1, s_2$  and  $s_3$  are constant on  $D$ .

Case 2.  $s_1$  is nonconstant on  $D$  and  $\varphi_1 = \varphi_2 = \varphi_3 = \varphi_4$  on  $D$ .

Case 3.  $s_1$  is constant on  $D$ ,  $s_2$  is nonconstant on  $D$  and there exists  $j_1, j_2, j_3, j_4$ ,  $\{j_1, j_2, j_3, j_4\} = \{1, 2, 3, 4\}$ ,  $j_1 < j_2, j_3 < j_4$ , with  $\varphi_{j_1} = \varphi_{j_2}$  and  $\varphi_{j_3} = \varphi_{j_4}$  on  $D$ .

**Proof.** Obvious by the above theorem.

**Example.** Let  $a_1, a_2, a_3, a_4 \in \mathbb{C}^2$ ,  $A_1, A_2, A_3, A_4 \in \mathbb{C}^m$  and  $F : \mathbb{C}^n \rightarrow \mathbb{C}^2$  be a

holomorphic function,  $n, m \geq 1$ . Define

$$\psi(z, w) = \left| \langle w/A_1 \rangle - \overline{\langle F(z)/a_1 \rangle} \right| \left| \langle w/A_2 \rangle - \overline{\langle F(z)/a_2 \rangle} \right| \cdot \left| \langle w/A_3 \rangle - \overline{\langle F(z)/a_3 \rangle} \right| \left| \langle w/A_4 \rangle - \overline{\langle F(z)/a_4 \rangle} \right|,$$

for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ .

In a slightly different direction, we can show all the conditions formulated by the constants  $a_1, a_2, a_3, a_4, A_1, A_2, A_3, A_4$ , which characterize the plurisubharmonicity of  $\psi$ .

**Example.** Let  $N \geq 2$  and  $p(\xi_1, \dots, \xi_N, w) \equiv (w - \xi_1) \dots (w - \xi_N)$ , for  $(\xi_1, \dots, \xi_N) \in \mathbb{C}^N$ ,  $w \in \mathbb{C}$ . Define  $F(\xi_1, \dots, \xi_N, w) = |p(\xi_1, \dots, \xi_N, w)|$ . Then for each Euclidean open ball  $B(a, R) \subset \mathbb{C}^N$ , ( $a \in \mathbb{C}^N, R > 0$ ), the function  $F$  is not psh on  $B(a, R) \times \mathbb{C}$ .

**Remark 10.** (I) Let  $g_1(z) = z^2, g_2(z) = -z^2, g_3(z) = iz^2, g_4(z) = -iz^2, z \in \mathbb{C}$ .  $g_1, g_2, g_3$  and  $g_4$  are holomorphic functions on  $\mathbb{C}$ . Let

$$u_1(z, w) = |w - \overline{g_1(z)}| |w - \overline{g_2(z)}| |w - \overline{g_3(z)}|,$$

$$u_2(z, w) = |w - \overline{g_1(z)}| |w - \overline{g_2(z)}| |w - \overline{g_4(z)}|,$$

$$u_3(z, w) = |w - \overline{g_1(z)}| |w - \overline{g_3(z)}| |w - \overline{g_4(z)}|,$$

$$u_4(z, w) = |w - \overline{g_2(z)}| |w - \overline{g_3(z)}| |w - \overline{g_4(z)}|,$$

$$u(z, w) = |w - \overline{g_1(z)}| |w - \overline{g_2(z)}| |w - \overline{g_3(z)}| |w - \overline{g_4(z)}|, (z, w) \in \mathbb{C}^2.$$

We have  $u_1, u_2, u_3$  and  $u_4$  are not psh functions on  $\mathbb{C}^2$ . But  $u$  is psh on  $\mathbb{C}^2$ .

(II)  $g_1(z) = g_2(z) = z + 1, g_3(z) = g_4(z) = -z + 1, z \in \mathbb{C}$ .

$g_1, g_2, g_3$  and  $g_4$  are holomorphic functions on  $\mathbb{C}$ .

$(g_1 + g_2 + g_3 + g_4)$  is constant on  $\mathbb{C}$ .

$(g_1g_2 + g_1g_3 + g_1g_4 + g_2g_3 + g_2g_4 + g_3g_4)$  is non constant on  $\mathbb{C}$ .

Let  $(z, w) \in \mathbb{C}^2$ . Put  $u(z, w) = |w - \overline{g_1(z)}| |w - \overline{g_2(z)}| |w - \overline{g_3(z)}| |w - \overline{g_4(z)}| = |w^4 - 4w^3 + [6 - 2(\overline{z})^2]w^2 - 4w[1 - (\overline{z})^2] + [1 - (\overline{z})^2]^2|$ . Observe that  $(g_1g_2g_3 + g_1g_2g_4 + g_1g_3g_4 + g_2g_3g_4)$  is nonconstant on  $\mathbb{C}$ . But  $u$  is psh on  $\mathbb{C}^2$ , because

$$u(z, w) = |w - 1 - \overline{z}|^2 |w - 1 + \overline{z}|^2 = |(w - 1)^2 - (\overline{z})^2|^2 = |h|^2,$$

where  $h$  is a prh function on  $\mathbb{C}^2$ .

**Question 8.** Let  $N_1, \dots, N_k, s_1, m_1, \dots, s_t, m_t \in \mathbb{N} \setminus \{0\}$ ,  $k, t \geq 1$  and  $g_1, \dots, g_k, \theta_1, \dots, \theta_t : \mathbb{C}^n \rightarrow \mathbb{C}$  be prh functions. Put

$$u(z, w) = |w - g_1(z)|^{N_1} \dots |w - g_k(z)|^{N_k} |w^{s_1} - \theta_1^{m_1}(z)| \dots |w^{s_t} - \theta_t^{m_t}(z)|,$$

for  $(z, w) \in \mathbb{C}^n \times \mathbb{C}$ . Find conditions  $g_1, \dots, g_k, \theta_1, \dots, \theta_t$  should satisfy so that  $u$  is psh on  $\mathbb{C}^n \times \mathbb{C}$ .

**Question 9.** Let  $N \in \mathbb{N} \setminus \{0, 1\}$  and  $A_0, \dots, A_{N-1} \in \mathbb{C}$ . Define  $v(z, w) = |w^N + A_{N-1}w^{N-1}\overline{z} + \dots + A_1wz^{N-1} + A_0z^N|$ . Find all conditions on  $N, A_0, \dots, A_{N-1}$  such that  $v$  is psh on  $\mathbb{C}^2$ .

Conclude that we can characterize all the holomorphic polynomials  $q$  on  $\mathbb{C}^2$ , such that  $F$  is psh on  $\mathbb{C}^2$ , where  $F(z, w) = |q(\bar{z}, w)|$  for  $(z, w) \in \mathbb{C}^2$ .

Let  $p$  be a holomorphic polynomial on  $\mathbb{C}^2$ . Put  $F_1(z, w) = |p(\bar{z}, w)|$  and  $F_2(z, w) = |p(z, \bar{w})|$ , for  $(z, w) \in \mathbb{C}^2$ . Moreover, thanks to the above characterization, we can prove that  $F_1$  is psh on  $\mathbb{C}^2$  if and only if  $F_2$  is psh on  $\mathbb{C}^2$ .

In the following question, we recall some properties and sharp results in the framework of complex analysis of the appeared function  $\theta$ , defined by  $\theta(z, w) = (w + \bar{z})^N$ , for  $N \in \mathbb{N} \setminus \{0, 1\}$  and  $(z, w) \in \mathbb{C}^2$ .

**Question 10.** Let  $N \in \mathbb{N} \setminus \{0, 1\}$ ,  $A \in \mathbb{C} \setminus \{0\}$ ,  $(B_1, \dots, B_n) \in \mathbb{C}^n \setminus \{0\}$  and  $s \in [1, +\infty[$ . Let  $g, f_0, \dots, f_{N-2} : D \rightarrow \mathbb{C}$  be continuous functions, where  $D$  is a domain on  $\mathbb{C}^n$ . Define  $\psi(z, w) = |(Aw + B_1\bar{z}_1 + \dots + B_n\bar{z}_n)^N + g(z)|^s$  and  $\varphi(z, w) = |(Aw + B_1\bar{z}_1 + \dots + B_n\bar{z}_n)^N + f_{N-2}(z)w^{N-2} + \dots + f_0(z)|$ , for  $(z, w) = (z_1, \dots, z_n, w) \in D \times \mathbb{C}$ . Assume that  $\psi$  is psh on  $D \times \mathbb{C}$ . Prove that  $g = 0$  on  $D$ . Suppose that  $\varphi$  is psh on  $D \times \mathbb{C}$ . Prove that  $f_{N-2} = \dots = f_0 = 0$ .

## References

- [1] J. Abidi, *Sur quelques problèmes concernant les fonctions holomorphes et plurisousharmoniques*, Rend. Circ. Mat. Palermo 51 (2002) 411–424.
- [2] J. Abidi, *Contribution à l'étude des fonctions plurisousharmoniques convexes et analytiques*, Serdica Math. J. 40 (2014) 329–388.
- [3] J. Abidi, *The real and complex convexity*, J. Math. Appl. 41 (2018) 123–156.
- [4] J. Abidi, *On convex and analytic functions*, An. Univ. Oradea Fasc. Mat. XXVI (2019) 107–128.
- [5] M. Andersson, M. Passare, R. Sigurdsson, *Complex Convexity and Analytic Functionals*, Progress in Mathematics, Vol. 225, Birkhäuser, Berlin, 2004.
- [6] H.J. Bremermann, *Complex convexity*, Trans. Amer. Math. Soc. 82 (1956) 17–51.
- [7] U. Cegrell, *Removable singularities for plurisubharmonic functions and related problems*, Proc. Lond. Math. Soc. 36 (1978) 310–336.
- [8] H. El Mir, *Sur le prolongement des courants positifs fermés*, Acta Math. 153 (1984) 1–45.
- [9] G.M. Henkin, J. Leiterer, *Theory of Functions on Complex Manifolds*, Birkhäuser, Boston, Mass., 1984.
- [10] L. Hörmander, *An Introduction to Complex Analysis in Several Variables*, Third Edition (revised), Mathematical Library, Vol. 7, North Holland, Amsterdam-New York-Oxford-Tokyo, 1990.
- [11] L. Hörmander, L., *Notions of Convexity*, Birkhäuser, Basel-Boston-Berlin, 1994.

- [12] M. Jarnicki, P. Pflug, *Extension of Holomorphic Functions*, de Gruyter, Berlin, 2000.
- [13] M. Klimek, *Pluripotential Theory*, Clarendon Press, Oxford, 1991.
- [14] S.G. Krantz, *Function Theory of Several Complex Variables*, Wiley, New York, 1982.
- [15] P. Lelong, *Définition des fonctions plurisousharmoniques*, C. R. Acad. Sci. Paris 215 (1942) 398–400.
- [16] P. Lelong, *Sur les suites des fonctions plurisousharmoniques*, C. R. Acad. Sci. Paris 215 (1942) 454–456.
- [17] P. Lelong, *Les fonctions plurisousharmoniques*, Ann. Sci. Ecole Norm. Sup. 62 (1945) 301–338.
- [18] W. Rudin, *Function Theory in Polydiscs*, Benjamin, New York, 1969.
- [19] V.S. Vladimirov, *Les Fonctions de Plusieurs Variables Complexes (et Leur Application a la Théorie Quantique des Champs)*, Paris: Dunod, 1967.

**DOI: 10.7862/rf.2022.2**

**Jamel Abidi**

email: abidijame11@gmail.com

ORCID: 0000-0002-2552-6848

Département de Mathématiques

Faculté des Sciences de Tunis

1060 Tunis

TUNISIA

*Received 25.11.2021*

*Accepted 14.05.2022*