

# **Journal of Mathematics and Applications**

(e-ISSN 2300-9926)

vol. 37 (2014)

Issued with the consent of the Rector

**Editor in Chief**  
**Publishing House of Rzeszow University of Technology**  
Leonard ZIEMIAŃSKI

**Journal of Mathematics and Applications (JMA)** will publish carefully selected original research papers in any area of pure mathematics and its applications. Occasionally, the very authoritative expository survey articles of exceptional value can be published.

Manuscript, written in English and prepared using LaTeX, may be submitted in duplicate to the Editorial Office or one of the Editors or members of the Editorial Board. Electronic submission (of pdf or dvi file) is strongly preferred. Detailed information for authors is given on the inside back cover.

**Editor in Chief**  
**Journal of Mathematics and Applications**  
Jan STANKIEWICZ (Poland)

**Editorial Committee (Thematic editors)**  
Józef BANAS (Poland), Janusz SOKÓŁ (Poland)  
Bronisław Dov WAJNRYB (Poland), Iwona WŁOCH (Poland)

**Statistical editor**  
Mariusz STARTEK (Poland)

**Editorial assistant**  
Katarzyna WILCZEK (Poland)

**Members**  
Antoni PARDAŁA (Poland), Krzysztof PIEJKO (Poland)  
Beata RZEPKA (Poland)

**Language editors**  
Richard Fournier (Canada)

Text prepared to print in L<sup>A</sup>T<sub>E</sub>X  
by K. Wilczek

The printed version of JMA is an original version.

p-ISSN 1733-6775  
e-ISSN 2300-9926

Publisher: Publishing House of Rzeszow University of Technology,  
12 Powstańców Warszawy Ave., 35-959 Rzeszow (e-mail: [oficyna1@prz.edu.pl](mailto:oficyna1@prz.edu.pl))  
<http://oficyna.portal.prz.edu.pl/en/>  
Editorial Office: Rzeszow University of Technology, Department of Mathematics, P.O. BOX 85  
8 Powstańców Warszawy Ave., 35-959 Rzeszów (e-mail: [jma@prz.edu.pl](mailto:jma@prz.edu.pl))  
<http://jma.prz.edu.pl>

Additional information and an imprint – p. 133

# Journal of Mathematics and Applications

vol. 37 (2014)

## Table of contents

1. **M.K. Aouf, R.M. El-Ashwah, J.Dziok, J. Stankiewicz:** Partial sums of a certain harmonic univalent meromorphic functions .....5
2. **R. Borzooei, A. Borumand Saeid, R. Ameri, A. Rezaei:** Involutionary  $BE$ -algebras ..... 13
3. **M. K. Dubey, R. Anuradha:** On Generalised Quasi-ideals and Bi-ideals in Ternary Semigroups .....27
4. **J. Górowski, J. Żabowski:** Remarkable identities .....39
5. **V.K. Jain:** On the zeros of an analytic function ..... 51
6. **L. Koczan, P. Zaprawa:** On circularly symmetric functions ..... 59
7. **Z. Latreuch, B. Belaïdi:** Properties of higher order differential polynomials generated by solutions of complex differential equations in the unit disc .....67
8. **L. Rybarska-Rusinek, D. Jaworski, A. Linkov:** On efficient evaluation of integrals entering boundary equations of 3D potential and elasticity theory .....85
9. **T. M. Seoudy, M. K. Aouf:** On differential sandwich theorems of analytic functions defined by certain generalized linear operator ..... 97
10. **P. Sharma, R. K. Maurya:** Certain subordination results on the convolution of analytic functions ..... 111
11. **A. Zireh, S. A. Hosseini:** Some inequalities for the polar derivative of a polynomial with restricted zeros ..119

## Partial sums of a certain harmonic univalent meromorphic functions

M.K. Aouf, R.M. El-Ashwah,  
J.Dziok, J. Stankiewicz

ABSTRACT: In the present paper we determine sharp lower bounds of the real part of the ratios of harmonic univalent meromorphic functions to their sequences of partial sums.

Let  $\Sigma_H$  denote the class of functions  $f$  that are harmonic univalent and sense-preserving in  $U^* = \{z : |z| > 1\}$  which are of the form

$$f(z) = h(z) + \overline{g(z)},$$

where

$$h(z) = z + \sum_{n=1}^{\infty} a_n z^{-n}, \quad g(z) = \sum_{n=1}^{\infty} b_n z^{-n}.$$

Now, we define the sequences of partial sums of functions  $f$  of the form

$$\begin{aligned} f_s(z) &= z + \sum_{n=1}^s a_n z^{-n} + \overline{g(z)}, \\ \tilde{f}_r(z) &= g(z) + \sum_{n=1}^r \overline{b_n z^{-n}}, \\ f_{s,r}(z) &= z + \sum_{n=1}^s a_n z^{-n} + \sum_{n=1}^r \overline{b_n z^{-n}}. \end{aligned}$$

In the present paper we will determine sharp lower bounds for  $\operatorname{Re} \left\{ \frac{f(z)}{f_s(z)} \right\}$ ,  $\operatorname{Re} \left\{ \frac{f_s(z)}{f(z)} \right\}$ ,  $\operatorname{Re} \left\{ \frac{f(z)}{\tilde{f}_r(z)} \right\}$ ,  $\operatorname{Re} \left\{ \frac{\tilde{f}_r(z)}{f(z)} \right\}$ ,  $\operatorname{Re} \left\{ \frac{f(z)}{f_{s,r}(z)} \right\}$ ,  $\operatorname{Re} \left\{ \frac{f_{s,r}(z)}{f(z)} \right\}$ .

AMS Subject Classification: 30C45, 30C50

Keywords and Phrases: Harmonic function, meromorphic, univalent, sense-preserving.

## 1 Introduction

A continuous function  $f = u + iv$  is a complex valued harmonic function in a simply connected complex domain  $D \subset \mathbb{C}$  if both  $u$  and  $v$  are real harmonic in  $D$ . It was shown by Clunie and Sheil-Small [4] that such harmonic function can be represented by  $f = h + \bar{g}$ , where  $h$  and  $g$  are analytic in  $D$ . Also, a necessary and sufficient condition for  $f$  to be locally univalent and sense preserving in  $D$  is that  $|h'(z)| > |g'(z)|$ . There are numerous papers on univalent harmonic functions defined in a domain  $U = \{z \in \mathbb{C} : |z| < 1\}$  (see [6,7], [14] and [15]). Hergartner and Schober [10] investigated functions harmonic in the exterior of the unit disc i.e  $U^* = \{z \in \mathbb{C} : |z| > 1\}$ . They showed that a complex valued, harmonic, sense preserving univalent function  $f$ , defined on  $U^*$  and satisfying  $f(\infty) = \infty$  must admit the representation

$$f(z) = h(z) + \overline{g(z)} + A \log |z| \quad (A \in \mathbb{C}), \quad (1.1)$$

where

$$h(z) = \alpha z + \sum_{n=1}^{\infty} a_n z^{-n}, \quad g(z) = \beta z + \sum_{n=1}^{\infty} b_n z^{-n} \quad (z \in U^*, 0 \leq |\beta| < |\alpha|), \quad (1.2)$$

and  $a = \bar{f}_z/f_z$  is analytic and satisfy  $|a(z)| < 1$  for  $z \in U^*$ .

Let us denote by  $\Sigma_H$  the class of functions  $f$  that are harmonic univalent and sense-preserving in  $U^*$ , which are of the form

$$f(z) = h(z) + \overline{g(z)} \quad (z \in U^*), \quad (1.3)$$

where

$$h(z) = z + \sum_{n=1}^{\infty} a_n z^{-n}, \quad g(z) = \sum_{n=1}^{\infty} b_n z^{-n}.$$

Now, we introduce a class  $\Sigma_H(c_n, d_n, \delta)$  consisting of functions of the form (1.3) such that

$$\sum_{n=1}^{\infty} c_n |a_n| + \sum_{n=1}^{\infty} d_n |b_n| < \delta \quad (d_n \geq c_n \geq c_2 > 0; \delta > 0). \quad (1.4)$$

It is easy to see that various subclasses of  $\Sigma_H$  consisting of functions  $f(z)$  of the form (1.3) can be represented as  $\Sigma_H(c_n, d_n, \delta)$  for suitable choices of  $c_n, d_n$  and  $\delta$  studies earlier by various authors.

- $\Sigma_H(n, n, 1) = H_0^*$  (see Jahangiri and Silverman. [8]);
- $\Sigma_H(n + \gamma, n - \gamma, 1 - \gamma) = \Sigma_H^*(\gamma)$  ( $0 \leq \gamma < 1, n \geq 1$ ) (see Jahangiri [5]);

- $\Sigma_H(|(n+1)\lambda - 1|, |(n-1)\lambda + 1|, 1 - \alpha) = \Sigma_H R(\alpha, \lambda) (0 \leq \alpha < 1, \lambda \geq 0, n \geq 1)$  (see Ahuja and Jahangiri [1]);
- $\Sigma_H(n + \alpha - \alpha\lambda(n+1), n - \alpha - \alpha\lambda(n-1), 1 - \alpha) = \Sigma_H S^*(\alpha, \lambda) (0 \leq \alpha < 1, 0 \leq \lambda \leq 1, n \geq 1)$  (see Janteng and Halim [9]),
- $\Sigma_H(n(n+2)^m, n(n-2)^m, 1) = MH^*(m) (m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, \mathbb{N} = \{1, 2, \dots\}, n \geq 1)$  (see Bostanci and Ozturk [2]);
- $\Sigma_H((n+\gamma)(n+2)^m, (n-\gamma)(n-2)^m, 1 - \gamma) = MH^*(m, \gamma) (0 \leq \gamma < 1, m \in \mathbb{N}_0, n \geq 1)$  (see Bostanci and Ozturk [3]).

Silvia [17] studied the partial sums of the convex functions of order  $\alpha$ , later on Silverman [16] studied partial sum for starlike and convex functions. Very recently, Porwal [12], Porwal and Dixit [13] and Porwal [11] studied analogues interesting results on the partial sums of certain harmonic univalent functions.

Since to a certain extent the work in the harmonic univalent meromorphic functions case has paralleled that of the harmonic analytic univalent case, one is tempted to search results analogous to those of Porwal [11] for meromorphic harmonic univalent functions in  $U^*$ .

Now, we define the sequences of partial sums of functions  $f$  of the form (1.3) by

$$\begin{aligned} f_s(z) &= z + \sum_{n=1}^s a_n z^{-n} + \sum_{n=1}^{\infty} \overline{b_n z^{-n}}, \\ \tilde{f}_r(z) &= z + \sum_{n=1}^{\infty} a_n z^{-n} + \sum_{n=1}^r \overline{b_n z^{-n}}, \\ f_{s,r}(z) &= z + \sum_{n=1}^s a_n z^{-n} + \sum_{n=1}^r \overline{b_n z^{-n}}, \end{aligned} \tag{1.5}$$

when the coefficients of  $f$  are sufficiently small to satisfy the condition (1.4).

In the present paper, motivated essentially by the work of Silverman [16] and Porwal [11], we will determine sharp lower bounds for  $\operatorname{Re} \left\{ \frac{f(z)}{f_s(z)} \right\}$ ,

$$\operatorname{Re} \left\{ \frac{f_s(z)}{f(z)} \right\}, \operatorname{Re} \left\{ \frac{f(z)}{\tilde{f}_r(z)} \right\}, \operatorname{Re} \left\{ \frac{\tilde{f}_r(z)}{f(z)} \right\}, \operatorname{Re} \left\{ \frac{f(z)}{f_{s,r}(z)} \right\} \text{ and } \operatorname{Re} \left\{ \frac{f_{s,r}(z)}{f(z)} \right\}.$$

## 2 Main Results

**Theorem 1.** *Let  $s \in \mathbb{N}$  and let  $f(z) = h(z) + \overline{g(z)} \in \Sigma_H$ . Then*

$$(i) \quad \operatorname{Re} \left\{ \frac{f(z)}{f_s(z)} \right\} > 1 - \frac{\delta}{c_{s+1}} \quad (z \in U), \tag{2.1}$$

and

$$(ii) \quad \operatorname{Re} \left\{ \frac{f_s(z)}{f(z)} \right\} > \frac{c_{s+1}}{\delta + c_{s+1}} \quad (z \in U), \tag{2.2}$$

whenever

$$c_n \geq \begin{cases} \delta, & n = 2, 3, \dots, s, \\ c_{s+1}, & n = s+1, s+2, \dots \end{cases} \quad (2.3)$$

The estimates in (2.1) and (2.2) are sharp for the function given by

$$f(z) = z + \frac{\delta}{c_{s+1}} z^{-s-1} \quad (z \in U^*). \quad (2.4)$$

**Proof.** (i) To obtain the sharp lower bound given by (2.1), let us put

$$\begin{aligned} g_1(z) &= \frac{c_{s+1}}{\delta} \left\{ \frac{f(z)}{f_s(z)} - \left(1 - \frac{\delta}{c_{s+1}}\right) \right\} \\ &= 1 + \frac{\frac{c_{s+1}}{\delta} \sum_{n=s+1}^{\infty} a_n z^{-n}}{z + \sum_{n=1}^s a_n z^{-n} + \sum_{n=1}^{\infty} b_n z^{-n}}. \end{aligned} \quad (2.5)$$

Then, it is sufficient to show that  $\operatorname{Re} g_1(z) > 0$  ( $z \in U^*$ ) or equivalently

$$\left| \frac{g_1(z) - 1}{g_1(z) + 1} \right| \leq 1 \quad (z \in U^*).$$

Since

$$\left| \frac{g_1(z) - 1}{g_1(z) + 1} \right| \leq \frac{\frac{c_{s+1}}{\delta} \sum_{n=s+1}^{\infty} |a_n|}{2 - 2 \left( \sum_{n=1}^s |a_n| + \sum_{n=1}^{\infty} |b_n| \right) - \frac{c_{s+1}}{\delta} \sum_{n=s+1}^{\infty} |a_n|}, \quad (2.6)$$

the last expression is bounded above by 1, if and only if

$$\sum_{n=1}^s |a_n| + \sum_{n=1}^{\infty} |b_n| + \frac{c_{s+1}}{\delta} \sum_{n=s+1}^{\infty} |a_n| \leq 1. \quad (2.7)$$

Then, it is sufficient to show that L.H.S. of (2.7) is bounded above by

$$\sum_{n=1}^{\infty} \frac{c_n}{\delta} |a_n| + \sum_{n=1}^{\infty} \frac{d_n}{\delta} |b_n|,$$

which is equivalent to the true inequality

$$\sum_{n=1}^s \frac{c_n - \delta}{\delta} |a_n| + \sum_{n=1}^{\infty} \frac{d_n - \delta}{\delta} |b_n| + \sum_{n=s+1}^{\infty} \frac{c_n - c_{s+1}}{\delta} |a_n| \geq 0. \quad (2.8)$$

If we take

$$f(z) = z + \frac{\delta}{c_{s+1}} z^{-s-1}, \quad (2.9)$$

with  $z = re^{\frac{i\pi}{s+2}}$  and let  $r \rightarrow 1^+$ , we obtain

$$\frac{f(z)}{f_s(z)} = 1 + \frac{\delta z^{-s-2}}{c_{s+1}} \rightarrow 1 - \frac{\delta}{c_{s+1}},$$

which shows that the bound in (2.1) is best possible.

(ii) Similarly, if we put

$$\begin{aligned} g_2(z) &= \left( \frac{\delta + c_{s+1}}{\delta} \right) \left( \frac{f_s(z)}{f(z)} - \frac{c_{s+1}}{\delta + c_{s+1}} \right) \\ &= 1 - \frac{\left( \frac{\delta + c_{s+1}}{\delta} \right) \left( \sum_{n=s+1}^{\infty} |a_n| z^n + \sum_{n=1}^{\infty} \overline{b_n} z^{-n} \right)}{z + \sum_{n=1}^{\infty} |a_n| z^n + \sum_{n=1}^{\infty} \overline{b_n} z^{-n}}, \end{aligned}$$

and make use of (2.3), we can deduce that

$$\left| \frac{g_2(z) - 1}{g_2(z) + 1} \right| \leq \frac{\left( \frac{c_{s+1} + \delta}{\delta} \right) \left( \sum_{n=s+1}^{\infty} |a_n| + \sum_{n=1}^{\infty} |b_n| \right)}{2-2 \left( \sum_{n=1}^{\infty} |a_n| + \sum_{n=1}^{\infty} |b_n| \right) - \left( \frac{c_{s+1} - \delta}{\delta} \right) \left( \sum_{n=s+1}^{\infty} |a_n| + \sum_{n=1}^{\infty} |b_n| \right)}. \quad (2.10)$$

This last expression is bounded above by 1, if and only if

$$\sum_{n=1}^s |a_n| + \sum_{n=1}^{\infty} |b_n| + \left( \frac{c_{s+1}}{\delta} \right) \sum_{n=s+1}^{\infty} |a_n| \leq 1. \quad (2.11)$$

Since L.H.S. of (2.11) is bounded above by

$$\sum_{n=1}^{\infty} \frac{c_n}{\delta} |a_n| + \sum_{n=1}^{\infty} \frac{d_n}{\delta} |b_n|,$$

the bound in (2.2) follows and is sharp with the extremal function  $f(z)$  given by (2.4). The proof of Theorem 1 is now complete.

Employing the techniques used in Theorem 1, we can prove the following theorems.

**Theorem 2.** Let  $r \in \mathbb{N}$  and let  $f(z) = h(z) + \overline{g(z)} \in \Sigma_H$ . Then

$$(i) \quad \operatorname{Re} \left\{ \frac{f(z)}{\tilde{f}_r(z)} \right\} > 1 - \frac{\delta}{d_{r+1}} \quad (z \in U), \quad (2.12)$$

and

$$(ii) \quad \operatorname{Re} \left\{ \frac{\tilde{f}_r(z)}{f(z)} \right\} > \frac{d_{r+1}}{\delta + d_{r+1}} \quad (z \in U), \quad (2.13)$$



whenever

$$d_n \geq \begin{cases} \delta, & n = 2, 3, \dots, r, \\ d_{r+1}, & n = r + 1, r + 2, \dots \end{cases}$$

The estimates in (2.12) and (2.13) are sharp for the function given by

$$f(z) = z + \frac{\delta}{d_{r+1}} \bar{z}^{-r-1} \quad (z \in U^*). \quad (2.14)$$

**Theorem 3.** Let  $s, r \in \mathbb{N}$  and let  $f(z) = h(z) + \overline{g(z)} \in \Sigma_H$ . Then

$$(i) \quad \operatorname{Re} \left\{ \frac{f(z)}{f_{s,r}(z)} \right\} > 1 - \frac{\delta}{c_{s+1}} \quad (z \in U), \quad (2.15)$$

and

$$(ii) \quad \operatorname{Re} \left\{ \frac{f_{s,r}(z)}{f(z)} \right\} > \frac{c_{s+1}}{\delta + c_{s+1}} \quad (z \in U), \quad (2.16)$$

whenever

$$\begin{aligned} c_n &\geq \begin{cases} \delta, & n = 2, 3, \dots, s, \\ c_{s+1}, & n = s + 1, s + 2, \dots, \end{cases} \\ d_n &\geq \begin{cases} \delta, & n = 2, 3, \dots, s, \\ c_{s+1}, & n = s + 1, s + 2, \dots \end{cases} \end{aligned} \quad (2.17)$$

Also,

$$(i) \quad \operatorname{Re} \left\{ \frac{f(z)}{f_{s,r}(z)} \right\} > 1 - \frac{\delta}{d_{r+1}} \quad (z \in U), \quad (2.18)$$

and

$$(ii) \quad \operatorname{Re} \left\{ \frac{f_{s,r}(z)}{f(z)} \right\} > \frac{d_{r+1}}{\delta + d_{r+1}} \quad (z \in U), \quad (2.19)$$

whenever

$$\begin{aligned} c_n &\geq \begin{cases} \delta, & n = 2, 3, \dots, r, \\ d_{r+1}, & n = r + 1, r + 2, \dots, \end{cases} \\ d_n &\geq \begin{cases} \delta, & n = 2, 3, \dots, r, \\ d_{r+1}, & n = r + 1, r + 2, \dots \end{cases} \end{aligned} \quad (2.20)$$

The estimates in (2.15), (2.16), (2.18) and (2.19) respectively, are sharp for the function given by (2.4) and (2.14), respectively.

**Remark.** By specializing the coefficients  $c_n, d_n$  and the parameters  $\delta$  we obtain corresponding results for various subclasses mentioned in the introduction.

## References

- [1] O. P. Ahuja and J. M. Jahangiri, Certain meromorphic harmonic functions, Bull. Malaysian Math. Sci. Soc., 25(2002), 1-10.

- [2] H. Bostanci and M. Ozturk, A new subclass of the meromorphic harmonic starlike functions, *Appl. Math. Letters*, 23(2010), 1027-1032.
- [3] H. Bostanci and M. Ozturk, A new subclass of the meromorphic harmonic  $\gamma$ -starlike functions, *Appl. Math. Comput.*, 218(2011), 683-688..
- [4] J. Clunie and T. Sheil-Small, Harmonic univalent functions, *Ann. Acad. Sci. Fenn. Ser. A. I. Math.*, 9(1984), 3-25.
- [5] J. M. Jahangiri, Harmonic meromorphic starlike functions, *Bull. Korean Math. Soc.*, 37(2002), no.2, 291-301.
- [6] J. M. Jahangiri, Harmonic functions starlike in the unit disc, *J. Math. Anal. Appl.*, 235(1999), 470-477.
- [7] J. M. Jahangiri, Coefficient bounds and univalent criteria for harmonic functions with negative coefficients, *Ann. Univ. Marie-Curie Sklodowska Sect. A*, 52(1998), 57-66.
- [8] J. M. Jahangiri and H. Silverman, Meromorphic univalent harmonic function with negative coefficients, *Bull. Korean Math. Soc.*, 36(1999), no.4, 763-770.
- [9] A. Janteng and S. A. Halim, A subclass of harmonic meromorphic functions, *Int. J. Contemp. Math. Sci.*, 2(2007), no. 24, 1167-1174.
- [10] W. Hergartner and G. Schober, Univalent harmonic function, *Trans. Amer. Math. Soc.*, 299(1987), 1-31.
- [11] S. Porwal, A convolution approach on partial sums of certain harmonic univalent functions, *Internat. J. Math. Math. Sci.*, Vol. 2012, Art. ID 509349, 1-12.
- [12] S. Porwal, Partial sums of certain harmonic univalent function, *Lobachevskii J. Math.*, 32(2011), no.4, 366-375.
- [13] S. Porwal and K. K. Dixit, Partial sums of stalike harmonic univalent function, *Kungpook Math. J.*, 50(2010), no. 3, 433-445.
- [14] H. Silverman, Harmonic univalent function with negative coefficients, *J. Math. Anal. Appl.*, 220(1998), 283-289.
- [15] H. Silverman and E. M. Silvia, Subclasses of harmonic univalent functions, *New Zealand J. Math.*, 28(1999), 275-284.
- [16] H. Silverman, Partial sums of starlike and convex functions, *J. Math. Anal. Appl.*, 209(1997), 221-227.
- [17] E. M. Silvia, On partial sums of convex functions of order  $\alpha$ , *Houston J. Math.*, 11(1985), 397-404.

**DOI: 10.7862/rf.2014.1**

**M.K. Aouf**

email: mkaouf127@yahoo.com

Department of Mathematics  
Faculty of Science  
Mansoura University  
Mansoura 35516, Egypt

**R.M. El-Ashwah**

email: r\_elashwah@yahoo.com

Department of Mathematics  
Faculty of Science  
Damietta University  
New Damietta 34517, Egypt

**J. Dziok - corresponding author**

email: jdziok@univ.rzeszow.pl

Institute of Mathematics  
University of Rzeszow  
ul.Rejtana16A,  
35-310 Rzeszow, Poland

**J. Stankiewicz**

email: jstan@prz.rzeszow.pl

Department of Mathematics  
Technical University of Rzeszow  
ul.Wincentego Pola 2  
35-959 Rzeszow, Poland

*Received 8.04.2013, Accepted 26.09.2013*

## Involutory $BE$ -algebras

*R. Borzooei, A. Borumand Saeid,  
R. Ameri, A. Rezaei*

ABSTRACT: This paper is devoted to the study of some structural properties of bounded and involutory  $BE$ -algebras and investigate the relationship between them. We construct a commutative monoid by definition of proper operation in an involutory  $BE$ -algebra. Some rules of calculus for  $BE$ -algebras with a semi-lattice structure are provided. Many results related to the natural order of a  $BE$ -algebras were found. Finally, we show that an involutory bounded  $BE$ -algebra  $X$  is semi-simple.

AMS Subject Classification: *06F35, 03G25*

Keywords and Phrases: *(bounded, involutory)BE-algebra, involution, semi-lattice, lattice, semi-simple.*

### 1 Introduction and Preliminaries

The study of  $BCK/BCI$ -algebras was initiated by K. Iséki in 1966 as a generalization of propositional logic. There exist several generalization of  $BCK/BCI$ -algebras, such as  $BCH$ -algebras,  $d$ -algebras,  $B$ -algebras,  $BH$ -algebras, etc.

Especially, the notion of  $BE$ -algebras was introduced by H. S. Kim and Y. H. Kim [7], in which was deeply studied by S. S. Ahn and et. al., in [1, 2, 3], Walendziak in [15], A. Rezaei and et. al., in [12, 13, 14]. Lattice-valued logic is becoming a research field strongly influences the development of Algebraic Logic, Computer Science and Artificial Intelligence Technology.  $BE$ -algebras are important tools for certain investigations in algebraic logic since they can be consider as fragments of any propositional logic containing a logical connective implication and the constant 1 which is considered as the logical value "true". In this paper, we develop the theory  $BE$ -algebras with define a new structure as bounded and involutory  $BE$ -algebras and investigate the relationship between them and proved some theorems.

The paper has been organized in tree sections. In section 1, we give some definitions and some previous results and in section 2 we define bounded  $BE$ -algebras and define a congruence relation on this algebra with respect to a filter which this

congruence relation allowed us to define a quotient algebra is also a bounded  $BE$ -algebra. In section 3 we discuss on involutory  $BE$ -algebra because it is well known this structure has an important and vital role in investigating the structure of a logical system. Since quotient algebra is a basic tool for exploring the structures of algebras and there are close contacts among congruences and quotient algebras, we introduce a new congruence relation on  $X$  and construct quotient algebra via this congruence relation.

**Definition 1.1.** [7] An algebra  $(X; *, 1)$  of type  $(2, 0)$  is called a  $BE$ -algebra if following axioms hold:

- (BE1)  $x * x = 1$ ,
- (BE2)  $x * 1 = 1$ ,
- (BE3)  $1 * x = x$ ,
- (BE4)  $x * (y * z) = y * (x * z)$ , for all  $x, y, z \in X$ .

We introduce a relation " $\leq$ " on  $X$  by  $x \leq y$  if and only if  $x * y = 1$ .

**Proposition 1.2.** [7] Let  $X$  be a  $BE$ -algebra. Then

- (i)  $x * (y * x) = 1$ ,
- (ii)  $y * ((y * x) * x) = 1$ , for all  $x, y \in X$ .

From now on, in this paper  $X$  is a  $BE$ -algebra, unless otherwise is stated. A subset  $F$  of  $X$  is called a filter of  $X$  if (F1)  $1 \in F$  and (F2)  $x \in F$  and  $x * y \in F$  imply  $y \in F$ . We denote by  $F(X)$  the set of all filters of  $X$  and  $Max(X)$  the set of all maximal filters of  $X$ . Let  $A$  be a non-empty subset of  $X$ , then the set

$$\langle A \rangle = \bigcap \{G \in F(X) \mid A \subseteq G\}$$

is called the filter generated by  $A$ , written  $\langle A \rangle$ . If  $A = \{a\}$ , we will denote  $\langle \{a\} \rangle$ , briefly by  $\langle a \rangle$ , and we call it a principal filter of  $X$ . For  $F \in F(X)$  and  $a \in X$ , we denote by  $F_a$  the filter generated by  $F \cup \{a\}$ .  $X$  is said to be self distributive if  $x * (y * z) = (x * y) * (x * z)$ , for all  $x, y, z \in X$ , (Example 8., [7]).

In a self distributive  $BE$ -algebra  $X$ ,  $F_a = \{x \in X : a * x \in F\}$ , ([3]).  $X$  is said to be transitive if  $y * z \leq (x * y) * (x * z)$  for all  $x, y, z \in X$ , [1]. We say that  $X$  is commutative if  $(x * y) * y = (y * x) * x$ , for all  $x, y \in X$ . In [15], A. Walendziak, showed that every dual  $BCK$ -algebra is a  $BE$ -algebra and any commutative  $BE$ -algebra is a dual  $BCK$ -algebra.

We note that " $\leq$ " is reflexive by (BE1). If  $X$  is self distributive, then relation " $\leq$ " is a transitive order set on  $X$ . Because if  $x \leq y$  and  $y \leq z$ , then

$$x * z = 1 * (x * z) = (x * y) * (x * z) = x * (y * z) = x * 1 = 1$$

and so  $x \leq z$ . If  $X$  is commutative, then by Proposition 3.3, [15], relation " $\leq$ " is antisymmetric. Hence if  $X$  is a commutative self distributive  $BE$ -algebra, then " $\leq$ " is a partial order set on  $X$ , (Example 3.4., [3]). We show that if  $I$  be an obstinate ideal of a self distributive  $BE$ -algebra  $X$ , then  $(X/I; *, C_1)$  is also a  $BE$ -algebra, which is called to be the quotient algebra via  $I$ , and  $C_1 = I$ , (see Theorem 3.13, [12]).

**Proposition 1.3.** [12] *Let  $X$  be self distributive. If  $x \leq y$ , then*

- (i)  $z * x \leq z * y$  and  $y * z \leq x * z$ ,
- (ii)  $y * z \leq (z * x) * (y * x)$ , for all  $x, y, z \in X$ .

**Theorem 1.4.** [13] *A dual BCK-algebra  $X$  is commutative if and only if  $(X; \leq)$  is an upper semi-lattice with  $x \vee y = (y * x) * x$ , for all  $x, y \in X$ .*

**Proposition 1.5.** [13] *Let  $X$  be a commutative  $BE$ -algebra. Then*

- (i) for each  $a \in X$ , the mapping  $f_a : x \rightarrow x * a$  is an anti-tone involution on the section  $[a, 1]$ .
- (ii)  $(A, \leq)$  is a near-lattice with section anti-tone involutions and for every  $a \in X$ , the anti-tone involutions  $f_a$  on  $[a, 1]$  is given by  $f_a(x) = x * a$ .

**Theorem 1.6.** [15, 13] *Let  $X$  be commutative. Then it is a semi-lattice with respect to  $\vee$ .*

**Definition 1.7.** [4] *A filter  $F$  of  $X$  is called an obstinate filter if  $x, y \notin F$  imply  $x * y \in F$  and  $y * x \in F$ .*

**Theorem 1.8.** [5] *Let  $X$  be self distributive.  $F \in F(X)$  and  $F \neq X$ . Then the following are equivalent:*

- (i)  $F$  is an obstinate filter,
- (ii) if  $x \notin F$ , then  $x * y \in F$ , for all  $y \in F$ .

## 2 On Bounded $BE$ -algebras

**Definition 2.1.**  $X$  is called bounded if there exists the smallest element  $0$  of  $X$  (i.e.,  $0 * x = 1$ , for all  $x \in X$ ).

**Example 2.2.** (i). The interval  $[0, 1]$  of real numbers with the operation " $*$ " defined by

$$x * y = \min\{1 - x + y, 1\}, \text{ for all } x, y \in X$$

is a bounded  $BE$ -algebra.

(ii). Let  $(X; *, 1)$  be a  $BE$ -algebra,  $0 \notin X$  and  $\bar{X} = X \cup \{0\}$ . If we extensively define

$$0 * x = 0 * 0 = 1 \text{ and } x * 0 = 0 \text{ for all } x \in X.$$

Then  $(\bar{X}; *, 0, 1)$  is a bounded  $BE$ -algebra with 0 as the smallest element.

(iii). Let  $X := \{0, a, b, 1\}$  be a set with the following table.

*	0	a	b	1
0	1	1	1	1
a	a	1	b	1
b	b	a	1	1
1	0	a	b	1

Then  $(X; *, 0, 1)$  is a bounded  $BE$ -algebra with 0 as the smallest element.

(iv). Let  $X := \{0, a, b, c, 1\}$  be a set with the following table.

*	0	a	b	c	1
0	1	1	1	1	1
a	0	1	b	c	1
b	0	a	1	c	1
c	0	1	b	1	1
1	0	a	b	c	1

Then  $(X; *, 0, 1)$  is a bounded  $BE$ -algebra with 0 as the smallest element.

(v). Let  $(X; \vee, \wedge, \neg, 0, 1)$  be a Boolean-lattice. Then  $(X; *, 1)$  is a bounded  $BE$ -algebra, where operation " $*$ " is defined by  $x * y = (\neg x) \vee y$ , for all  $x, y \in X$ .

**Remark.** The following example shows that the bounded  $BE$ -algebra is not a dual  $BCK$ -algebra and Hilbert algebra in general (see Definition 2.3, [15] and Definition 3.1, [14]).

**Example 2.3.** Let  $X := \{0, a, b, 1\}$  be a set with the following table.

*	0	a	b	1
0	1	1	1	1
a	0	1	1	1
b	0	1	1	1
1	0	a	b	1

Then  $(X; *, 0, 1)$  is a bounded  $BE$ -algebra with 0 as the smallest element but it is not a dual  $BCK$ -algebra, Hilbert algebra. Because

$$a * b = b * a = 1 \quad \text{while} \quad a \neq b.$$

Also, it is not an implication algebra. Because

$$(a * b) * b = 1 * b = b \neq (b * a) * a = 1 * a = a.$$

Given a bounded  $BE$ -algebra  $X$  with 0 as the smallest element, we denote  $x * 0$  by  $Nx$ , then  $N$  can be regarded as a unary operation on  $X$ .

**Proposition 2.4.** Let  $X$  be bounded with the smallest element 0. Then the following hold:

- (i)  $N0 = 1$  and  $N1 = 0$ ,
- (ii)  $x \leq NNx$ ,
- (iii)  $x * Ny = y * Nx$ , for all  $x, y \in X$ .

*Proof.* (i). By (BE1) and (BE2) we have  $N0 = 0 * 0 = 1$  and  $N1 = 1 * 0 = 0$ .

(ii). Since  $x * (Nxx) = x * ((x * 0) * 0) = (x * 0) * (x * 0) = 1$ , then  $x \leq NNx$ .

(iii). By (BE4) we have  $x * Ny = x * (y * 0) = y * (x * 0) = y * Nx$ .  $\square$

**Proposition 2.5.** *Let  $X$  be a self distributive and bounded. Then*

- (i)  $y * x \leq Nx * Ny$ ,
- (ii)  $x \leq y$ , implies  $Ny \leq Nx$ , for all  $x, y \in X$ .

*Proof.* (i). We have

$$\begin{aligned}
 (y * x) * (Nx * Ny) &= Nx * ((y * x) * Ny) &= (x * 0) * ((y * x) * (y * 0)) \\
 & &= (x * 0) * (y * (x * 0)) \\
 & &= y * ((x * 0) * (x * 0)) \\
 & &= y * 1 = 1.
 \end{aligned}$$

Hence  $y * x \leq Nx * Ny$ .

(ii). By (BE3) and assumption we have

$$\begin{aligned}
 Ny * Nx &= (y * 0) * (x * 0) &= (y * 0) * (1 * (x * 0)) \\
 & &= (y * 0) * ((x * y) * (x * 0)) \\
 & &= (y * 0) * (x * (y * 0)) \\
 & &= x * ((y * 0) * (y * 0)) \\
 & &= x * 1 = 1.
 \end{aligned}$$

Hence  $Ny \leq Nx$ .  $\square$

In the following example we show that the self-distributivity condition in the above theorem is necessary.

**Example 2.6.** *Example 2.2(iii), is a bounded BE-algebra with 0 as the smallest element, while it is not self-distributive. Because*

$$b * (0 * a) = 2 * 1 = 1 \neq (b * 0) * (b * a) = b * a = a.$$

We can seen easily that,  $b = a * b \not\leq Nb * Na = b * a = a$ .

**Proposition 2.7.** *Let  $X$  be bounded implicative self distributive. Then the following hold:*

- (i)  $X$  is commutative,



- (ii)  $x = Nx * x$ ,  
 (iii)  $x \vee y = y \vee x = Nx * y$ , for all  $x, y \in X$ .

*Proof.* (i). See proof of Theorem 3.12([14]).

(ii). Assume that  $X$  is a bounded implicative. Then  $Nx * x = (x * 0) * x = x$ .

(iii). Let  $X$  be bounded implicative self distributive and  $x, y \in X$ . then by Proposition 1.3,  $0 \leq y$  and  $x * 0 \leq x * y$ . Furthermore, by Propositions 1.2 and 1.3, we get

$$x \leq (x * y) * y \leq (x * 0) * y = Nx * y$$

Since by Proposition 1.2,  $y \leq Nx * y$ , then  $Nx * y$  is an upper bound of  $x$  and  $y$ . Hence  $x \vee y \leq Nx * y$ . Also, we have

$$Nx * y \leq (y * x) * (Nx * x) = (y * x) * x.$$

Since  $X$  is commutative, then by Theorem 1.6, we have  $(y * x) * x = x \vee y = y \vee x$  and so by Proposition 3.3([15]), the proof is complete.  $\square$

**Corollary 2.8.** *Let  $X$  be self distributive,  $F \in F(X)$  and  $F \neq X$ . Then the following are equivalent:*

- (i)  $F$  is an obstinate filter,  
 (ii) if  $x \notin F$ , then  $Nx \in F$ .

**Definition 2.9.** *Let  $X$  and  $Y$  be bounded. A homomorphism from  $X$  to  $Y$  is a function  $f : X \rightarrow Y$  such that*

- (i)  $f(x * y) = f(x) * f(y)$ ,  
 (ii)  $f(Nx) = N(f(x))$ ,  
 (iii)  $f(0) = 0$ , for all  $x, y \in X$ .

**Example 2.10.** *Consider  $X$  as Example 2.2(iii) and  $Y$  as Example 2.3. Define  $f : X \rightarrow Y$  such that  $f(1) = f(a) = f(b) = 1$  and  $f(0) = 0$ . Then  $f$  is a homomorphism.*

**Theorem 2.11.** *Let  $f : X \rightarrow Y$  be a homomorphism. Then  $\ker(f) = \{x \in X : f(x) = 1\}$  is a filter in  $X$ . Moreover, if  $f(x) = f(y)$ , then  $x * y \in \ker(f)$  and  $y * x \in \ker(f)$ , for all  $x, y \in X$ . If  $Y$  is commutative, then the converse is valid.*

*Proof.* We have  $f(1) = f(x * x) = f(x) * f(x) = 1$ . Hence  $1 \in \ker(f)$ . Now, let  $x \in \ker(f)$  and  $x * y \in \ker(f)$ . Then  $f(x) = f(x * y) = 1$ . But  $f(x * y) = f(x) * f(y) = 1$ . Hence  $f(y) = 1 * f(y) = 1$ . Therefore,  $y \in \ker(f)$ .

Now, let  $f(x) = f(y)$ . By using (BE1),  $f(x) * f(y) = 1$  and  $f(y) * f(x) = 1$ . But  $1 = f(x) * f(y) = f(x * y)$  and  $1 = f(y) * f(x) = f(y * x)$  implies  $x * y \in \ker(f)$  and  $y * x \in \ker(f)$ .

Assume that  $Y$  is commutative,  $x * y \in \ker(f)$  and  $y * x \in \ker(f)$ . Then  $f(x * y) = f(y * x) = 1$  which implies that  $f(x) * f(y) = f(y) * f(x) = 1$ . Hence by Proposition 3.3([15]),  $f(x) = f(y)$ .  $\square$

**Theorem 2.12.** *Let  $X$  be bounded transitive,  $F$  be a filter and  $X/F$  be the corresponding quotient algebra. Then the map  $f : X \rightarrow X/F$  which is defined by  $f(a) = [a]$ , for all  $a \in X$ , is a homomorphism and  $\ker(f) = F$ .*

*Proof.* By Propositions 5.4 and 5.7([11]),  $X/F$  is a quotient BE-algebra. Now, we have  $f(0) = [0]$  and

$$f(Nx) = f(x * 0) = f(x) * f(0) = f(x) * [0] = N(f(x)).$$

Now, let  $x \in \ker(f)$ . Then  $f(x) = [x] = [1]$  if and only if  $1 = x * 1 \in F$  and  $x = 1 * x \in F$  if and only if  $1 \in F$  and  $x \in F$ . Therefore,  $\ker(f) = F$ .  $\square$

### 3 Involutory BE-algebras

If  $NNx = x$ , then  $x$  is called an involution of  $X$ . The smallest element 0 and the greatest element 1 are two involutions of  $X$ , because

$$NN0 = N(0 * 0) = N1 = 1 * 0 = 0,$$

$$NN1 = N(1 * 0) = N0 = 0 * 0 = 1.$$

**Definition 3.1.** *A bounded BE-algebra  $X$  is called involutory if any element of  $X$  is involution.*

**Example 3.2.** (i). *Examples 2.2(i), (iii), (v), are involutory.*

(ii). *Let  $X := \{0, a, b, 1\}$  be a set with the following table.*

$*$	0	a	b	1
0	1	1	1	1
a	0	1	b	1
b	0	a	1	1
1	0	a	b	1

*Then  $(X; *, 0, 1)$  is a bounded BE-algebra but it is not an involutory. Because*

$$NNb = N(b * 0) = N0 = 0 * 0 = 1 \neq b.$$

(iii). *Let  $X := \{0, a, b, 1\}$  be a set with the following table.*

$*$	0	a	b	1
0	1	1	1	1
a	b	1	1	1
b	a	1	1	1
1	0	a	b	1

*Then  $(X; *, 0, 1)$  is an involutory BE-algebra but it is not an involutory dual BCK-algebra and involutory Hilbert algebra. Because*

$$a * b = 1 \text{ and } b * a = 1 \text{ while, } a \neq b.$$

Also, it is not an involutory implication algebra. Because

$$(a * b) * b = 1 * b = b \neq (b * a) * a = 1 * a = a.$$

**Proposition 3.3.** *If  $X$  is a bounded commutative, then  $X$  is an involutory.*

*Proof.* By using the commutativity we get

$$NNx = (x * 0) * 0 = (0 * x) * x = 1 * x = x.$$

Hence  $X$  is an involutory. □

In the following example we show that the commutativity condition in the above theorem is necessary.

**Example 3.4.** *Example 3.2(ii), is not an involutory. Because it is not commutative.*

**Proposition 3.5.** *If  $X$  is an involutory, then*

- (i)  $x * y = Ny * Nx$ ,
- (ii)  $x \leq Ny$  implies  $y \leq Nx$ , for all  $x, y \in X$ .

*Proof.* (i). Since  $X$  is an involutory, then we have  $NNx = x$ , for all  $x, y \in X$ . Hence by Proposition 2.4(iii),  $x * y = x * NNy = Ny * Nx$ .

(ii). Since  $x \leq Ny$ , we get  $x * Ny = 1$ . Hence by Proposition 2.4(iii),  $1 = x * Ny = y * Nx$ . So,  $y \leq Nx$ . □

**Lemma 3.6.** *Let  $X$  be bounded self distributive and  $x, y \in X$ .*

- (i) *if the smallest upper bound  $x \vee y$  of  $x$  and  $y$  exists, then the greatest lower bound  $Nx \wedge Ny$  of  $Nx$  and  $Ny$  exists and  $Nx \wedge Ny = N(x \vee y)$ .*
- (ii) *if  $X$  is involutory and the greatest lower bound  $x \wedge y$  exists, then the least upper bound  $Nx \vee Ny$  exists and  $Nx \vee Ny = N(x \wedge y)$ .*

*Proof.* (i). Assume that the smallest upper bound  $x \vee y$  of  $x$  and  $y$  exists. Since  $x \leq x \vee y$ , then by Proposition 1.3,  $(x \vee y) * 0 \leq x * 0$ , (i.e.,  $N(x \vee y) \leq Nx$ ). By the similar way  $N(x \vee y) \leq Ny$ . Hence  $N(x \vee y)$  is a lower bound of  $Nx$  and  $Ny$ . Also, assume that  $u$  is any lower bound of  $Nx$  and  $Ny$ . Then  $u \leq Nx$  and  $u \leq Ny$ . Hence by (BE4), we have  $x * (u * 0) = u * (x * 0) = u * Nx = 1$ . Hence  $x \leq Nu$  and by the similar way  $y \leq Nu$ . So,  $x \vee y \leq Nu$ . Now, by (BE4), we have  $(x \vee y) * (u * 0) = u * ((x \vee y) * 0) = 1$ . So,  $u \leq N(x \vee y)$ . Hence  $N(x \vee y)$  is a greatest lower bound of  $Nx$  and  $Ny$ . Therefore, the greatest lower bound  $Nx \wedge Ny$  of  $Nx$  and  $Ny$  exists, and  $Nx \wedge Ny = N(x \vee y)$ .

(ii). Assume that  $x \wedge y$  exists. Since  $x \wedge y \leq x$  and  $x \wedge y \leq y$ , then by Proposition 2.5, we have  $N(x) \leq N(x \wedge y)$  and  $N(y) \leq N(x \wedge y)$ . Hence  $N(x \wedge y)$  is an upper bound of  $Nx$  and  $Ny$ . Also, let  $u$  be any upper bound of  $Nx$  and  $Ny$ . Then  $Nx \leq u$  and  $Ny \leq u$ . Since  $X$  is involutory, then by Proposition 2.5, we derive  $Nu \leq NNx = x$  and  $Nu \leq NNy = y$ . So,  $Nu \leq x \wedge y$ . By Proposition 2.5, we have  $N(x \wedge y) \leq NNu = u$ . Hence  $N(x \wedge y)$  is the smallest upper bound of  $Nx$  and  $Ny$ . Then the least upper bound  $Nx \vee Ny$  exists, and  $Nx \vee Ny = N(x \wedge y)$ . □

**Theorem 3.7.** *Let  $X$  be involutory self distributive. Then the following are equivalent:*

- (i)  $(X; \leq)$  is an upper semi-lattice,
- (ii)  $(X; \leq)$  is a lower semi-lattice,
- (iii)  $(X; \leq)$  is a lattice.

Moreover, if  $(X; \leq)$  is a lattice, then the following identities hold:

$$x \wedge y = N(Nx \vee Ny) \text{ and } x \vee y = N(Nx \wedge Ny).$$

*Proof.* (i)  $\Rightarrow$  (ii). Since  $(X; \leq)$  is an upper semi-lattice, then  $Nx \vee Ny$  exists for all  $x, y \in X$ . By the first half part of Lemma 3.6,  $NNx \wedge NNy$  exists. Also, since  $X$  is involutory, we have  $NNx \wedge NNy = x \wedge y$ . Then  $x \wedge y$  exists. So,  $(X; \leq)$  is a lower semi-lattice.

(ii)  $\Rightarrow$  (iii). Since  $(X; \leq)$  is a lower semi-lattice,  $Nx \wedge Ny$  exists and using the second half part of Lemma 3.6,  $NNx \vee NNy$  exists, for all  $x, y \in X$ . Also, since  $X$  is involutory, we have  $NNx \vee NNy = x \vee y$ . Then  $x \vee y$  exists. So,  $(X; \leq)$  is an upper semi-lattice.

(iii)  $\Rightarrow$  (i). The proof is obvious.

Now, let  $(X; \leq)$  is a lattice. Since by Lemma 3.6,  $X$  is involutory, then we have

$$x \wedge y = NNx \wedge NNy = N(Nx \vee Ny),$$

$$x \vee y = NNx \vee NNy = N(Nx \wedge Ny).$$

□

**Theorem 3.8.** *Let  $I$  be an obstinate ideal of involutory(bounded) self-distributive  $X$ . Then  $(X/I; *, C_1)$  is involutory(bounded) self-distributive, too.*

*Proof.* By Theorems 3.13 and 3.16([12]),  $(X/I; *, C_1)$  is a self-distributive  $BE$ -algebra. Let  $x \in X$ . Then  $C_0 * C_x = C_{0*x} = C_1$ . Hence  $X/I$  is a bounded  $BE$ -algebra. Now,

$$NNC_x = (C_x * C_0) * C_0 = C_{x*0} * C_0 = C_{(x*0)*0} = C_{NNx} = C_x.$$

Therefore,  $X/I$  is an involutory  $BE$ -algebra. □

**Proposition 3.9.** *Let  $X$  be involutory and operation "  $\circ$  " is defined on  $X$  by  $x \circ y = Nx * y$ , for all  $x, y \in X$ . Then  $(X; \circ, 0)$  is a commutative monoid.*

*Proof.* By Proposition 2.4(iii),

$$x \circ y = Nx * y = Nx * NNy = Ny * NNx = Ny * x = y \circ x$$

and so  $X$  is commutative. Now, by Proposition 2.4(iii), and (BE4) we have

$$\begin{aligned}
 x \circ (y \circ z) = Nx * (y \circ z) &= Nx * (z \circ y) \\
 &= Nx * (Nz * y) \\
 &= Nz * (Nx * y) \\
 &= z \circ (Nx * y) \\
 &= (Nx * y) \circ z \\
 &= (x \circ y) \circ z.
 \end{aligned}$$

Hence " $\circ$ " is associative operation on  $X$ . Moreover, for any  $x \in X$

$$x \circ 0 = Nx * 0 = NNx = x \quad \text{and} \quad 0 \circ x = N0 * x = 1 * x = x.$$

□

In the following example we show that the converse of the Proposition 3.9, is not valid in general.

**Example 3.10.** Let  $X := \{0, a, b, 1\}$  be a set with the following table.

$*$	1	$a$
1	1	$a$
$a$	$a$	$a$

Then  $(X; *, 1)$  is a commutative monoid, but it is not a BE-algebra. Because  $a * a = a \neq 1$  and  $a * 1 = a \neq 1$ , (i.e., conditions (BE1) and (BE2) are not holds).

**Lemma 3.11.** Let  $X$  be bounded. Then

- (i) filter  $F$  of  $X$  is proper if and only if  $0 \notin F$ .
- (ii) each proper filter  $F$  is contained in a maximal filter.

*Proof.* (i). Let  $F$  be a proper filter of  $X$  and  $0 \in F$ . If  $x \in X$ , since  $0 * x = 1 \in F$ , which implies  $x \in F$ . Hence  $X = F$ , which is a contradiction. The converse is clear.

(ii). The proof is obvious. □

**Theorem 3.12.** Every bounded BE-algebra contains at least one maximal filter.

*Proof.* Let  $X$  be a bounded BE-algebra. Since  $\{1\}$  is a proper filter of  $X$ , then the proof is clear by Lemma 3.11. □

**Definition 3.13.** Let  $X$  be bounded. Then the radical of  $X$ , written  $Rad(X)$ , is defined by

$$Rad(X) = \cap \{F : F \in Max(X)\}.$$

In view of Theorem 3.12,  $Rad(X)$  always exists for a bounded algebra  $X$ . Following a standard terminology in the contemporary algebra, we shall call an algebra  $X$  semi-simple if  $Rad(X) = \{1\}$ .

**Example 3.14.** In Example 2.2(iv),  $F_1 = \{1\}$ ,  $F_2 = \{1, a\}$ ,  $F_3 = \{1, a, b, c\}$  and  $X$  are filters in  $X$  and  $F_3$  is only maximal filter of  $X$ . Hence  $\text{Rad}(X) = F_3$ .

**Example 3.15.** In Example 2.2(iii),  $F_1 = \{1\}$ ,  $F_2 = \{1, a\}$ ,  $F_3 = \{1, b\}$  and  $X$  are filters in  $X$  and  $F_2, F_3$  are maximal filters of  $X$ , also  $F_2 \cap F_3 = \{1\}$ . Hence  $\text{Rad}(X) = \{1\}$  and therefore  $X$  is semi-simple.

**Lemma 3.16.** Let  $X$  be an involutory bounded BE-algebra. Then for every  $x \in X$  with  $x \neq 1$ , there exists a maximal filter  $F$  of  $X$  such that  $x \notin F$ .

*Proof.* Let  $1 \neq x \in X$ . We claim that  $\langle Nx \rangle$  is a proper filter of  $X$ . By contrary, if it is not, then  $\langle Nx \rangle = X$ . Hence  $0 \in \langle Nx \rangle$  and therefore  $Nx * 0 = NNx = 1$ . Since  $X$  is involutory, then  $x = NNx = 1$ , which is a contradiction. By Lemma 3.11(ii), there is a maximal filter  $F$  of  $X$  such that  $\langle Nx \rangle \subseteq F$ , and  $x \notin F$ . Suppose  $x \in F$ . Since  $Nx = x * 0 \in F$ , then  $0 \in F$ , which is contrary by Lemma 3.11(i).  $\square$

**Theorem 3.17.** Let  $X$  be involutory and bounded. Then  $X$  is a semi-simple.

*Proof.* By Lemma 3.16, the proof is clear.  $\square$

In this section we define a congruence relation " $\theta$ " on involutory bounded BE-algebra  $X$  and construct quotient algebra  $(X/\theta; *, \theta_0, \theta_1)$  induced by the congruence relation " $\theta$ ", where, we denote  $\theta_x$  for the equivalence class  $[x]$  containing  $x$ . Since " $\theta$ " is a congruence on  $X$ , then the operation " $*$ " on  $X/\theta$  given by  $\theta_x * \theta_y = \theta_{x*y}$  is well-defined, because " $\theta$ " satisfied of the substitution property. Then  $(X/\theta; *, \theta_0, \theta_1)$  is an algebra of type  $(2, 0, 0)$  where,

$$\theta_0 = \{x : N0 = Nx\} = \{x : Nx = 1\}$$

is the zero equivalence class containing 0 and

$$\theta_1 = \{x : N1 = Nx\} = \{x : Nx = 0\}$$

is the one equivalence class containing 1. Now, in the following theorem define and prove this results.

**Theorem 3.18.** Let  $X$  be involutory and bounded. The relation " $\theta$ " defined on  $X$  by:

$$(x, y) \in \theta \text{ if and only if } Nx = Ny$$

is a congruence relation on  $X$  and the quotient algebra  $(X/\theta; *, \theta_0, \theta_1)$  is an involutory bounded BE-algebra.

*Proof.* It is clear that " $\theta$ " is an equivalence relation on  $X$ . Now, Let  $(x, y) \in \theta$  and  $(u, v) \in \theta$ . Then  $Nx = Ny$  and  $Nu = Nv$ . Hence,

$$Nx * Nu = (x * 0) * (u * 0) = u * ((x * 0) * 0) = u * NNx = u * x.$$

Thus  $N(u * x) = N(Nx * Nu) = N(Ny * Nv) = N(v * y)$ , and so  $(u * x, v * y) \in \theta$ . By the similarly way we have  $(x * u, y * v) \in \theta$ . Hence " $\theta$ " is a congruence relation on  $X$ .

Let  $\theta_x, \theta_y, \theta_z \in X/\theta$ . Then

$$(BE1) \quad \theta_x * \theta_x = \theta_{x*x} = \theta_1,$$

$$(BE2) \quad \theta_x * \theta_1 = \theta_{x*1} = \theta_1,$$

$$(BE3) \quad \theta_1 * \theta_x = \theta_{1*x} = \theta_x,$$

$$(BE4) \quad \theta_x * (\theta_y * \theta_z) = \theta_x * \theta_{y*z} = \theta_{x*(y*z)} = \theta_{y*(x*z)} = \theta_y * \theta_{x*z} = \theta_y * (\theta_x * \theta_z).$$

Now, since  $\theta_0 * \theta_x = \theta_{0*x} = \theta_1$ . Hence  $\theta_0$  is as the smallest element of  $X/\theta$ . Also,

$$NN\theta_x = (\theta_x * \theta_0) * \theta_0 = \theta_{x*0} * \theta_0 = \theta_{(x*0)*0} = \theta_{NNx} = \theta_x.$$

Therefore,  $(X/\theta; *, \theta_0, \theta_1)$  is an involutory bounded  $BE$ -algebra.  $\square$

**Example 3.19.** Consider Example 2.2(iv),  $\theta_0 = \{0\}$  and  $\theta_a = \theta_b = \theta_c = \theta_1 = \{a, b, c, 1\}$ . Then  $X/\theta = \{\theta_0, \theta_1\}$ . Thus  $(X/\theta; *, \theta_0, \theta_1)$  is an involutory bounded  $BE$ -algebra.

**Corollary 3.20.** Let  $X$  be an involutory bounded  $BE$ -algebra and  $X_0 := \{Nx : x \in X\}$ . Then  $(X_0; *, N0)$  is a  $BE$ -algebra.


**Example 3.21.** In Example 2.2(ii), (iv), respectively,  $X_0 = \{0, 1\}$  and  $X_0 = \{0, a, b, 1\}$ .

**Proposition 3.22.** Let  $X$  be involutory, bounded and self-distributive(commutative). Then  $X/\theta$  is involutory, bounded and self-distributive(commutative), too.

## 4 Conclusion and future research

In this paper, we introduced the notion of bounded and involutory  $BE$ -algebras and get some results. In addition, we have defined a congruence relation on involutory bounded  $BE$ -algebras and construct the quotient  $BE$ -algebra via this relations. In [10], J. Meng proved that implication algebras are dual to implicative  $BCK$ -algebras. Also R. Halaš in [9], showed commutative *Hilbert* algebras are implication algebras and A. Digo in [6], proved implication algebras are *Hilbert* algebras. Recently, A. Walendziak in [15], showed that an implication algebra is a  $BE$ -algebra and commutative  $BE$ -algebras are dual  $BCK$ -algebras. In [14], we showed that every *Hilbert* algebra is a self distributive  $BE$ -algebra and commutative self distributive  $BE$ -algebra is a *Hilbert* algebra. Then in the following diagram we summarize the results of this paper and we give the relations among such structures of involutory algebras.

" $A \rightarrow B$ ," means that  $A$  conclude  $B$ .



We think such results are very useful for study in this structure. In the future work we try assemble of calculus relative to different kinds of  $BE$ -algebras, as example, latticical structure and Boolean lattices.

**Acknowledgments:** The authors wish to thank the reviewers for their excellent suggestions that have been incorporated into this paper.

## References

- [1] S. S. AHN AND Y. H. SO, *On ideals and upper sets in  $BE$ -algebras*, Sci. Math. Jpn. 68, No. 2(2008), 279-285.
- [2] S. S. AHN AND K. S. SO, *On generalized upper sets in  $BE$ -algebras*, Bull. Korean Math. Soc. 46, No. 2(2009), 281-287.
- [3] S. S. AHN, Y. H. KIM AND J. M. KO, *Filters in commutative  $BE$ -algebras*, Bull. Korean Math. Soc. 27, No. 2(2012), 233-242.
- [4] A. BORUMAND SAEID, A. REZAEI AND R. A. BORZOOEI, *Some types of filters in  $BE$ -algebras*, Math. Comput. Sci. Vol. 6, No 3 (2013), 341-352.
- [5] R. A. BORZOOEI, A. BORUMAND SAEID, R. AMERI AND A. REZAEI, *States in  $BE$ -algebras*, Kochi Journal of Mathematics, Kochi Journal of Mathematics, 9 (2014), 27-42.
- [6] A. DIEGO, *Sur algébra de Hilbert*, Logique Math. Ser. A, 21(1967), 177-198.
- [7] H. S. KIM AND Y. H. KIM, *On  $BE$ -algebras*, Sci, Math, Jpn. 66, No. 1(2007), 113-116.
- [8] Y. HUANG, *BCI-algebra*, Science Press, Beijing 100717, China, (2006).
- [9] R. HALAŠ, *Remarks on commutative Hilbert algebras*, Mathematica Bohemica, 127, No. 4(2002), 525-529.
- [10] J. MENG, *Implication algebras are dual to Implicative BCK-algebras*, Soochow Journal of Math. V. 22, No. 4(1996), 567-571.
- [11] B. L. MENG, *On filters in  $BE$ -algebras*, Sci. Math. Jpn. 71(2010), 201-207.
- [12] A. REZAEI AND A. BORUMAND SAEID, *Some results in  $BE$ -algebras*, Analele Universitatii Oradea Fasc. Matematica, Tom XIX (2012), 33-44.
- [13] A. REZAEI AND A. BORUMAND SAEID, *Commutative ideals in  $BE$ -algebras*, Kyungpook Math. J. 52(2012), 483-494.
- [14] A. REZAEI, A. BORUMAND SAEID AND R. A. BORZOOEI, *Relation between Hilbert algebras and  $BE$ -algebras*, Appl. Math. 8. No 1 (2013), 573-574.



- [15] A. WALENDZIAK, *On commutative BE-algebras*, Sci. Math. Jpn. 69, No. 2(2008), 585-588.

**DOI: 10.7862/rf.2014.2**

**Rajab Ali Borzooei**

email: [borzooei@sbu.ac.ir](mailto:borzooei@sbu.ac.ir)

Dept. of Math.

Shahid Beheshti University of Tehran,

Tehran, Iran

**Arsham Borumand Saeid - corresponding author**

email: [rAmeri@ut.ac.ir](mailto:rAmeri@ut.ac.ir)

Dept. of Math.

Tehran University

Tehran, Iran

**Reza Ameri**

email: [arsham@uk.ac.ir](mailto:arsham@uk.ac.ir)

Dept. of Math.

Shahid Bahonar University of Kerman

Kerman, Iran

**Akbar Rezaei**

[mailRezaei@pnu.ac.ir](mailto:mailRezaei@pnu.ac.ir)

Dept. of Math.

Payame Noor University

p. o. box. 19395-3697, Tehran, Iran

*Received 13.12.2013, Accepted 12.02.2014*

# On Generalised Quasi-ideals and Bi-ideals in Ternary Semigroups

*Manish Kant Dubey, Rohatgi Anuradha*

ABSTRACT: In this paper, we introduce the notions of generalised quasi-ideals and generalised bi-ideals in a ternary semigroup. We also characterised these notions in terms of minimal quasi-ideals and minimal bi-ideals in a ternary semigroup.

AMS Subject Classification: *16Y30, 16Y60*

Keywords and Phrases: *Ternary Semigroup, Quasi-ideals, Bi-ideals, Minimal Quasi-ideals, Minimal Bi-ideals*

## 1 Introduction and Preliminaries

Good and Hughes[5] introduced the notion of bi-ideals and Steinfeld [2] introduced the notion of quasi-ideals in semigroups. In [1], Sioson studied the concept of quasi-ideals in ternary semigroups. He also introduced the notion of regular ternary semigroups and characterised them by using the notion of quasi-ideals. In [7], Dixit and Dewan studied about the quasi-ideals and bi-ideals of ternary semigroups.

A nonempty set  $S$  with a ternary operation  $S \times S \times S \mapsto S$ , written as  $(x_1, x_2, x_3) \mapsto [x_1x_2x_3]$  is called a ternary semigroup if it satisfies the following associative law:  $[[x_1x_2x_3]x_4x_5] = [x_1[x_2x_3x_4]x_5] = [x_1x_2[x_3x_4x_5]]$  for any  $x_1, x_2, x_3, x_4, x_5 \in S$ . In this paper, we denote  $[x_1x_2x_3]$  by  $x_1x_2x_3$ .

A non-empty subset  $T$  of a ternary semigroup  $S$  is called a ternary subsemigroup if  $t_1t_2t_3 \in T$  for all  $t_1, t_2, t_3 \in T$ . A ternary subsemigroup  $I$  of a ternary semigroup  $S$  is called a left ideal of  $S$  if  $SSI \subseteq I$ , a lateral ideal if  $SIS \subseteq I$ , a right ideal of  $S$  if  $ISS \subseteq I$ , a two-sided ideal of  $S$  if  $I$  is both left and right ideal of  $S$ , and an ideal of  $S$  if  $I$  is a left, a right and a lateral ideal of  $S$ . An ideal  $I$  of a ternary semigroup  $S$  is called a proper ideal if  $I \neq S$ . Let  $S$  be a ternary semigroup. If there exists an element  $0 \in S$  such that  $0xy = x0y = xy0 = 0$  for all  $x, y \in S$ , then "0" is called the zero element or simply the zero of the ternary semigroup  $S$ . In this case  $S \cup \{0\}$  becomes a ternary semigroup with zero. For example, the set of all non-positive integers  $Z_0^{-1}$  forms a ternary semigroup with usual ternary multiplication

and '0' forms a ternary semigroup with zero element and also the zero element satisfy  $(SS)^0S = S^0SS^0 = S(SS)^0 = S$ . Throughout this paper  $S$  will always denote a ternary semigroup with zero. A ternary subsemigroup  $Q$  of a ternary semigroup  $S$  is called a quasi-ideal of  $S$  if  $QSS \cap (SQS \cup SSQSS) \cap SSQ \subseteq Q$  and a ternary subsemigroup  $B$  of a ternary semigroup  $S$  is called a bi-ideal of  $S$  if  $BSBSB \subseteq B$ . It is easy to see that every quasi-ideal in a ternary semigroup is a bi-ideal of  $S$ . An element  $a$  in a ternary semigroup  $S$  is called regular if there exists an element  $x$  in  $S$  such that  $axa = a$ . A ternary semigroup is called regular if all of its elements are regular. A ternary semigroup  $S$  is regular if and only if  $R \cap M \cap L = RML$  for every right ideal  $R$ , lateral ideal  $M$  and left ideal  $L$  of  $S$ .

## 2 Generalised Quasi-ideals in Ternary Semigroup

In this section, we introduce the concept of generalised quasi-ideals in ternary semigroups and prove some results related to the same.

**Definition 2.1.** A ternary subsemigroup  $Q$  of a ternary semigroup  $S$  is called a generalised quasi-ideal or  $(m, (p, q), n)$ -quasi-ideal of  $S$  if  $Q(SS)^m \cap (S^pQS^q \cup S^pSSQSS^q) \cap (SS)^nQ \subseteq Q$ , where  $m, n, p, q$  are positive integers greater than 0 and  $p + q = \text{even}$ .

**Remark 2.1.** Every quasi-ideal of a ternary semigroup  $S$  is  $(1, (1, 1), 1)$ -quasi-ideal of  $S$ . But  $(m, (p, q), n)$ -quasi-ideal of a ternary semigroup  $S$  need not be a quasi-ideal of  $S$ .

**Example 1.** Let  $Z^- \setminus \{-1\}$  be the set of all negative integers excluding  $\{0\}$ . Then  $Z^- \setminus \{-1\}$  is a ternary semigroup with usual ternary multiplication. Consider  $Q = \{-3\} \cup \{k \in Z^- : k \leq -14\}$ . Clearly  $Q$  is a non-empty ternary subsemigroup of  $S$  and also  $Q$  is  $(2, (1, 1), 3)$ -quasi-ideal of  $S$ . Now,  $\{-12\} \in QSS \cap (SQS \cup SSQSS) \cap SSQ$ . But  $\{-12\} \notin Q$ . Therefore  $QSS \cap (SQS \cup SSQSS) \cap SSQ \not\subseteq Q$ . Hence  $Q$  is not quasi-ideal of  $Z^- \setminus \{-1\}$ .

**Lemma 2.1.** Non-empty intersection of arbitrary collection of ternary subsemigroups of a ternary semigroup  $S$  is a ternary subsemigroup of  $S$ .

*Proof.* Let  $T_i$  be a ternary subsemigroup of  $S$  for all  $i \in I$  such that  $\bigcap_{i \in I} T_i \neq \emptyset$ . Let  $t_1, t_2, t_3 \in \bigcap_{i \in I} T_i$ . Then  $t_1, t_2, t_3 \in T_i$  for all  $i \in I$ . Since  $T_i$  is a ternary subsemigroup of  $S$  for all  $i \in I$ , therefore  $t_1t_2t_3 \in T_i$  for all  $i \in I$ . Therefore  $t_1t_2t_3 \in \bigcap_{i \in I} T_i$ . Hence  $\bigcap_{i \in I} T_i$  is a ternary subsemigroup of  $S$ .  $\square$

**Theorem 2.1.** Let  $S$  be a ternary semigroup and  $Q_i$  be an  $(m, (p, q), n)$ -quasi-ideal of  $S$  such that  $\bigcap_{i \in I} Q_i \neq \emptyset$ . Then  $\bigcap_{i \in I} Q_i$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ .

*Proof.* Clearly  $\bigcap_{i \in I} Q_i$  is a ternary subsemigroup of  $S$  (by Lemma 2.1).

Let  $x \in \left[ \bigcap_{i \in I} Q_i (SS)^m \right] \cap \left[ S^p \bigcap_{i \in I} Q_i S^q \cup S^p S \bigcap_{i \in I} Q_i SS^q \right] \cap \left[ (SS)^n \bigcap_{i \in I} Q_i \right]$ . Then  $x \in$

$\bigcap_{i \in I} Q_i(SS)^m$ ,  $x \in S^p \bigcap_{i \in I} Q_i S^q \cup S^p S \bigcap_{i \in I} Q_i S S^q$  and  $x \in (SS)^n \bigcap_{i \in I} Q_i$ . This implies  $x \in Q_i(SS)^m$ ,  $x \in [S^p Q_i S^q \cup S^p S Q_i S S^q]$  and  $x \in (SS)^n Q_i$  for all  $i \in I$ . Therefore  $x \in [Q_i(SS)^m] \cap [S^p Q_i S^q \cup S^p S Q_i S S^q] \cap [(SS)^n Q_i] \subseteq Q_i$  for all  $i \in I$ , since  $Q_i$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ . Thus  $x \in Q_i$  for all  $i \in I$ . Therefore  $x \in \bigcap_{i \in I} Q_i$ . Hence  $\bigcap_{i \in I} Q_i$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ .  $\square$

**Remark 2.2.** Let  $Z^-$  be the set of all negative integers under ternary multiplication and  $Q_i = \{k \in Z^- : k \leq -i\}$  for all  $i \in I$ . Then  $Q_i$  is an  $(2, (1, 1), 3)$ -quasi-ideal of  $Z^-$  for all  $i \in I$ . But  $\bigcap_{i \in I} Q_i = \emptyset$ . So condition  $\bigcap_{i \in I} Q_i \neq \emptyset$  is necessary.

**Definition 2.2.** Let  $S$  be a ternary semigroup. Then a ternary subsemigroup

- (i)  $R$  of  $S$  is called an  $m$ -right ideal of  $S$  if  $R(SS)^m \subseteq R$ .
- (ii)  $M$  of  $S$  is called an  $(p, q)$ -lateral ideal of  $S$  if  $S^p M S^q \cup S^p S M S S^q \subseteq M$ ,
- (iii)  $L$  of  $S$  is called an  $n$ -left ideal of  $S$  if  $(SS)^n L \subseteq L$ ,

where  $m, n, p, q$  are positive integers and  $p + q$  is an even positive integer.

**Theorem 2.2.** Every  $m$ -right,  $(p, q)$ -lateral and  $n$ -left ideal of a ternary semigroup  $S$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ . But converse need not be true.

*Proof.* One way is straight forward. Conversely, let  $S = M_2(Z_0^-)$  be the ternary semigroup of  $2 \times 2$  square matrices over  $Z_0^-$ . Consider  $Q = \left\{ \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} : a \in Z_0^- \right\}$ . Then  $Q$  is an  $(2, (1, 1), 3)$ -quasi-ideal of  $S$ . But it is not 2-right ideal,  $(1, 1)$ -lateral ideal and 3-left ideal of  $S$ .  $\square$

**Theorem 2.3.** Let  $S$  be a ternary semigroup. Then the following statements hold:

- (i) Let  $R_i$  be an  $m$ -right ideal of  $S$  such that  $\bigcap_{i \in I} R_i \neq \emptyset$ . Then  $\bigcap_{i \in I} R_i$  is an  $m$ -right ideal of  $S$ .
- (ii) Let  $M_i$  be an  $(p, q)$ -lateral ideal of  $S$  such that  $\bigcap_{i \in I} M_i \neq \emptyset$ . Then  $\bigcap_{i \in I} M_i$  is an  $(p, q)$ -lateral ideal of  $S$ .
- (iii) Let  $L_i$  be an  $n$ -left ideal of  $S$  such that  $\bigcap_{i \in I} L_i \neq \emptyset$ . Then  $\bigcap_{i \in I} L_i$  is an  $n$ -left ideal of  $S$ .

*Proof.* Similar to the proof of Theorem 2.1  $\square$

**Theorem 2.4.** Let  $R$  be an  $m$ -right ideal,  $M$  be an  $(p, q)$ -lateral ideal and  $L$  be an  $n$ -left ideal of a ternary semigroup  $S$ . Then  $R \cap M \cap L$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ .

*Proof.* Suppose  $Q = R \cap M \cap L$ . Since every  $m$ -right,  $(p, q)$ -lateral and  $n$ -left ideal of ternary semigroup  $S$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ , therefore  $R, M$  and  $L$  are  $(m, (p, q), n)$ -quasi-ideals of  $S$ . Clearly,  $R \cap M \cap L$  is non-empty. By Theorem 2.1, we have  $Q = R \cap M \cap L$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ .  $\square$

**Lemma 2.2.** *Let  $Q$  be an  $(m, (p, q), n)$ -quasi-ideal of a ternary semigroup  $S$ . Then*

- (i)  $R = Q \cup Q(SS)^m$  is an  $m$ -right ideal of  $S$ .
- (ii)  $M = Q \cup (S^pQS^q \cup S^pSQSS^q)$  is an  $(p, q)$ -lateral ideal of  $S$ .
- (iii)  $L = Q \cup (SS)^nQ$  is an  $n$ -left ideal of  $S$ .

*Proof.* It is easy to show that  $R$  is ternary subsemigroup of  $S$ . Now to show that  $R$  is an  $m$ -right ideal of  $S$ .

$$\begin{aligned} R(SS)^m &= [(Q \cup Q(SS)^m)(SS)^m] \\ &= Q(SS)^m \cup Q(SS)^m(SS)^m \\ &= Q(SS)^m \cup Q(SSSS)^m \\ &\subseteq Q(SS)^m \cup Q(SS)^m \\ &= Q(SS)^m \subseteq R. \end{aligned}$$

Therefore  $R$  is an  $m$ -right ideal of  $S$ . Similarly, we can show that  $M$  is an  $(p, q)$ -lateral ideal of  $S$  and  $L$  is an  $n$ -left ideal of  $S$ .  $\square$

**Theorem 2.5.** *Every  $(m, (p, q), n)$ -quasi-ideal in a regular ternary semigroup  $S$  is the intersection of  $m$ -right,  $(p, q)$ -lateral and  $n$ -left ideal of  $S$ .*

*Proof.* Let  $S$  be regular ternary semigroup and  $Q$  be an  $(m, (p, q), n)$ -quasi-ideal of  $S$ . Then  $R = Q \cup Q(SS)^m$ ,  $M = Q \cup (S^pQS^q \cup S^pSQSS^q)$  and  $L = Q \cup (SS)^nQ$  are  $m$ -right,  $(p, q)$ -lateral and  $n$ -left ideal of  $S$  respectively. Clearly  $Q \subseteq R$ ,  $Q \subseteq M$  and  $Q \subseteq L$  implies  $Q \subseteq R \cap M \cap L$ . Since  $S$  is regular therefore  $Q \subseteq Q(SS)^m$ ,  $Q \subseteq S^pQS^q \cup S^pSQSS^q$  and  $Q \subseteq (SS)^nQ$ .

Thus  $R = Q(SS)^m$ ,  $M = S^pQS^q \cup S^pSQSS^q$  and  $L = (SS)^nQ$ . Now

$$R \cap M \cap L = Q(SS)^m \cap (S^pQS^q \cup S^pSQSS^q) \cap (SS)^nQ \subseteq Q$$

Hence,  $Q = R \cap M \cap L$ .  $\square$

### 3 Generalised Minimal Quasi-ideals

In this section, we study the concept of generalised minimal quasi-ideal or minimal  $(m, (p, q), n)$ -quasi-ideals of ternary semigroup  $S$ .

An  $(m, (p, q), n)$ -quasi-ideal  $Q$  of a ternary semigroup  $S$  is called minimal  $(m, (p, q), n)$ -quasi-ideal of  $S$  if  $Q$  does not properly contain any  $(m, (p, q), n)$ -quasi-ideal of  $S$ . Similarly, we can define minimal  $m$ -right ideals, minimal  $(p, q)$ -lateral ideals and minimal  $n$ -left ideals of a ternary semigroup.

**Lemma 3.1.** *Let  $S$  be a ternary semigroup and  $a \in S$ . Then the following statements hold:*

- (i)  $a(SS)^m$  is an  $m$ -right ideal of  $S$ .
- (ii)  $(S^paS^q \cup S^pSaSS^q)$  is an  $(p, q)$ -lateral ideal of  $S$ .

- (iii)  $(SS)^n a$  is an  $n$ -left ideal of  $S$ .  
 (iv)  $a(SS)^m \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ .

*Proof.* (i), (ii) and (iii) are obvious. (iv) follows from (i), (ii), (iii) and Theorem 2.4.  $\square$

**Theorem 3.1.** *Let  $S$  be a ternary semigroup and  $Q$  be an  $(m, (p, q), n)$ -quasi-ideal of  $S$ . Then  $Q$  is minimal iff  $Q$  is the intersection of some minimal  $m$ -right ideal  $R$ , minimal  $(p, q)$ -lateral ideal  $M$  and minimal  $n$ -left ideal  $L$  of  $S$ .*

*Proof.* Suppose  $Q$  is minimal  $(m, (p, q), n)$ -quasi-ideal of  $S$ . Let  $a \in Q$ . Then by above Lemma, we have  $a(SS)^m$  is an  $m$ -right ideal,  $(S^p a S^q \cup S^p S a S S^q)$  is an  $(p, q)$ -lateral ideal,  $(SS)^n a$  is an  $n$ -left ideal and  $a(SS)^m \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ . Now,

$$\begin{aligned} & a(SS)^m \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a \\ & \subseteq Q(SS)^m \cap (S^p Q S^q \cup S^p S Q S S^q) \cap (SS)^n Q \\ & \subseteq Q. \end{aligned}$$

Since  $Q$  is minimal therefore  $a(SS)^m \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a = Q$ . Now, to show that  $a(SS)^m$  is minimal  $m$ -right ideal of  $S$ . Let  $R$  be an  $m$ -right ideal of  $S$  contained in  $a(SS)^m$ . Then

$$\begin{aligned} & R \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a \\ & \subseteq a(SS)^m \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a \\ & = Q. \end{aligned}$$

Since  $R \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$  and  $Q$  is minimal, therefore  $R \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a = Q$ . This implies  $Q \subseteq R$  and therefore

$$a(SS)^m \subseteq Q(SS)^m \subseteq R(SS)^m \subseteq R$$

implies  $R = a(SS)^m$ . Thus  $m$ -right ideal  $a(SS)^m$  is minimal. Similarly, we can prove that  $(S^p a S^q \cup S^p S a S S^q)$  is minimal  $(p, q)$ -lateral ideal of  $S$  and  $(SS)^n a$  is minimal  $n$ -left ideal of  $S$ .

Conversely, assume that  $Q = R \cap M \cap L$  for some minimal  $m$ -right ideal  $R$ , minimal  $(p, q)$ -lateral ideal  $M$  and minimal  $n$ -left ideal  $L$ . So,  $Q \subseteq R$ ,  $Q \subseteq M$  and  $Q \subseteq L$ . Let  $Q'$  be an  $(m, (p, q), n)$ -quasi-ideal of  $S$  contained in  $Q$ . Then  $Q'(SS)^m \subseteq Q(SS)^m \subseteq R(SS)^m \subseteq R$ . Similarly,  $(S^p Q' S^q \cup S^p S Q' S S^q) \subseteq M$  and  $(SS)^n Q' \subseteq (SS)^n Q \subseteq L$ .

Now  $Q'(SS)^m$  is an  $m$ -right ideal of  $S$ , as  $Q'(SS)^m(SS)^m \subseteq Q'(SS)^m$ . Similarly,  $(S^p Q' S^q \cup S^p S Q' S S^q)$  is an  $(p, q)$ -lateral ideal of  $S$  and  $(SS)^n Q'$  is an  $n$ -left ideal of  $S$ . Since  $R$ ,  $M$  and  $L$  are minimal  $m$ -right ideal, minimal  $(p, q)$ -lateral ideal and minimal  $n$ -left ideal of  $S$  respectively, therefore  $Q'(SS)^m = R$ ,  $S^p Q' S^q \cup S^p S Q' S S^q = M$  and  $(SS)^n Q' = L$ .

Thus  $Q = R \cap M \cap L = Q'(SS)^m \cap (S^p Q' S^q \cup S^p S Q' S S^q) \cap (SS)^n Q' \subseteq Q'$ . Hence  $Q = Q'$ . Thus  $Q$  is minimal  $(m, (p, q), n)$ -quasi-ideal of  $S$ .  $\square$

**Note.** A ternary semigroup  $S$  need not contains a minimal  $(m, (p, q), n)$ -quasi-ideal of  $S$ .

For example, let  $Z^-$  be the set of all negative integers. Then  $Z^-$  is a ternary semigroup with usual ternary multiplication. Let  $Q = \{-2, -3, -4, \dots\}$ . Then  $Q$  is an  $(2, (1, 1), 3)$ -quasi-ideal of  $Z^-$ . Suppose  $Q$  is minimal  $(2, (1, 1), 3)$ -quasi-ideal of  $Z^-$ . Let  $Q' = Q \setminus \{-2\}$ . Then we can easily show that  $Q'$  is an  $(2, (1, 1), 3)$ -quasi-ideal of  $Z^-$ . But  $Q'$  is proper subset of  $Q$ . This is contradiction. Hence,  $Z^-$  does not contain a minimal  $(m, (p, q), n)$ -quasi-ideal.

**Theorem 3.2.** Let  $S$  be a ternary semigroup. Then the following holds:

- (i) An  $m$ -right ideal  $R$  is minimal iff  $a(SS)^m = R$  for all  $a \in R$ .
- (ii) An  $(p, q)$ -lateral ideal  $M$  is minimal iff  $(S^p a S^q \cup S^p S a S S^q) = M$  for all  $a \in M$ .
- (iii) An  $n$ -left ideal  $L$  is minimal iff  $(SS)^n a = L$  for all  $a \in L$ .
- (iv) An  $(m, (p, q), n)$ -quasi-ideal  $Q$  is minimal iff  $a(SS)^m \cap (S^p a S^q \cup S^p S a S S^q) \cap (SS)^n a = Q$  for all  $a \in Q$ .

*Proof.* (i) Suppose  $m$ -right ideal  $R$  is minimal. Let  $a \in R$ . Then  $a(SS)^m \subseteq R(SS)^m \subseteq R$ . By Lemma 3.1, we have  $a(SS)^m$  is an  $m$ -right ideal of  $S$ . Since  $R$  is minimal  $m$ -right ideal of  $S$  therefore  $a(SS)^m = R$ .

Conversely, Suppose that  $a(SS)^m = R$  for all  $a \in R$ . Let  $R'$  be an  $m$ -right ideal of  $S$  contained in  $R$ . Let  $x \in R'$ . Then  $x \in R$ . By assumption, we have  $x(SS)^m = R$  for all  $x \in R$ .  $R = x(SS)^m \subseteq R'(SS)^m \subseteq R'$ . This implies  $R \subseteq R'$ . Thus,  $R = R'$ . Hence,  $R$  is minimal  $m$ -right ideal.

Similarly we can prove (ii), (iii) and (iv). □

## 4 Generalised Bi-ideals in Ternary Semigroup

In this section, we define generalised bi-ideals in a ternary semigroup and give their characterizations.

**Definition 4.1.** A ternary subsemigroup  $B$  of a ternary semigroup  $S$  is called a generalised bi-ideal or  $(m, (p, q), n)$  bi-ideal of  $S$  if  $B(SS)^{m-1} S^p B S^q (SS)^{n-1} B \subseteq B$ , where  $m, n, p, q$  are positive integers greater than zero and  $p$  and  $q$  are odd.

**Remark.** Every bi-ideal of a ternary semigroup  $S$  is  $(1, (1, 1), 1)$ -bi-ideal of  $S$ . But every  $(m, (p, q), n)$ -bi-ideal of a ternary semigroup  $S$  need not be a bi-ideal of  $S$  which is illustrated by the following example.

**Example 2.** Let  $Z^- \setminus \{-1\}$  be the set of all negative integers excluding  $\{0\}$ . Then  $Z^- \setminus \{-1\}$  is a ternary semigroup with usual ternary multiplication. Consider  $B = \{-3, -27\} \cup \{k \in Z^- : k \leq -110\}$ . Clearly  $B$  is a non-empty ternary subsemigroup of  $S$  and also  $B$  is  $(3, (1, 1), 4)$ -bi-ideal of  $S$ . Now  $-108 \in BSBSB$ . But  $-108 \notin B$ . Therefore  $BSBSB \not\subseteq B$ . Hence  $B$  is not a bi-ideal of  $Z^- \setminus \{-1\}$ .

**Theorem 4.1.** Let  $S$  be a ternary semigroup and  $B_i$  be an  $(m, (p, q), n)$ -bi-ideals of  $S$  such that  $\bigcap_{i \in I} B_i \neq \emptyset$ . Then  $\bigcap_{i \in I} B_i$  is an  $(m, (p, q), n)$  bi-ideal of  $S$ .

*Proof.* It is straight forward.  $\square$

**Remark.** Let  $Z^-$  be the set of all negative integers. Then  $Z^-$  is a ternary semigroup under usual ternary multiplication and  $B_i = \{k \in Z^- : k \leq -i\}$  for all  $i \in I$ . Then  $B_i$  is an  $(3, (1, 1), 4)$ -bi-ideal of  $Z^-$  for all  $i \in I$ . But  $\bigcap_{i \in I} B_i = \emptyset$ . So condition  $\bigcap_{i \in I} B_i \neq \emptyset$  is necessary.

**Theorem 4.2.** Every  $(m, (p, q), n)$ -quasi-ideal of a ternary semigroup  $S$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ .

*Proof.* Let  $Q$  be an  $(m, (p, q), n)$ -quasi-ideal of  $S$ . Then

$$Q(SS)^{m-1}S^pQS^q(SS)^{n-1}Q \subseteq Q(SS)^{m-1}S^pSS^q(SS)^{n-1}S \subseteq Q(SS)^m.$$

Similarly,

$$Q(SS)^{m-1}S^pQS^q(SS)^{n-1}Q \subseteq S(SS)^{m-1}(S^pQS^q)(SS)^{n-1}S \subseteq S^{p+1}QS^{q+1}.$$

Again  $\{0\} \subseteq S^pQS^q$ . So

$$Q(SS)^{m-1}S^pQS^q(SS)^{n-1}Q \subseteq S^pQS^q \cup S^{p+1}QS^{q+1}.$$

Also,

$$Q(SS)^{m-1}S^pQS^q(SS)^{n-1}Q \subseteq S(SS)^{m-1}S^pSS^q(SS)^{n-1}Q \subseteq (SS)^nQ.$$

Consequently,

$$Q(SS)^{m-1}S^pQS^q(SS)^{n-1}Q \subseteq Q(SS)^m \cap (S^pQS^q \cup S^{p+1}QS^{q+1}) \cap (SS)^nQ \subseteq Q.$$

Hence  $Q$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ .  $\square$

**Remark.** Every  $(m, (p, q), n)$ -bi-ideal need not be an  $(m, (p, q), n)$ -quasi-ideal of  $S$  which is illustrated by the following example.

**Example 3.** Consider the ternary semigroup  $S = Z^- \setminus \{-1\}$  with usual ternary multiplication and let  $B = \{-3, -27\} \cup \{k \in Z^- : k \leq -194\}$ . Clearly,  $B$  is non-empty ternary subsemigroup of  $S$  and also  $B$  is  $(2, (1, 1), 3)$ -bi-ideal of  $S$ . Now,  $-192 \in B(SS)^2 \cap (SBS \cup SSBSS) \cap (SS)^3B$ . But  $-192 \notin B$ . Therefore  $B(SS)^2 \cap (SBS \cup SSBSS) \cap (SS)^3B \not\subseteq B$ . Hence  $B$  is not  $(2, (1, 1), 3)$ -quasi-ideal of  $S$ .

**Theorem 4.3.** A ternary subsemigroup  $B$  of a regular ternary semigroup  $S$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$  if and only if  $B = BSB$ .

*Proof.* Suppose  $B$  is an  $(m, (p, q), n)$ -bi-ideal of a regular ternary semigroup  $S$ . Let  $b \in B$ . Then there exists  $x \in S$  such that  $b = bxb$ . This implies that  $b \in BSB$ . Hence  $B \subseteq BSB$ . Now,

$$BSB \subseteq BSBSBSBSB \subseteq B(SS)(SBS)(SS)B \subseteq B.$$



Therefore  $B = BSB$ .

Conversely, if  $B = BSB$ , then

$$B(SS)^{m-1}S^pBS^q(SS)^{n-1}B \subseteq B(SS)^{m-1}S^pSS^q(SS)^{n-1}B \subseteq BSB = B.$$

Hence  $B$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ .  $\square$

**Theorem 4.4.** *Let  $S$  be a regular ternary semigroup. Then every  $(m, (p, q), n)$ -bi-ideal of  $S$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ .*

*Proof.* Let  $B$  be an  $(m, (p, q), n)$ -bi-ideal of  $S$ . Let  $a \in B(SS)^m \cap (S^pBS^q \cup S^pSBSS^q) \cap (SS)^nB$ . Then  $a \in B(SS)^m$ ,  $a \in (S^pBS^q \cup S^pSBSS^q)$  and  $a \in (SS)^nB$ . Thus  $a = b(SS)^m = S^pb'S^q \cup S^pSb''SS^q = (SS)^nb'''$  for some  $b, b', b'', b''' \in B$ . Since  $S$  is regular, therefore for  $a \in S$  there exists an element  $x$  in  $S$  such that  $a = axa$ . Then

$$\begin{aligned} a &= axa = axaxa \\ &= b(SS)^m x (S^pb'S^q \cup S^pSb''SS^q) x (SS)^nb''' \\ &\in B(SS)^m S (S^pBS^q \cup S^pSBSS^q) S (SS)^n B \\ &= [B(SS)^m S S^pBS^q S (SS)^n B] \cup [B(SS)^m S S^pSBSS^q S (SS)^n B] \\ &\subseteq B[(SS)^m S S^pSS^q S (SS)^n] B \cup B[(SS)^m S S^pSSSS^q S (SS)^n] B \\ &\subseteq BSB \cup BSB = B \cup B = B. \end{aligned}$$

Thus  $a \in B$ . Therefore  $B(SS)^m \cap (S^pBS^q \cup S^pSBSS^q) \cap (SS)^nB \subseteq B$ . Hence  $B$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ .  $\square$

It is easy to prove the following propositions:

**Proposition 4.5.** *The intersection of an  $(m, (p, q), n)$ -bi-ideal  $B$  of a ternary semigroup  $S$  with a ternary subsemigroup  $T$  of  $S$  is either empty or an  $(m, (p, q), n)$ -bi-ideal of  $T$ .*

**Proposition 4.6.** *Let  $B$  be an  $(m, (p, q), n)$ -bi-ideal of a ternary semigroup  $S$  and  $T_1, T_2$  are two ternary subsemigroups of  $S$ . Then  $BT_1T_2, T_1BT_2$  and  $T_1T_2B$  are  $(m, (p, q), n)$ -bi-ideals of  $S$ .*

**Proposition 4.7.** *Let  $B_1, B_2$  and  $B_3$  are three  $(m, (p, q), n)$ -bi-ideals of a ternary semigroup  $S$ . Then  $B_1B_2B_3$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ .*

**Proposition 4.8.** *Let  $Q_1, Q_2$  and  $Q_3$  are three  $(m, (p, q), n)$ -quasi-ideals of a ternary semigroup  $S$ . Then  $Q_1Q_2Q_3$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ .*

**Proposition 4.9.** *Let  $R$  be an  $m$ -right,  $M$  be an  $(p, q)$ -lateral and  $L$  be an  $n$ -left ideal of a ternary semigroup  $S$ . Then the ternary subsemigroup  $B = RML$  of  $S$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ .*

**Theorem 4.10.** *Let  $S$  be a regular ternary semigroup. If  $B$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ , then  $B(SS)^{m-1}S^pBS^q(SS)^{n-1}B = B$ .*

*Proof.* Let  $B$  be an  $(m, (p, q), n)$ -bi-ideal of  $S$ . Let  $a \in B$ . Then  $a \in S$ . Since  $S$  is regular, therefore there exists  $x \in S$  such that  $a = axa$ . Now  $a = axa = a(xa)(xax)(ax)a \in B(SS)(SBS)(SS)B$ . Similarly, by property of regularity it is easy to show that  $a \in B(SS)^{m-1}S^pBS^q(SS)^{n-1}B$ . Thus,  $B \subseteq B(SS)^{m-1}S^pBS^q(SS)^{n-1}B$ . Since  $B$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ , therefore  $B(SS)^{m-1}S^pBS^q(SS)^{n-1}B \subseteq B$ . Hence  $B(SS)^{m-1}S^pBS^q(SS)^{n-1}B = B$   $\square$

**Corollary 4.1.** *Let  $S$  be a regular ternary semigroup. If  $Q$  is an  $(m, (p, q), n)$ -quasi-ideal of  $S$ , then  $Q(SS)^{m-1}S^pQS^q(SS)^{n-1}Q = Q$ .*

*Proof.* Since every  $(m, (p, q), n)$ -quasi-ideal of  $S$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ , therefore result follows directly.  $\square$

## 5 Generalised Minimal Bi-ideals

In this section, we introduce the concept of generalised minimal bi-ideal or minimal  $(m, (p, q), n)$ -bi-ideals in ternary semigroups.

**Definition 5.1.** *An  $(m, (p, q), n)$ -bi-ideal  $B$  of a ternary semigroup  $S$  is called minimal  $(m, (p, q), n)$ -bi-ideal of  $S$  if  $B$  does not properly contain any  $(m, (p, q), n)$ -bi-ideal of  $S$ .*

**Lemma 5.1.** *Let  $S$  be a ternary semigroup and  $a \in S$ . Then the following holds:*

- (i)  $a(SS)^{m-1}$  is an  $m$ -right ideal of  $S$ .
- (ii)  $S^paS^q$  is an  $(p, q)$ -lateral ideal of  $S$ .
- (iii)  $(SS)^{n-1}a$  is an  $n$ -left ideal of  $S$ .
- (iv)  $a(SS)^{m-1}S^paS^q(SS)^{n-1}a$  is an  $(m, (p, q), n)$ -bi-ideal.

*Proof.* (i), (ii) and (iii) are obvious and (iv) follows from (i), (ii), (iii).  $\square$

**Theorem 5.1.** *Let  $S$  be a ternary semigroup and  $B$  be an  $(m, (p, q), n)$ -bi-ideal of  $S$ . Then  $B$  is minimal if and only if  $B$  is the product of some minimal  $m$ -right ideal  $R$ , minimal  $(p, q)$ -lateral ideal  $M$  and minimal  $n$ -left ideal  $L$  of  $S$ .*

*Proof.* Suppose  $B$  is minimal  $(m, (p, q), n)$ -bi-ideal of  $S$ . Let  $a \in B$ . Then by above Lemma,  $a(SS)^{m-1}$  is an  $m$ -right ideal,  $S^paS^q$  is an  $(p, q)$ -lateral ideal,  $(SS)^{n-1}a$  is an  $n$ -left ideal and  $a(SS)^{m-1}S^paS^q(SS)^{n-1}a$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ . Now  $a(SS)^{m-1}S^paS^q(SS)^{n-1}a \subseteq B(SS)^{m-1}S^pBS^q(SS)^{n-1}B \subseteq B$ . Since  $B$  is minimal, therefore  $a(SS)^{m-1}S^paS^q(SS)^{n-1}a = B$ . Now to show that  $a(SS)^{m-1}$  is minimal  $m$ -right ideal of  $S$ . Let  $R$  be an  $m$ -right ideal of  $S$  contained in  $a(SS)^{m-1}$ . Then  $R(S^paS^q)(SS)^{n-1}a \subseteq a(SS)^{m-1}(S^paS^q)(SS)^{n-1}a = B$ . Since  $R(S^paS^q)(SS)^{n-1}a$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$  and  $B$  is minimal, therefore  $R(S^paS^q)(SS)^{n-1}a = B$ . This implies  $B \subseteq R$ . Therefore  $a(SS)^{m-1} \subseteq B(SS)^{m-1} \subseteq R(SS)^{m-1} \subseteq R$ . Thus  $a(SS)^{m-1}$  is minimal. Similarly we can prove that  $S^paS^q$  is minimal  $(p, q)$ -lateral ideal of  $S$  and  $(SS)^{n-1}a$  is minimal  $n$ -left ideal of  $S$ .

Conversely, assume that  $B = RML$  for some minimal  $m$ -right ideal  $R$ , minimal  $(p, q)$ -lateral ideal  $M$  and minimal  $n$ -left ideal  $L$ . So  $B \subseteq R$ ,  $B \subseteq M$  and  $B \subseteq L$ . Let  $B'$  be an  $(m, (p, q), n)$ -bi-ideal of  $S$  contained in  $B$ . Then  $B'(SS)^{m-1} \subseteq B(SS)^{m-1} \subseteq R(SS)^{m-1} \subseteq R$ . Similarly,  $S^p B' S^q \subseteq S^p B S^q \subseteq S^p M S^q \subseteq M$  and  $(SS)^{n-1} B' \subseteq (SS)^{n-1} B \subseteq (SS)^{n-1} L \subseteq L$ . Now,  $B'(SS)^{m-1}(SS)^m \subseteq B'(SS)^{m-1}$ . So  $B'(SS)^{m-1}$  is an  $m$ -right ideal of  $S$ . Similarly  $S^p B' S^q$  is an  $(p, q)$ -lateral ideal and  $(SS)^{n-1} B'$  is an  $n$ -left ideal of  $S$ . Since  $R$ ,  $M$  and  $L$  are minimal  $m$ -right ideal, minimal  $(p, q)$ -lateral ideal and minimal  $n$ -left ideal of  $S$  respectively, therefore  $B'(SS)^{m-1} = R$ ,  $S^p B' S^q = M$  and  $(SS)^{n-1} B' = L$ . Thus  $B = RML = B'(SS)^{m-1} S^p B' S^q (SS)^{n-1} B' \subseteq B'$ . Hence  $B = B'$ . Consequently,  $B$  is minimal  $(m, (p, q), n)$ -bi-ideal of  $S$ .  $\square$

**Definition 5.2.** Let  $S$  be a ternary semigroup. Then  $S$  is called a bi-simple ternary semigroup if  $S$  is the unique  $(m, (p, q), n)$ -bi-ideal of  $S$ .

**Theorem 5.2.** Let  $S$  be a ternary semigroup and  $B$  be an  $(m, (p, q), n)$ -bi-ideal of  $S$ . Then  $B$  is a minimal  $(m, (p, q), n)$ -bi-ideal of  $S$  if and if  $B$  is a bi-simple ternary semigroup.

*Proof.* Suppose  $B$  is a minimal  $(m, (p, q), n)$ -bi-ideal of  $S$ . Let  $C$  be an  $(m, (p, q), n)$ -bi-ideal of  $B$ . Then  $C(BB)^{m-1} B^p C B^q (BB)^{n-1} C \subseteq C \subseteq B$ . By Proposition 4.9,  $BCC$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ . Therefore

$(BCC)(SS)^{m-1} S^p (BCC) S^q (SS)^{n-1} BCC \subseteq BCC \subseteq BBB \subseteq B$ . Since  $B$  is minimal, therefore  $BCC = B$ . It is easy to show that  $C(BB)^{m-1} B^p C B^q (BB)^{n-1} C$  is an  $(m, (p, q), n)$ -bi-ideal of  $S$ .

Since  $B$  is minimal, therefore  $C(BB)^{m-1} B^p C B^q (BB)^{n-1} C = B$ . This implies  $B = C(BB)^{m-1} B^p C B^q (BB)^{n-1} C \subseteq C$ . Hence  $C = B$ . Consequently,  $B$  is a bi-simple ternary semigroup.

Conversely, suppose  $B$  is a bi-simple ternary semigroup. Let  $C$  be an  $(m, (p, q), n)$ -bi-ideal of  $S$  such that  $C \subseteq B$ . Then

$$C(BB)^{m-1} B^p C B^q (BB)^{n-1} C \subseteq C(SS)^{m-1} S^p C S^q (SS)^{n-1} C \subseteq C$$

which implies that  $C$  is an  $(m, (p, q), n)$ -bi-ideal of  $B$ . Since  $B$  is bi-simple ternary semigroup, therefore  $C = B$ . Hence  $B$  is minimal.  $\square$

## References

- [1] F. M. Sioson, Ideal theory in ternary semigroups, *Math. Japon.*,10 (1965), 63–84.
- [2] O. Steinfeld, *Quasi-ideals in rings and semigroups*, Akademiai Kiado, Budapest (1978).
- [3] R. Chinram, A note on  $(m, n)$  Quasi-ideals in semigroups, *International Journal of Pure and Applied Mathematics*, **49** (1), (2008), 45–50.
- [4] R. Chinram, Generalised quasi-ideals of semigroups, *KKU Sci. J.*, **37** (2), (2009), 213–210.

- [5] R. A. Good and D. R. Hughes: Associated groups for a semigroup, Bull. Amer. Math. Soc. 58 (1952), 624–625:
- [6] S. Kar, On quasi-ideal and bi-ideals in ternary semiring, *Int. J. Math. & Math. Sci.*, **18**, (2005), 3015–3023.
- [7] V.N. Dixit and S. Dewan, A note on quasi and bi-ideals in ternary semigroups, *Int. J. Math. & Math. Sci.* **18** (3), (1995), 501–508.

**DOI: 10.7862/rf.2014.3**

**Manish Kant Dubey**

email: kantmanish@yahoo.com

SAG, DRDO, Metcalf House,  
Delhi 110054, India.

**Rahatgi Anuradha - corresponding author**

email: anuvikaspuri@yahoo.co.in

Department of Mathematics,  
University of Delhi,  
Delhi 110007, India.

*Received 5.02.2013, Accepted 25.10.2013*

## Remarkable identities

Jan Górowski, Jerzy Żabowski

ABSTRACT: In the paper a number of identities involving even powers of the values of functions tangent, cotangent, secans and cosecans are proved. Namely, the following relations are shown:

$$\begin{aligned}\sum_{j=1}^{m-1} f^{2n} \left( \frac{\pi j}{2m} \right) &= w_f(m), \\ \sum_{j=0}^{m-1} f^{2n} \left( \frac{2j+1}{4m} \pi \right) &= v_f(m), \\ \sum_{j=1}^m f^{2n} \left( \frac{\pi j}{2m+1} \right) &= \tilde{w}_f(m),\end{aligned}$$

where  $m, n$  are positive integers,  $f$  is one of the functions: tangent, cotangent, secans or cosecans and  $w_f(x), v_f(x), \tilde{w}_f(x)$  are some polynomials from  $\mathbb{Q}[x]$ .

One of the remarkable identities is the following:

$$\sum_{j=0}^{m-1} \sin^{-2} \frac{(2j+1)\pi}{2m} = m^2, \quad \text{provided } m \geq 1.$$

Some of these identities are used to find, by elementary means, the sums of the series of the form  $\sum_{j=1}^{\infty} \frac{1}{j^{2n}}$ , where  $n$  is a fixed positive integer. One can also notice that Bernoulli numbers appear in the leading coefficients of the polynomials  $w_f(x), v_f(x)$  and  $\tilde{w}_f(x)$ .

AMS Subject Classification: 11M06, 11Y60, 10H10

Keywords and Phrases: *trigonometric identity, zeta function*

In [7] the following formulas have been proved

$$\sum_{j=1}^m \cot^2 \frac{\pi j}{2m+1} = \frac{m(2m-1)}{3}, \quad \sum_{j=1}^m \sin^{-2} \frac{\pi j}{2m+1} = \frac{2m(m+1)}{3}, \quad (1)$$

where  $m \in \mathbb{N}_1$ . By  $\mathbb{N}_k$  for a positive integer  $k$  we mean  $\mathbb{N} \setminus \{0, 1, 2, \dots, k-1\}$ . The above identities were then used in an elementary proof of the formula  $\sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}$ .

In this paper we develop the ideas from [7] to prove more generalized identities than (1). Next we use some of them to find the sum of  $\sum_{k=1}^{\infty} \frac{1}{k^{2n}}$ , where  $n \in \mathbb{N}_1$ . The general identities given in this article yield, in particular, the following identity of uncommon beauty

$$\sum_{j=0}^{m-1} \sin^{-2} \frac{2j+1}{2m} \pi = m^2, \quad m \in \mathbb{N}_1.$$

Some elementary methods of finding the sums of the series of the form  $\sum_{j=1}^{\infty} \frac{1}{j^{2n}}$  may be found for example in [1], [3], [5], [6], [8].

We start by recalling some basic facts on symmetric polynomials in  $m$  variables. Put

$$\sigma_n = \sum_{j=1}^m x_j^n \quad \text{for } n \in \mathbb{N}_1,$$

$$\tau_k = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq m} x_{j_1} x_{j_2} \dots x_{j_k} \quad \text{for } k \in \{1, 2, \dots, m\}.$$

Moreover, for the convenience set  $\tau_k = 0$  for  $k > m$ .

The following lemma comes from [2].

**Lemma 1** (Newton). *Let  $n \in \mathbb{N}_1$ , then*

$$\sigma_n - \tau_1 \sigma_{n-1} + \tau_2 \sigma_{n-2} - \dots + (-1)^{n-1} \tau_{n-1} \sigma_1 + (-1)^n n \tau_n = 0. \quad (2)$$

In view of Lemma 1 we have

$$\sigma_n = \det \begin{pmatrix} (-1)^{n+1} n \tau_n & -\tau_1 & \tau_2 & \dots & (-1)^{n-2} \tau_{n-2} & (-1)^{n-1} \tau_{n-1} \\ (-1)^n (n-1) \tau_{n-1} & 1 & -\tau_1 & \dots & (-1)^{n-3} \tau_{n-3} & (-1)^{n-2} \tau_{n-2} \\ (-1)^{n-1} (n-2) \tau_{n-2} & 0 & 1 & \dots & (-1)^{n-4} \tau_{n-4} & (-1)^{n-3} \tau_{n-3} \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ -2\tau_2 & 0 & 0 & \dots & 1 & -\tau_1 \\ \tau_1 & 0 & 0 & \dots & 0 & 1 \end{pmatrix} \quad (3)$$

for every  $n \in \mathbb{N}_1$ . Indeed, putting in (2) instead of  $n$  respectively  $n-1, n-2, \dots, 1$  we get, together with (2), the system of  $n$  equations in  $n$  variables:  $\sigma_1, \dots, \sigma_n$ . Such a system is a Cramer's system and by the Cramer's rule we get (3).

From now on by  $D_{\tan}$  and  $D_{\cot}$  we denote the domains of the trigonometric functions tangent and cotangent, respectively.

**Lemma 2.** *The following identities hold true:*

- (A)  $\frac{\sin 2mx}{\cos^{2m} x} \cot x = \sum_{j=0}^m \binom{2m}{2j+1} (-1)^j \tan^{2j} x, \quad (m, x) \in \mathbb{N} \times (D_{\tan} \cap D_{\cot});$
- (B)  $\frac{\cos 2mx}{\cos^{2m} x} = \sum_{j=0}^m \binom{2m}{2j} (-1)^j \tan^{2j} x, \quad (m, x) \in \mathbb{N} \times D_{\tan};$

$$\begin{aligned}
\text{(C)} \quad & \frac{\sin(2m+1)x}{\cos^{2m+1}x} \cot x = \sum_{j=0}^m \binom{2m+1}{2j+1} (-1)^j \tan^{2j} x, \quad (m, x) \in \mathbb{N} \times (D_{\tan} \cap D_{\cot}); \\
\text{(D)} \quad & \frac{\sin(2m+1)x}{\sin^{2m+1}x} = \sum_{j=0}^m \binom{2m+1}{2j+1} (-1)^j \cot^{2m-2j} x, \quad (m, x) \in \mathbb{N} \times D_{\cot}; \\
\text{(E)} \quad & \frac{\cos(2m+1)x}{\cos^{2m+1}x} = \sum_{j=0}^m \binom{2m+1}{2j} (-1)^j \tan^{2j} x, \quad (m, x) \in \mathbb{N} \times D_{\tan}; \\
\text{(F)} \quad & \frac{\cos(2m+1)x}{\sin^{2m+1}x} \tan x = \sum_{j=0}^m \binom{2m+1}{2j} (-1)^j \cot^{2m-2j} x, \quad (m, x) \in \mathbb{N} \times (D_{\tan} \cap D_{\cot}).
\end{aligned}$$

*Proof.* It is a known fact that

$$\sum_{j=0}^k \binom{k}{j} \cos^{k-j} x (i \sin x)^j = (\cos x + i \sin x)^k = \cos kx + i \sin kx$$

for  $k \in \mathbb{N}$  and  $x \in \mathbb{R}$ . Putting  $k = 2m$  in the above equation and comparing real and imaginary parts of the both sides we obtain (A) and (B). Similarly, with  $k = 2m + 1$  we get (C), (D), (E) and (F).  $\square$

Now we prove the following result.

**Theorem 1.** For every  $m \in \mathbb{N}_2$  and any  $n \in \mathbb{N}_1$ ,

$$\sigma_{n,m}(A) = \sum_{j=1}^{m-1} \tan^{2n} \frac{\pi j}{2m} = \sum_{j=1}^{m-1} \cot^{2n} \frac{\pi j}{2m},$$

where  $\sigma_{n,m}(A)$  denotes the determinant given by (3) in which  $\tau_j = \frac{\binom{2m}{2j+1}}{2m}$  for  $j \in \{1, 2, \dots, n\}$ .

*Proof.* Replace in the identity (A) of Lemma 2,  $\tan^2 x$  by  $t$  and set

$$w_A(t) = \sum_{j=0}^m \binom{2m}{2j+1} (-1)^j t^j, \quad (4)$$

then  $w_A(t)$  is a polynomial of order  $m - 1$  in the real variable  $t$ .

On the other hand, substituting  $\frac{\pi l}{2m}$ , where  $l \in \{1, 2, \dots, m - 1\}$ , for  $x$  in (A) we get

$$0 = \sum_{j=0}^m \binom{2m}{2j+1} (-1)^j \tan^{2j} \frac{\pi l}{2m}, \quad l \in \{1, 2, \dots, m - 1\}.$$

Hence and by (4) we obtain

$$w_A(t) = (-1)^{m-1} \binom{2m}{2m-1} \prod_{j=1}^{m-1} \left( t - \tan^2 \frac{\pi j}{2m} \right) = (-1)^{m-1} 2m \prod_{j=1}^{m-1} \left( t - \tan^2 \frac{\pi j}{2m} \right).$$

This and the Vieta's formulas give

$$\sum_{1 \leq k_1 < k_2 < \dots < k_j \leq m-1} \tan^2 \frac{\pi k_1}{2m} \tan^2 \frac{\pi k_2}{2m} \dots \tan^2 \frac{\pi k_j}{2m} = \frac{\binom{2m}{2j+1}}{2m}$$

and in view of (3) we have

$$\sigma_{n,m}(A) = \sum_{j=1}^{m-1} \tan^{2n} \frac{\pi j}{2m}.$$

As  $\tan \frac{\pi j}{2m} = \cot \frac{\pi(m-j)}{2m}$  for  $j \in \{1, 2, \dots, m-1\}$  we get

$$\sum_{j=1}^{m-1} \tan^{2n} \frac{\pi j}{2m} = \sum_{j=1}^{m-1} \cot^{2n} \frac{\pi(m-j)}{2m} = \sum_{j=1}^{m-1} \cot^{2n} \frac{\pi j}{2m},$$

which completes the proof.  $\square$

**Theorem 2.** For every  $m, n \in \mathbb{N}_1$  the following identity holds true:

$$\sigma_{n,m}(B) = \sum_{j=0}^{m-1} \tan^{2n} \frac{2j+1}{4m} \pi = \sum_{j=0}^{m-1} \cot^{2n} \frac{2j+1}{4m} \pi,$$

where  $\sigma_{n,m}(B)$  denotes the determinant given by (3) in which  $\tau_j = \binom{2m}{2j}$  for  $j \in \{1, 2, \dots, n\}$ .

*Proof.* Similarly as in the proof of Theorem 1, replace in the right hand side of the identity (B) of Lemma 2,  $\tan^2 x$  by  $t$  and set

$$w_B(t) = \sum_{j=0}^m \binom{2m}{2j} (-1)^j t^j.$$

Next, substitute  $\frac{2l+1}{4m}\pi$ , where  $l \in \{0, 1, \dots, m-1\}$ , for  $x$  in (B). This yields

$$0 = \sum_{j=0}^m \binom{2m}{2j} (-1)^j \tan^{2j} \frac{2l+1}{4m} \pi, \quad l \in \{0, 1, \dots, m-1\}.$$

Hence and by the definition of  $w_B(t)$  we get

$$w_B(t) = (-1)^m \prod_{j=0}^{m-1} \left( t - \tan^2 \frac{2j+1}{4m} \pi \right),$$

which in view of the Vieta's formulas gives

$$\sum_{1 \leq k_1 < k_2 < \dots < k_j \leq m-1} \tan^2 \frac{2k_1+1}{4m} \tan^2 \frac{2k_2+1}{4m} \dots \tan^2 \frac{2k_j+1}{4m} = \binom{2m}{2j}.$$



By this and (3),

$$\sigma_{n,m}(B) = \sum_{j=0}^{m-1} \tan^{2n} \frac{2j+1}{4m} \pi.$$

Using the same argument as in the proof of Theorem 1 we get

$$\sum_{j=0}^{m-1} \tan^{2n} \frac{2j+1}{2m} \pi = \sum_{j=0}^{m-1} \cot^{2n} \frac{2j+1}{4m} \pi$$

and the proof is completed.  $\square$

Using identities (C) and (D) of Lemma 2 and the same method as in proofs of Theorems 1 and 2 one may obtain

**Theorem 3.** For every  $m, n \in \mathbb{N}_1$  the following identity holds true:

$$\sigma_{n,m}(C) = \sum_{j=1}^m \tan^{2n} \frac{\pi j}{2m+1},$$

where  $\sigma_{n,m}(C)$  denotes the determinant given by (3) in which  $\tau_j = \binom{2m+1}{2j}$  for  $j \in \{1, 2, \dots, n\}$ .

**Theorem 4.** For every  $m, n \in \mathbb{N}_1$  the following identity holds true:

$$\sigma_{n,m}(D) = \sum_{j=1}^m \cot^{2n} \frac{\pi j}{2m+1},$$

where  $\sigma_{n,m}(D)$  denotes the determinant given by (3) in which  $\tau_j = \frac{1}{2m+1} \binom{2m+1}{2j+1}$  for  $j \in \{1, 2, \dots, n\}$ .

Finally, applying the same reasoning as in the proof of Theorem 1 from (E) and (F) of Lemma 2 we have

**Theorem 5.** For every  $m, n \in \mathbb{N}_1$  the following identity holds true:

$$\sigma_{n,m}(E) = \sum_{j=0}^{m-1} \tan^{2n} \frac{2j+1}{2(2m+1)} \pi,$$

where  $\sigma_{n,m}(E)$  denotes the determinant given by (3) in which  $\tau_j = \frac{1}{2m+1} \binom{2m+1}{2j+1}$  for  $j \in \{1, 2, \dots, n\}$ .

**Theorem 6.** For every  $m, n \in \mathbb{N}_1$  the following identity holds true:

$$\sigma_{n,m}(F) = \sum_{j=0}^{m-1} \cot^{2n} \frac{2j+1}{2(2m+1)} \pi,$$

where  $\sigma_{n,m}(F)$  denotes the determinant given by (3) in which  $\tau_j = \binom{2m+1}{2j}$  for  $j \in \{1, 2, \dots, n\}$ .

The following formulas

$$\cot^{2n} x = \left( \frac{1 - \sin^2 x}{\sin^2 x} \right)^n, \quad \tan^{2n} x = \left( \frac{1 - \cos^2 x}{\cos^2 x} \right)^n$$

yield

**Lemma 3.** *The following identities hold true:*

$$(G) \sum_{j=0}^{n-1} \binom{n}{j} (-1)^j \sin^{2j-2n} x = (-1)^{n-1} + \cot^{2n} x, \quad (n, x) \in \mathbb{N}_1 \times D_{\cot};$$

$$(H) \sum_{j=0}^{n-1} \binom{n}{j} (-1)^j \cos^{2j-2n} x = (-1)^{n-1} + \tan^{2n} x, \quad (m, x) \in \mathbb{N}_1 \times D_{\tan}.$$

**Lemma 4.** *Assume that  $n \in \mathbb{N}_1$  and  $x \in D_{\cot}$ , then*

$$\frac{1}{\sin^{2n} x} = \det \begin{pmatrix} (-1)^{n-1} + \cot^{2n} x & -\binom{n}{1} & \binom{n}{2} & -\binom{n}{3} & \dots & (-1)^{n-1} \binom{n}{n-1} \\ (-1)^{n-2} + \cot^{2n-2} x & 1 & -\binom{n-1}{1} & \binom{n-1}{2} & \dots & (-1)^{n-2} \binom{n-1}{n-2} \\ (-1)^{n-3} + \cot^{2n-4} x & 0 & 1 & -\binom{n-2}{1} & \dots & (-1)^{n-3} \binom{n-2}{n-3} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 + \cot^2 x & 0 & 0 & 0 & \dots & 1 \end{pmatrix}.$$

*Proof.* Replacing  $n$  in (G) (Lemma 3) by  $n-1, n-2, \dots, 1$ , respectively we get, together with (G), the system of  $n$  equations in  $n$  variables:

$$\frac{1}{\sin^{2n} x}, \frac{1}{\sin^{2n-2} x}, \dots, \frac{1}{\sin^2 x}.$$

Such a system is a Cramer's system and the assertion follows by the Cramer's rule.  $\square$

Using (H) in the same manner as in Lemma 4 we obtain

**Lemma 5.** *Let  $n \in \mathbb{N}_1$  and  $x \in D_{\tan}$ , then*

$$\frac{1}{\cos^{2n} x} = \det \begin{pmatrix} (-1)^{n-1} + \tan^{2n} x & -\binom{n}{1} & \binom{n}{2} & -\binom{n}{3} & \dots & (-1)^{n-1} \binom{n}{n-1} \\ (-1)^{n-2} + \tan^{2n-2} x & 1 & -\binom{n-1}{1} & \binom{n-1}{2} & \dots & (-1)^{n-2} \binom{n-1}{n-2} \\ (-1)^{n-3} + \tan^{2n-4} x & 0 & 1 & -\binom{n-2}{1} & \dots & (-1)^{n-3} \binom{n-2}{n-3} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 + \tan^2 x & 0 & 0 & 0 & \dots & 1 \end{pmatrix}.$$

To shorten notation from now on we set

$$\mu(a_n, a_{n-1}, \dots, a_1) = \det \begin{pmatrix} a_n & -\binom{n}{1} & \binom{n}{2} & -\binom{n}{3} & \dots & (-1)^{n-1} \binom{n}{n-1} \\ a_{n-1} & 1 & -\binom{n-1}{1} & \binom{n-1}{2} & \dots & (-1)^{n-2} \binom{n-1}{n-2} \\ a_{n-2} & 0 & 1 & -\binom{n-2}{1} & \dots & (-1)^{n-3} \binom{n-2}{n-3} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_1 & 0 & 0 & 0 & \dots & 1 \end{pmatrix},$$

thus the identities of Lemmas 4 and 5 can be written as

$$\frac{1}{\sin^{2n} x} = \mu \left( (-1)^{n-1} + \cot^{2n} x, (-1)^{n-2} + \cot^{2n-2} x, \dots, 1 + \cot^2 x \right) \quad (5)$$

and

$$\frac{1}{\cos^{2n} x} = \mu \left( (-1)^{n-1} + \tan^{2n} x, (-1)^{n-2} + \tan^{2n-2} x, \dots, 1 + \tan^2 x \right), \quad (6)$$

respectively.

**Theorem 7.** For every  $m \in \mathbb{N}_2$  and each  $n \in \mathbb{N}_1$  the following identity holds true:

$$\begin{aligned} \sum_{j=1}^{m-1} \sin^{-2n} \frac{\pi j}{2m} &= \sum_{j=1}^{m-1} \cos^{-2n} \frac{\pi j}{2m} \\ &= \mu \left( (-1)^{n-1} (m-1) + \sigma_{n,m}(A), (-1)^{n-2} (m-1) + \sigma_{n-1,m}(A), \right. \\ &\quad \left. \dots, (m-1) + \sigma_{1,m}(A) \right), \end{aligned} \quad (7)$$

where the numbers  $\sigma_{k,m}(A)$  for  $k \in \{1, 2, \dots, n\}$  are defined in Theorem 1.

*Proof.* In view of (5) we can write

$$\sin^{-2n} \frac{\pi j}{2m} = \mu \left( (-1)^{n-1} + \cot^{2n} \frac{\pi j}{2m}, (-1)^{n-2} + \cot^{2n-2} \frac{\pi j}{2m}, \dots, 1 + \cot^2 \frac{\pi j}{2m} \right)$$

for  $j \in \{1, 2, \dots, m-1\}$ . This by the definition of  $\mu$ , properties of determinants and Theorem 1 gives

$$\begin{aligned} \sum_{j=1}^{m-1} \sin^{-2n} \frac{\pi j}{2m} &= \mu \left( \sum_{j=1}^{m-1} \left( (-1)^{n-1} + \cot^{2n} \frac{\pi j}{2m} \right), \sum_{j=1}^{m-1} \left( (-1)^{n-2} + \cot^{2n-2} \frac{\pi j}{2m} \right), \right. \\ &\quad \left. \dots, \sum_{j=1}^{m-1} \left( 1 + \cot^2 \frac{\pi j}{2m} \right) \right) \\ &= \mu \left( (-1)^{n-1} (m-1) + \sigma_{n,m}(A), (-1)^{n-2} (m-1) + \sigma_{n-1,m}(A), \right. \\ &\quad \left. \dots, (m-1) + \sigma_{1,m}(A) \right). \end{aligned}$$

The same reasoning applies to the second identity.  $\square$

Analysis similar to that in the proof of Theorem 7 and the use of Theorems 2 – 6 give

**Theorem 8.** For every  $n, m \in \mathbb{N}_1$  the following identities holds true:

$$\begin{aligned} \sum_{j=0}^{m-1} \sin^{-2n} \frac{2j+1}{4m} \pi &= \sum_{j=0}^{m-1} \cos^{-2n} \frac{2j+1}{4m} \pi \\ &= \mu \left( (-1)^{n-1} m + \sigma_{n,m}(B), (-1)^{n-2} m + \sigma_{n-1,m}(B), \dots, m + \sigma_{1,m}(B) \right), \end{aligned} \quad (8)$$

$$\begin{aligned} \sum_{j=1}^m \sin^{-2n} \frac{\pi j}{2m+1} \\ &= \mu \left( (-1)^{n-1} m + \sigma_{n,m}(D), (-1)^{n-2} m + \sigma_{n-1,m}(D), \dots, m + \sigma_{1,m}(D) \right), \end{aligned} \quad (9)$$

$$\begin{aligned} \sum_{j=1}^m \cos^{-2n} \frac{\pi j}{2m+1} \\ &= \mu \left( (-1)^{n-1} m + \sigma_{n,m}(C), (-1)^{n-2} m + \sigma_{n-1,m}(C), \dots, m + \sigma_{1,m}(C) \right), \end{aligned} \quad (10)$$

$$\begin{aligned} \sum_{j=0}^{m-1} \sin^{-2n} \frac{(2j+1)\pi}{2(2m+1)} \\ &= \mu \left( (-1)^{n-1} m + \sigma_{n,m}(F), (-1)^{n-2} m + \sigma_{n-1,m}(F), \dots, m + \sigma_{1,m}(F) \right), \end{aligned} \quad (11)$$

$$\begin{aligned} \sum_{j=0}^{m-1} \cos^{-2n} \frac{(2j+1)\pi}{2(2m+1)} \\ &= \mu \left( (-1)^{n-1} m + \sigma_{n,m}(E), (-1)^{n-2} m + \sigma_{n-1,m}(E), \dots, m + \sigma_{1,m}(E) \right), \end{aligned} \quad (12)$$

where  $\sigma_{k,m}(B)$ ,  $\sigma_{k,m}(C)$ ,  $\sigma_{k,m}(D)$ ,  $\sigma_{k,m}(E)$ ,  $\sigma_{k,m}(F)$  for  $k \in \{1, 2, \dots, n\}$  are defined in Theorems 2 – 6.

Now we show that the general identities from Theorems 1 – 8 yield some particular equalities, including the one considered by the authors as remarkable.

**Theorem 9.** If  $m \in \mathbb{N}$ , then

$$\sum_{j=1}^{m-1} \sin^{-2} \frac{\pi j}{m} = \frac{m^2 - 1}{3}, \quad \text{provided } m \geq 2, \quad (13)$$

$$\sum_{j=1}^{m-1} \cot^2 \frac{\pi j}{m} = \frac{(m-1)(m-2)}{3}, \quad \text{provided } m \geq 2, \quad (14)$$

$$\sum_{j=0}^{m-1} \sin^{-2} \frac{(2j+1)\pi}{2m} = m^2, \quad \text{provided } m \geq 1, \quad (15)$$

*Proof.* According to Theorem 1 we have

$$\sum_{j=1}^{m-1} \tan^2 \frac{\pi j}{2m} = \sum_{j=1}^{m-1} \cot^2 \frac{\pi j}{2m} = \frac{1}{2m} \binom{2m}{3}. \quad (16)$$

On the other hand, in view of

$$\tan^2 x + \cot^2 x = \frac{4}{\sin^2 2x} - 2$$

we get

$$\sum_{j=1}^{m-1} \tan^{2n} \frac{\pi j}{2m} + \sum_{j=1}^{m-1} \cot^{2n} \frac{\pi j}{2m} = 4 \sum_{j=1}^{m-1} \sin^{-2} \frac{\pi j}{m} - 2(m-1).$$

Combining this with (16) gives

$$\sum_{j=1}^{m-1} \sin^{-2} \frac{\pi j}{m} = \frac{m^2 - 1}{3}$$

for  $m \geq 2$ . This proves (13).

To prove (14) notice that the identity

$$\cot^2 x - \frac{1}{\sin^2 x} = -1$$

yields

$$\sum_{j=1}^{m-1} \cot^2 \frac{\pi j}{m} - \sum_{j=1}^{m-1} \sin^{-2} \frac{\pi j}{m} = -(m-1), \quad m \geq 2.$$

Thus by (13) we obtain (14).

Finally we show the remarkable (15). Theorem 8 leads to

$$\sum_{j=0}^{m-1} \sin^{-2} \frac{(2j+1)\pi}{4m} = \sum_{j=0}^{m-1} \cos^{-2} \frac{(2j+1)\pi}{4m} = m + \binom{2m}{2} = 2m^2 \quad (17)$$

for  $m \geq 1$ . Since

$$\frac{1}{\sin^2 x} + \frac{1}{\cos^2 x} = \frac{4}{\sin^2 2x}$$

we have

$$\sum_{j=0}^{m-1} \sin^{-2} \frac{(2j+1)\pi}{4m} + \sum_{j=0}^{m-1} \cos^{-2} \frac{(2j+1)\pi}{4m} = 4 \sum_{j=0}^{m-1} \sin^{-2} \frac{(2j+1)\pi}{2m}, \quad m \geq 1,$$

which by (17) implies (15), and the theorem follows.  $\square$

Next we use the the identities proved here to find the sums of the series of the form  $\sum_{k=1}^{\infty} \frac{1}{k^{2n}}$ , where  $n \in \mathbb{N}_1$ . We begin with the following lemma.

**Lemma 6.** *Let  $n \in \mathbb{N}_1$ , then expression  $\sigma_{n,m}(A)$ , defined in Theorem 1, is a value of some polynomial from  $\mathbb{Q}[x]$ , where  $x = m$ . The order of such a polynomial does not exceed  $2n$ .*

*Proof.* The proof is by induction on  $n$ . For  $n = 1$  we have

$$\sigma_{1,m}(A) = \frac{1}{2m} \binom{2m}{3} = \frac{2}{3}m^2 - m - \frac{1}{3},$$

and the assertion follows. Fix  $n \geq 2$ . Assuming Lemma 6 to hold for any  $k \in \mathbb{N}_1$ ,  $k \leq n - 1$  we prove it for  $n$ . By (2),

$$\sigma_{n,m}(A) = \sum_{j=1}^{n-1} (-1)^{j-1} \tau_j \sigma_{n-j,m}(A) - (-1)^n n \tau_n,$$

where  $\tau_j = \frac{1}{2m} \binom{2m}{2j+1}$  for  $j \in \{1, 2, \dots, n\}$ . Hence by the inductive assumption  $\sigma_{n,m}(A)$  is a value of some polynomial from  $\mathbb{Q}[x]$  of order not greater than  $2n$  with  $x = m$ , as claimed.  $\square$

**Theorem 10.** *For every  $n \in \mathbb{N}_1$ ,*

$$\sum_{j=1}^{\infty} \frac{1}{j^{2n}} = \lim_{m \rightarrow \infty} \frac{\pi^{2n} \sigma_{n,m}(A)}{(2m)^{2n}},$$

where  $\sigma_{n,m}(A)$  is defined in Theorem 1.

*Proof.* Observe that

$$0 < \cot x < \frac{1}{x} < \frac{1}{\sin x} \quad x \in \left(0, \frac{\pi}{2}\right),$$

thus

$$\cot^{2n} \frac{\pi j}{2m} < \left(\frac{2m}{\pi j}\right)^{2n} < \frac{1}{\sin^{2n} \frac{\pi j}{2m}}$$

and in consequence

$$\sum_{j=1}^{m-1} \cot^{2n} \frac{\pi j}{2m} < \left(\frac{2m}{\pi}\right)^{2n} \sum_{j=1}^{m-1} \frac{1}{j^{2n}} < \sum_{j=1}^{m-1} \frac{1}{\sin^{2n} \frac{\pi j}{2m}}$$

for  $n \in \mathbb{N}_1$ ,  $m \in \mathbb{N}_2$  and  $j \in \{1, 2, \dots, n\}$ . By the definitions of  $\sigma_{n,m}(A)$  and the function  $\mu$  we have

$$\begin{aligned} \frac{\pi^{2n} \sigma_{n,m}(A)}{(2m)^{2n}} &< \sum_{j=1}^{m-1} \frac{1}{j^{2n}} < \frac{\pi^{2n}}{(2m)^{2n}} \mu((-1)^{n-1}(m-1) + \sigma_{n,m}(A), \\ &\quad (-1)^{n-2}(m-1) + \sigma_{n-1,m}(A), \\ &\quad \dots, m-1 + \sigma_{1,m}(A)). \end{aligned} \quad (18)$$

The formula for  $\mu$  and the properties of determinants give

$$\begin{aligned} & \mu((-1)^{n-1}(m-1) + \sigma_{n,m}(A), (-1)^{n-2}(m-1) + \sigma_{n-1,m}(A), \dots, m-1 + \sigma_{1,m}(A)) \\ &= (m-1)\mu((-1)^{n-1}, (-1)^{n-2}, \dots, (-1)^{n-n}) + \mu(\sigma_{n,m}(A), \sigma_{n-1,m}(A), \dots, \sigma_{1,m}(A)) \\ &= (m-1)C_1 + \sigma_{n,m}(A) + C_2\sigma_{n-1,m}(A) + \dots + C_n\sigma_{1,m}(A), \end{aligned}$$

where  $C_1, \dots, C_n$  are constants depending on  $n$ . Hence by Lemma 6 and inequality (18) we obtain

$$\lim_{m \rightarrow \infty} \frac{\pi^{2n}\sigma_{n,m}(A)}{(2m)^{2n}} \leq \sum_{j=1}^{\infty} \frac{1}{j^{2n}} \leq \lim_{m \rightarrow \infty} \frac{\pi^{2n}\sigma_{n,m}(A)}{(2m)^{2n}},$$

which establishes the formula.  $\square$

**Remark 1.** Note that in the proof Theorem 10 (the last step of the proof) we have actually proved more, namely that the order of the polynomial from Lemma 6 equals exactly  $2n$ . Indeed, if it was not true, we would have

$$\lim_{m \rightarrow \infty} \frac{\pi^{2n}\sigma_{n,m}(A)}{(2m)^{2n}} = 0$$

and consequently

$$\sum_{j=1}^{\infty} \frac{1}{j^{2n}} \leq 0,$$

which is impossible.

**Remark 2.** Treating  $\sigma_{n,m}(A)$  as a polynomial in  $m$  of order  $2n$  we have

$$\lim_{m \rightarrow \infty} \frac{\pi^{2n}\sigma_{n,m}(A)}{(2m)^{2n}} = a_{2n} \frac{\pi^{2n}}{4^n},$$

where  $a_{2n}$  denotes the leading coefficient of  $\sigma_{n,m}(A)$ . On the other hand,

$$B_{2n} \frac{2^{2n-1}\pi^{2n}}{(2n)!} (-1)^{n-1} = \sum_{j=1}^{\infty} \frac{1}{j^{2n}}, \quad n \in \mathbb{N}_1,$$

where  $B_{2n}$  stands for the  $2n$ -th Bernoulli number (see [4], p.320). Thus we get the following relation between Bernoulli numbers and the coefficients of  $\sigma_{n,m}(A)$

$$a_{2n} = B_{2n} \frac{2^{4n-1}}{(2n)!} (-1)^{n-1}, \quad n \in \mathbb{N}_1.$$

**Remark 3.** Similarly as Theorem 10 one can show that

$$\sum_{j=1}^{\infty} \frac{1}{j^{2n}} = \lim_{m \rightarrow \infty} \frac{\pi^{2n}\sigma_{n,m}(B)}{(2m)^{2n}} = \lim_{m \rightarrow \infty} \frac{\pi^{2n}\sigma_{n,m}(D)}{(2m)^{2n}} = \lim_{m \rightarrow \infty} \frac{\pi^{2n}\sigma_{n,m}(F)}{(2m)^{2n}}, \quad n \in \mathbb{N}_1.$$

where  $\sigma_{n,m}(B)$ ,  $\sigma_{n,m}(D)$  and  $\sigma_{n,m}(F)$  are defined in Theorems 2, 4 and 6, respectively.

## References

- [1] T. M. Apostol, *Another elementary proof of Euler's formula for  $\zeta(2n)$* , Amer. Math. Monthly **80** (1973), No. 4, 425-431.
- [2] P.M. Cohn, *Algebra. Vol. 1*, Second edition, John Wiley & Sons, Ltd., Chichester, 1982.
- [3] D. Cvijović, J. Klinowski, *Finite cotangent sums and the Riemann zeta function*, Math. Slovaca **50** (2000), no. 2, 149-157.
- [4] R.L. Graham, D.E. Knuth, O. Patashnik, *Concrete mathematics. A foundation for computer science*, Second edition, Addison-Wesley Publishing Company, Reading, MA, 1994.
- [5] R.A. Kortram, *Simple proofs for  $\sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}$  and  $\sin x = x \prod_{k=1}^{\infty} \left(1 - \frac{x^2}{k^2\pi^2}\right)$* . Math. Mag. **69** (1996), no. 2, 122-125.
- [6] I. Papadimitriou, *A simple proof of the formula  $\sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}$* , Amer. Math. Monthly **80** (1973), No. 4, 424-425.
- [7] T.J. Ransford, *An elementary proof of  $\sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}$* , Eureka No. 42, Summer 1982, 3-5.
- [8] H. Tsumura, *An elementary proof of Euler's formula for  $\zeta(2m)$* , Amer. Math. Monthly **111** (2004), no. 5, 430-431.

DOI: 10.7862/rf.2014.4

### Jan Górowski – Corresponding author

email: jangorowski@interia.pl

Institute of Mathematics,  
Pedagogical University,  
ul. Podchorążych 2,  
PL-30-084 Kraków, Poland

### Jerzy Żabowski

email: jerzy.zabowski@wlochkowic.pl

Paweł Włódkowic University College in Płock,  
al. Kilińskiego 12,  
PL-09-402 Płock, Poland

Received 01.10.2013, Accepted 12.01.2014



## On the zeros of an analytic function

V.K. Jain

ABSTRACT: Kuniyeda, Montel and Toya had shown that the polynomial  $p(z) = \sum_{k=0}^n a_k z^k; a_0 \neq 0$ , of degree  $n$ , does not vanish in

$$|z| \leq \{1 + (\sum_{j=1}^n |a_j/a_0|^p)^{q/p}\}^{-1/q},$$

where  $p > 1, q > 1, (1/p) + (1/q) = 1$  and we had proved that  $p(z)$  does not vanish in  $|z| \leq \alpha^{1/q}$ , where

$$\alpha = \text{unique root in } (0, 1) \text{ of } D_n x^3 - D_n S x^2 + (1 + D_n S)x - 1 = 0,$$

$$D_n = (\sum_{j=1}^n |a_j/a_0|^p)^{q/p},$$

$$S = (|a_1| + |a_2|)^q (|a_1|^p + |a_2|^p)^{-(q-1)},$$

a refinement of Kuniyeda et al.'s result under the assumption

$$D_n < (2 - S)/(S - 1).$$

Now we have obtained a generalization of our old result and proved that the function

$$f(z) = \sum_{k=0}^{\infty} a_k z^k, (\neq \text{a constant}); a_0 \neq 0,$$

analytic in  $|z| \leq 1$ , does not vanish in  $|z| < \alpha_m^{1/q}$ , where

$$\alpha_m = \text{unique root in } (0, 1) \text{ of } D x^{m+1} - D M_m x^2 + (1 + D M_m)x - 1 = 0,$$

$$D = (\sum_{k=1}^{\infty} |a_k/a_0|^p)^{q/p},$$

$$M_m = (\sum_{k=1}^m |a_k|)^q (\sum_{k=1}^m |a_k|^p)^{-q/p},$$

$m =$  any positive integer with the characteristic that there exists a positive integer  $k(\leq m)$  with  $a_k \neq 0$ .

AMS Subject Classification: *Primary 30C15, Secondary 30C10*

Keywords and Phrases: *zeros, polynomial, function analytic in  $|z| \leq 1$ , generalization, Hölder's inequality.*

## 1 Introduction and statement of results

Let

$$P(z) = b_0 + b_1z + \dots + b_nz^n$$

be a polynomial of degree  $n$ . Then according to a classical result of Kuniyeda, Montel and Toya [3, p. 124] on the location of zeros of a polynomial we have

**Theorem A.** *All the zeros of the polynomial  $P(z)$  lie in*

$$|z| < \left\{ 1 + \left( \sum_{j=0}^{n-1} |b_j/b_n|^p \right)^{q/p} \right\}^{1/q},$$

where

$$p > 1, \quad q > 1, \quad (1/p) + (1/q) = 1. \quad (1.1)$$

On applying Theorem A to the polynomial  $z^n p(1/z)$ , we have the following equivalent formulation of Theorem A.

**Theorem B.** *The polynomial*

$$p(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n; a_0 \neq 0, \quad (1.2)$$

of degree  $n$  does not vanish in

$$|z| \leq (1 + D_n)^{-1/q}, \quad (1.3)$$

where  $p, q$  are given in (1.1) and

$$D_n = \left( \sum_{j=1}^n |a_j/a_0|^p \right)^{q/p}. \quad (1.4)$$

We [2] had obtained

**Theorem C.** *All the zeros of  $P(z)$  lie in*

$$|z| < \chi^{1/q},$$

where  $\chi$  is the unique root of the equation

$$x^3 - (1 + LM)x^2 + LMx - L = 0,$$

in  $(1, \infty)$ ,

$$L = \left( \sum_{j=0}^{n-1} |b_j/b_n|^p \right)^{q/p},$$

$$M = (|b_{n-1}| + |b_{n-2}|)^q (|b_{n-1}|^p + |b_{n-2}|^p)^{-(q-1)}.$$

Theorem C is a refinement of Theorem A, under the assumption

$$L < (2 - M)/(M - 1).$$

The equivalent formulation of Theorem C, (similar to the formulation of Theorem B from Theorem A) is

**Theorem D.** *The polynomial*

$$p(z) = a_0 + a_1z + \dots + a_nz^n; a_0 \neq 0,$$

of degree  $n$  does not vanish in

$$|z| \leq \alpha^{1/q},$$

where  $\alpha$  is the unique root of the equation

$$D_n x^3 - D_n S x^2 + (1 + D_n S)x - 1 = 0,$$

in  $(0, 1)$ ,

$$S = (|a_1| + |a_2|)^q (|a_1|^p + |a_2|^p)^{-(q-1)},$$

and  $D_n$  is as in Theorem B.

Theorem D is a refinement of Theorem B, under the assumption

$$D_n < (2 - S)/(S - 1).$$

In this paper we have obtained a generalization of Theorem D for the functions, analytic in  $|z| \leq 1$ . More precisely we have proved

**Theorem 1.** *Let*

$$f(z) = \sum_{k=0}^{\infty} a_k z^k, (\neq \text{a constant}); a_0 \neq 0, \tag{1.5}$$

be analytic in  $|z| \leq 1$ . Then  $f(z)$  does not vanish in

$$|z| < \alpha_m^{1/q}, \tag{1.6}$$

where

$$\begin{aligned} q &> 1, \quad p > 1, \quad (1/p) + (1/q) = 1, \\ m &= \text{any positive integer with the characteristic that} \end{aligned} \tag{1.7}$$

$$\begin{aligned} &\text{there exists a positive integer } k(\leq m) \text{ with } a_k \neq 0, \\ \alpha_m &= \text{unique root in } (0, 1), \text{ of} \\ &\{g(x) \equiv\}, \quad D x^{m+1} - D M_m x^2 + (1 + D M_m)x - 1 = 0, \end{aligned} \tag{1.8}$$

$$D = \left( \sum_{k=1}^{\infty} |a_k/a_0|^p \right)^{q/p}, (> 0, \text{ by (1.5)}), \tag{1.9}$$

$$M_m = \left( \sum_{k=1}^m |a_k| \right)^q \left( \sum_{k=1}^m |a_k|^p \right)^{-q/p}, (> 0, \text{ by (1.7)}). \tag{1.10}$$

From Theorem 1 we easily get

**Corollary 1.** *Under the same hypothesis as in Theorem 1,  $f(z)$  does not vanish in*

$$|z| < \sup_{m \geq M, q > 1} \alpha_m^{1/q},$$

where

$$M = \text{least positive integer } k \text{ such that } a_k \neq 0.$$

## 2 Lemmas

For the proof of the theorem, we require the following lemmas.

**Lemma 1.** *Let*

$$\begin{aligned} \alpha_j > 0, \quad \beta_j > 0, \quad \text{for } j = 1, 2, \dots, n, \\ q > 1, \quad p > 1, \quad (1/p) + (1/q) = 1, \\ 1 \leq m < n. \end{aligned}$$

Then

$$\sum_{j=1}^n \alpha_j \beta_j \leq \left( \left( \sum_{j=1}^n \beta_j^p \right)^{1/p} \left( \sum_{j=1}^m \beta_j^p \right)^{-1/p} \right) \left\{ \left( \sum_{j=1}^m \alpha_j \beta_j \right)^q + \left( \sum_{j=1}^m \beta_j^p \right)^{q-1} \left( \sum_{j=m+1}^n \alpha_j^q \right) \right\}^{1/q}. \quad (2.1)$$

This lemma is due to Beckenbach [1].

From Lemma 1 we easily obtain

**Lemma 2.** *Inequality (2.1) is true even if*

$$\begin{aligned} \alpha_j \geq 0, \quad j = 1, 2, \dots, n, \\ \beta_j \geq 0, \quad j = 1, 2, \dots, n, \end{aligned}$$

with

$$\beta_j \neq 0, \text{ for at least one } j, 1 \leq j \leq m.$$

**Lemma 3.** *The equation*

$$Dx^{m+1} - DM_m x^2 + (1 + DM_m)x - 1 = 0 \quad (2.2)$$

has a unique root  $\alpha_m$  in  $(0, 1)$  where  $m$ ,  $D$  and  $M_m$  are as in Theorem 1.

*Proof of Lemma 3.* We firstly assume that

$$m > 1.$$

Now we consider the transformation

$$x = 1/t$$

in equation (2.2), thereby giving the transformed equation

$$t^{m+1} - (1 + DM_m)t^m + DM_mt^{m-1} - D = 0, \quad (2.3)$$

and then the transformation

$$t = 1 + y$$

in (2.3), thereby giving the transformed equation

$$(1 + y)^{m+1} - (1 + DM_m)(1 + y)^m + DM_m(1 + y)^{m-1} - D = 0, \quad (2.4)$$

i.e.

$$\begin{aligned} & y^{m+1} + y^m((m/1) - DM_m) + ((m-1)/1!)((m/2) - DM_m)y^{m-1} \\ & + ((m-1)(m-2)/2!)((m/3) - DM_m)y^{m-2} + \dots \\ & + ((m-1)(m-2)\dots(m-j+1)/(j-1)!)((m/j) - DM_m)y^{m+1-j} + \dots \\ & + ((m-1)(m-2)\dots(m-m+1)/(m-1)!)((m/m) - DM_m)y - D \\ & = 0. \end{aligned} \quad (2.5)$$

By using Descartes's rule of signs we can say that equation (2.5) (i.e. equation (2.4)) will have a unique positive root and accordingly the equation (2.3) will have a unique root in  $(1, \infty)$ . Hence the equation (2.2) will have a unique root  $\alpha_m$ , (say), in  $(0, 1)$ , thereby proving Lemma 3 for the possibility under consideration.

For the possibility

$$m = 1,$$

the transformed equation, similar to equation (2.5), (i.e. equation (2.4)), is

$$y^2 + y(1 - DM_m) - D = 0.$$

Now Lemma 3 follows for this possibility, by using arguments similar to those used for proving Lemma 3 for the possibility

$$m > 1.$$

This completes the proof of Lemma 3.

### 3 Proof of Theorem 1

Let

$$f_n(z) = \sum_{k=0}^n a_k z^k, n = 1, 2, 3, \dots .$$

Then for  $|z| < 1$  and  $n > m$

$$\begin{aligned}
|f_n(z)| &\geq |a_0| - \sum_{k=1}^n |z|^k |a_k|, \\
&\geq |a_0| - \left\{ \left( \sum_{k=1}^n |a_k|^p \right)^{1/p} \left( \sum_{k=1}^m |a_k|^p \right)^{-1/p} \right\} \left[ \left( \sum_{k=1}^m |z|^k |a_k| \right)^q \right. \\
&\quad \left. + \left\{ \left( \sum_{k=1}^m |a_k|^p \right)^{q-1} \right\} \left( \sum_{k=m+1}^n |z|^{kq} \right) \right]^{1/q}, \text{ (by Lemma 2),} \\
&\geq |a_0| - \left( \sum_{k=1}^n |a_k|^p \right)^{1/p} \left[ \left( \sum_{k=1}^m |a_k| |z|^k \right)^q \left( \sum_{k=1}^m |a_k|^p \right)^{-q/p} \right. \\
&\quad \left. + \left( \sum_{k=m+1}^n |z|^{kq} \right) \right]^{1/q}, \text{ (by 1.1),} \\
&\geq |a_0| - \left( \sum_{k=1}^n |a_k|^p \right)^{1/p} \left[ M_m |z|^q + \left( \sum_{k=m+1}^n |z|^{kq} \right) \right]^{1/q}, \text{ (by 1.10),}
\end{aligned}$$

which, by making

$$n \rightarrow \infty,$$

implies that

$$\begin{aligned}
|f(z)| &\geq |a_0| - \left( \sum_{k=1}^{\infty} |a_k|^p \right)^{1/p} \left[ M_m |z|^q + \left( \sum_{k=m+1}^{\infty} |z|^{kq} \right) \right]^{1/q}, \text{ (} \sum_{k=1}^{\infty} |a_k|^p \text{ will converge} \\
&\quad \text{as } \sum_{k=1}^{\infty} |a_k| \text{ converges and } \left( \sum_{k=1}^n |a_k|^p \right)^{1/p} \leq \sum_{k=1}^n |a_k|, n = 1, 2, \dots), \\
&= |a_0| \left[ 1 - \{ D(M_m |z|^q + (|z|^{(m+1)q} / (1 - |z|^q))) \}^{1/q} \right], \text{ (by 1.9),} \\
&> 0, \tag{3.1}
\end{aligned}$$

if

$$D|z|^{(m+1)q} - DM_m |z|^{2q} + (1 + DM_m) |z|^q - 1 < 0. \tag{3.2}$$

Now as

$$g(0) = -1, \text{ (by (1.8)),}$$

we can say by Lemma 3, (3.1) and (3.2) that

$$|f(z)| > 0,$$

if

$$|z|^q < \alpha_m,$$

thereby proving Theorem 1.

**Remark 1.** Theorem 1 gives better bound than that given by the result, that  $f(z)$  does not vanish in

$$|z| < \{1/(1 + D)\}^{1/q},$$

obtained by using Hölder's inequality instead of Lemma 2 and following the method of proof of Theorem 1, provided

$$\begin{aligned} m = 1 & \quad \& \quad M_m < m, \\ m \geq 2 & \quad \& \quad M_m \leq 1, \\ m \geq 2, 1 < M_m < m & \quad \text{and} \quad D < D_0, \end{aligned} \tag{3.3}$$

where  $D_0$  is the unique positive root of the equation

$$\begin{aligned} (M_m - 1)D^{m-1} & + (m - 1)(M_m - (m/(m - 1)))D^{m-2} \\ & + ((m - 1)(m - 2)/2)(M_m - (m/(m - 2)))D^{m-3} \\ & + \dots + (m - 1)(M_m - (m/2))D + (M_m - m) \\ = 0, & \quad (m \geq 2 \& 1 < M_m < m), \end{aligned}$$

as for  $m = 1 \& M_m < m$

$$g(1/(1 + D)) < 0,$$

and for  $m \geq 2$

$$g(1/(1 + D)) < 0,$$

is equivalent to

$$\begin{aligned} (M_m - 1)D^{m-1} & + (m - 1)(M_m - (m/(m - 1)))D^{m-2} \\ & + ((m - 1)(m - 2)/2)(M_m - (m/(m - 2)))D^{m-3} \\ & + \dots + (m - 1)(M_m - (m/2))D + (M_m - m) \\ < 0. \end{aligned}$$

The function

$$f(z) = 1 + z + (z/(2i))^3 + (z/(2i))^4 + (z/(2i))^5 + \dots$$

satisfies (3.3) with

$$p = q = m = 2$$

and the corresponding  $\alpha_m^{1/q}$  is .752.

## References

- [1] Beckenbach, E. F., On Hölder's inequality, *J. Math. Anal. Appl.* 15 (1966), 21–29.
- [2] Jain, V. K., On the zeros of a polynomial, *Publ. L'inst. Math. (Nouvelle série)* 41 (1987), 75–77.

- [3] Marden, M., The geometry of polynomials, *A. M. S. Surveys* 3 (1966), New York.

**DOI: 10.7862/rf.2014.5**

**V.K. Jain**

email: [vkj@maths.iitkgp.ernet.in](mailto:vkj@maths.iitkgp.ernet.in)

Mathematics Department,  
I.I.T. Kharagpur - 721302, INDIA

*Received 31.12.2013, Accepted 10.06.2014*



## On circularly symmetric functions

*Leopold Koczan, Paweł Zaprawa*

ABSTRACT: Let  $D \subset \mathbb{C}$  and  $0 \in D$ . A set  $D$  is circularly symmetric if for each  $\varrho \in \mathbb{R}^+$  a set  $D \cap \{\zeta \in \mathbb{C} : |\zeta| = \varrho\}$  is one of three forms: an empty set, a whole circle, a curve symmetric with respect to the real axis containing  $\varrho$ . A function  $f \in \mathcal{A}$  is circularly symmetric if  $f(\Delta)$  is a circularly symmetric set. The class of all such functions we denote by  $X$ . The above definitions were given by Jenkins in [2].

In this paper besides  $X$  we also consider some of its subclasses:  $X(\lambda)$  and  $Y \cap S^*$  consisting of functions in  $X$  with the second coefficient fixed and univalent starlike functions respectively. According to the suggestion, in Abstract we add one more paragraph at the end of the section:

For  $X(\lambda)$  we find the radii of starlikeness, starlikeness of order  $\alpha$ , univalence and local univalence. We also obtain some distortion results. For  $Y \cap S^*$  we discuss some coefficient problems, among others the Fekete-Szegő inequalities.

AMS Subject Classification: *30C45*

Keywords and Phrases: *symmetric function, radius of starlikeness, zeros of polynomials*

### 1 The class of circularly symmetric functions and some its subclasses.

Let  $\tilde{\mathcal{A}}$  denote the class of all functions analytic in  $\Delta \equiv \{\zeta \in \mathbb{C} : |\zeta| < 1\}$  and let  $\mathcal{A}$  denote the class of all functions analytic in  $\Delta$  normalized by  $f(0) = f'(0) - 1 = 0$ . Similar notation is applied to the class of typically real functions, i.e. functions satisfying the following condition:  $\operatorname{Im} z \operatorname{Im} f(z) \geq 0$  for  $z \in \Delta$ . The set of all analytic and typically real functions is denoted by  $\tilde{T}$ ; the subset of  $\tilde{T}$  consisting of normalized functions is denoted by  $T$ . Hence  $T = \tilde{T} \cap \mathcal{A}$ . It follows from the definition of a typically real function that  $z \in \Delta^+ \Leftrightarrow f(z) \in \mathbb{C}^+$  and  $z \in \Delta^- \Leftrightarrow f(z) \in \mathbb{C}^-$ . The symbols  $\Delta^+$ ,  $\Delta^-$ ,  $\mathbb{C}^+$ ,  $\mathbb{C}^-$  mean the following open sets: the upper and the lower half of the unit disk  $\Delta$  and the upper and the lower halfplane.

In this paper we focus on so called circularly symmetric functions, which were defined by Jenkins in [2]. Let us start with the following definitions.

Let  $D \subset \mathbb{C}$  and  $0 \in D$ .

*Definition 1.* A set  $D$  is circularly symmetric if for each  $\varrho \in \mathbb{R}^+$  a set  $D \cap \{\zeta \in \mathbb{C} : |\zeta| = \varrho\}$  is one of three forms: an empty set, a whole circle, a curve symmetric with respect to the real axis containing  $\varrho$ .

*Definition 2.* A function  $f \in \mathcal{A}$  is circularly symmetric if  $f(\Delta)$  is a circularly symmetric set. The class of all such functions we denote by  $X$ .

In fact, Jenkins claimed more than it was stated in the above definition. He considered only these circularly symmetric functions which are univalent. This assumption is rather restrictive. Furthermore, there are no objections to reject it. The number of interesting problems appear while discussing non-univalent circularly symmetric functions. For these reasons we decided to define a circularly symmetric function as in Definition 2. In order to distinguish the classes of non-univalent and univalent circularly symmetric functions we will denote the latter by  $Y$ .

Besides  $X$  we will also consider some of its subclasses:  $X(\lambda)$  and  $Y \cap S^*$  consisting of functions in  $X$  with the fixed second coefficient of the Taylor series expansion and univalent starlike functions respectively. As it was shown in [2], for all  $r \in (0, 1)$  and for a circularly symmetric function  $f$  the expression  $|f(re^{i\varphi})|$  is a nonincreasing function for  $\varphi \in (0, \pi)$  and a nondecreasing function for  $\varphi \in (\pi, 2\pi)$ . From this fact and the equality

$$-\frac{\partial}{\partial \varphi} (\log |f(re^{i\varphi})|) = \operatorname{Im} \left( re^{i\varphi} \frac{f'(re^{i\varphi})}{f(re^{i\varphi})} \right)$$

it follows that on the circle  $|z| = r$  there is

$$\operatorname{Im} \frac{zf'(z)}{f(z)} \geq 0 \quad \text{if and only if} \quad \operatorname{Im} z \geq 0.$$

Hence

**Theorem 1.** [2]

$$f \in X \Leftrightarrow \frac{zf'(z)}{f(z)} \in \tilde{T}.$$

The condition  $\frac{zf'(z)}{f(z)} \in \tilde{T}$  is not sufficient for univalence of  $f$ . We have only

**Theorem 2.** If  $f \in Y$  then  $\frac{zf'(z)}{f(z)} \in \tilde{T}$ .

According to Theorem 1, all coefficients of the Taylor expansion of  $f \in X$  are real. Some other results concerning  $Y$  one can find in [1] and [4].

Similar, but more general, functions were discussed by Libera in [3]. He considered so called disk-like functions. The functions  $f$  of this class have the property: there exists a number  $\varrho$  depending on  $f$  that for each fixed  $r$ ,  $r \in (\varrho, 1]$ , there exist numbers  $\varphi_1, \varphi_2$  depending on  $r$  that  $|f(re^{i\varphi})|$  is decreasing if  $\varphi$  increases in some interval  $I_1 = (\varphi_1, \varphi_2)$  and increasing in  $I_2 = (\varphi_2, \varphi_1 + 2\pi)$ . The class of these functions Libera

denoted by  $\mathcal{D}$ . In particular, if  $f$  has real coefficients and  $|f(re^{i\varphi})|$  is increasing on the lower half of the circle  $|z| = r$  and is decreasing on the upper half of this circle, then  $f$  is a circularly symmetric function. Although  $\mathcal{D}$  is more general than  $X$ , some of the results of the paper [3] are still valid for the class  $X$ .

Let us assume that  $f$  is of the form  $f(z) = z + \lambda z^2 + \dots$ . From Theorem 1 it follows that a function

$$\frac{1}{\lambda} \left( \frac{zf'(z)}{f(z)} - 1 \right)$$

is in  $T$ ; let us denote it by  $h(z)$ . Hence

$$(1) \quad f(z) = z \exp \left( \lambda \int_0^z \frac{h(\zeta)}{\zeta} d\zeta \right).$$

Applying the very well known relation between  $T$  and  $CVR(i)$  consisting of functions with real coefficients  $g$  which are convex in the direction of the imaginary axis and normalized by  $g(0) = g'(0) - 1 = 0$ , we obtain

**Corollary 1.**

$$(2) \quad f \in X \Leftrightarrow f(z) = z \exp \{ \lambda g(z) \}, \quad g \in CVR(i), \quad \lambda > 0.$$

The conclusion similar to the above corollary one can find in the paper of Libera (corollary on page 253).

Basing on the equivalence (2) we can define the subclass of  $X$  containing these circularly symmetric functions for which the second coefficient is fixed and equal to  $\lambda \geq 0$ . We denote this class by  $X(\lambda)$ . For  $\lambda = 0$  the set  $X(0)$  has only one element - the identity function. We shall present the properties of  $X(\lambda)$  in next section.

## 2 Properties of $X(\lambda)$ .

**Theorem 3.** *The radius of starlikeness for  $X(\lambda)$  is equal to  $r_{S^*}(X(\lambda)) = r_\lambda$ , where  $r_\lambda = \frac{1}{4} \left( \sqrt{\lambda + 4} - \sqrt{\lambda} \right)^2$ . The extremal function is  $f_\lambda(z) = z \exp \left( \lambda \frac{z}{1+z} \right)$ .*

*Proof*

It follows from (1) that  $\frac{zf'(z)}{f(z)} = 1 + \lambda z g'(z) = 1 + \lambda h(z)$ , where  $g \in CVR(i)$ ,  $h \in T$ . The well-known estimate of the real part of a typically real function leads to

$$\operatorname{Re} \frac{zf'(z)}{f(z)} \geq 1 - \lambda \frac{r}{(1-r)^2}.$$

Therefore,  $\operatorname{Re} \frac{zf'(z)}{f(z)} \geq 0$  if and only if  $r \leq \frac{2}{2 + \lambda + \sqrt{\lambda^2 + 4\lambda}}$ , or equivalently, if  $r \leq r_\lambda$ . Equality in the above estimate holds for  $h(z) = \frac{z}{(1+z)^2}$  and  $z = -r$ . It means that the extremal function is  $f_\lambda$ . ■

The result of Theorem 3 can be generalized in order to finding the radius of starlikeness of order  $\alpha$ ,  $\alpha \in [0, 1)$ . It suffices to replace the inequality  $1 - \lambda \frac{r}{(1-r)^2} \geq 0$  by  $1 - \lambda \frac{r}{(1-r)^2} \geq \alpha$ . Hence

**Theorem 4.** *The radius of starlikeness of order  $\alpha$ ,  $\alpha \in [0, 1)$  for  $X(\lambda)$  is equal to  $r_{S^*(\alpha)}(X(\lambda)) = \frac{1}{4} \left( \sqrt{\frac{\lambda}{1-\alpha} + 4} - \sqrt{\frac{\lambda}{1-\alpha}} \right)^2$ . The extremal function is  $f_\lambda(z) = z \exp\left(\lambda \frac{z}{1+z}\right)$ .*

Observe that for all  $f \in X(\lambda)$  the condition  $\frac{zf'(z)}{f(z)} \neq 0$  holds if only  $z \in \Delta_{r_\lambda}$ . Moreover, for  $f_\lambda$  and  $z = -r_\lambda$  there is

$$\frac{zf'(z)}{f(z)} \Big|_{z=-r_\lambda} = \left( 1 + \lambda \frac{z}{(1+z)^2} \right) \Big|_{z=-r_\lambda} = 1 - \lambda \frac{r_\lambda}{(1-r_\lambda)^2} = 0.$$

This results in

**Theorem 5.** *The radius of local univalence for  $X(\lambda)$  is equal to  $r_{LU}(X(\lambda)) = r_\lambda$ .*

Because of  $r_{S^*} \leq r_S \leq r_{LU}$ , which is true for any class of analytic functions, we obtain

**Corollary 2.** *The radius of univalence for  $X(\lambda)$  is equal to  $r_S(X(\lambda)) = r_\lambda$ .*

It is known that the second coefficients of the Taylor expansion of functions in the following subclasses of  $\mathcal{A}$  consisting of: convex functions, univalent functions and locally univalent functions have the upper bounds: 1, 2, 4 respectively. For this reason it is worth observing that

$$r_S(X(1)) = \frac{1}{2}(3 - \sqrt{5}), \quad r_S(X(2)) = 2 - \sqrt{3}, \quad r_S(X(4)) = (\sqrt{2} - 1)^2.$$

**Theorem 6.** *If  $f \in X(\lambda)$  and  $r = |z| \in (0, 1)$  then*

$$(3) \quad r \exp\left(\frac{-\lambda r}{1-r}\right) \leq |f(z)| \leq r \exp\left(\frac{\lambda r}{1-r}\right),$$

*Equalities in the above estimates hold for  $f(z) = z \exp\left(\frac{\lambda z}{1+z}\right)$ ,  $z = -r$  and  $f(z) = z \exp\left(\frac{\lambda z}{1-z}\right)$ ,  $z = r$  respectively.*

*Proof*

For  $g \in CVR(i)$  the exact estimate holds (see for example [5])

$$(4) \quad |\operatorname{Re} g(z)| \leq \frac{r}{1-r},$$

with equality for  $g(z) = \frac{z}{1+z}$ ,  $z = -r$  and  $g(z) = \frac{z}{1-z}$ ,  $z = r$  respectively. From Corollary 1 it follows that

$$|f(z)| = |z| \exp(\lambda \operatorname{Re} g(z)).$$

Combining it with (4) completes the proof.  $\blacksquare$

**Theorem 7.** *If  $f \in X(\lambda)$  and  $r = |z| \in (0, 1)$  then*

$$(5) \quad |f'(z)| \leq \left(1 + \lambda \frac{r}{(1-r)^2}\right) \exp\left(\frac{\lambda r}{1-r}\right),$$

*Equality in the above estimate holds for  $f(z) = z \exp\left(\frac{\lambda z}{1-z}\right)$  and  $z = r$ .*

*Proof*

From Corollary 1 and (1) we have

$$|f'(z)| = \left|\frac{f(z)}{z}\right| |1 + \lambda h(z)|,$$

where  $h \in T$ . Applying Theorem 6 and the estimate of the modulus of a function in  $T$  in the above equality leads to the assertion.  $\blacksquare$

According to Theorems 6 and 7, both  $|f(z)|$  and  $|f'(z)|$  can be arbitrarily large while considering functions in the whole class  $X$ , not only functions with the second coefficient fixed.

### 3 Properties of $Y \cap S^*$ .

In the paper of Szapiel [4] one can find the relation between the class of circularly symmetric functions which are starlike with the class of typically real functions  $T$ :

**Theorem 8.**

$$f \in Y \cap S^* \Leftrightarrow \frac{zf'(z)}{f(z)} \in \tilde{T} \cap P_R.$$

Szapiel also proved the representation formula for functions in the class  $\mathcal{R}^2 = \{q \in \mathcal{A} : q = p^2, p \in \tilde{T} \cap P_R\}$ . Namely,  $q \in \mathcal{R}^2$  if and only if

$$(6) \quad q(z) = \int_{-1}^1 \frac{(1+z)^2}{1-2zt+z^2} d\mu(t).$$

From this formula one can establish the relationship between  $\mathcal{R}^2$  and  $T$ :

$$(7) \quad q \in \mathcal{R}^2 \Leftrightarrow g \in T,$$

where

$$q(z) = (1+z)^2 \frac{g(z)}{z}.$$

From the above we get

**Corollary 3.**

$$(8) \quad f \in Y \cap S^* \Leftrightarrow \frac{zf'(z)}{f(z)} = (1+z) \sqrt{\frac{g(z)}{z}}, \quad g \in T.$$

**Examples.** Putting functions of the class  $T$  into (8) we obtain associated functions from  $Y \cap S^*$ :

1. If  $g(z) = z$ , then  $\frac{zf'(z)}{f(z)} = 1 + z$  and hence  $f(z) = ze^z$ .
2. If  $g(z) = \frac{z}{(1+z)^2}$ , then  $\frac{zf'(z)}{f(z)} = 1$  and hence  $f(z) = z$ .
3. If  $g(z) = \frac{z}{(1-z)^2}$ , then  $\frac{zf'(z)}{f(z)} = \frac{1+z}{1-z}$  and hence  $f(z) = \frac{z}{(1-z)^2}$ .

Many of the properties of  $Y \cap S^*$  follow directly from obvious inclusion  $Y \cap S^* \subset S^*$  and the fact that the Koebe function  $f(z) = \frac{z}{(1-z)^2}$ , which is starlike, belongs also to  $Y \cap S^*$ . This observation gives us the following sharp results:

1. If  $f \in Y \cap S^*$  and  $f(z) = \sum_{n=1}^{\infty} a_n z^n$ , then  $|a_n| \leq n$ .
2. If  $f \in Y \cap S^*$  and  $r = |z| \in (0, 1)$ , then  $\frac{r}{(1+r)^2} \leq |f(z)| \leq \frac{r}{(1-r)^2}$ .
3. If  $f \in Y \cap S^*$  and  $r = |z| \in (0, 1)$ , then  $\frac{1-r}{(1+r)^3} \leq |f'(z)| \leq \frac{1+r}{(1-r)^3}$ .
4. Every function in  $Y \cap S^*$  is convex in the disk  $|z| < 2 - \sqrt{3}$ .
5. Every function in  $Y \cap S^*$  is strongly starlike of order  $\alpha$  in the disk  $|z| < \tan(\alpha \frac{\pi}{4})$ .

Now we shall find the lower bounds of the second and the third coefficients in  $Y \cap S^*$ . Let  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in Y \cap S^*$  and  $g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in T$ . From (8) we conclude

$$(9) \quad 2a_2 = b_2 + 2,$$

$$(10) \quad 4a_3 - a_2^2 = b_3 + 2b_2 + 1.$$

Let us denote by  $A_{2,3}(A)$  a set  $\{(a_2(f), a_3(f)) : f \in A\}$ . This set for  $T$  is known:  $A_{2,3}(T) = \{(x, y) : -2 \leq x \leq 2, x^2 - 1 \leq y \leq 3\}$ . This results in the following bound for a function in  $Y \cap S^*$ :

$$(11) \quad 0 \leq a_2 \leq 2.$$

Taking into account (9) and (10) in  $A_{2,3}(T)$  we obtain

**Theorem 9.**

$$A_{2,3}(Y \cap S^*) = \left\{ (x, y) : 0 \leq x \leq 2, \frac{1}{4}(5x^2 - 4x) \leq y \leq \frac{1}{4}(x^2 + 4x) \right\}.$$

Consequently

**Corollary 4.** Let  $f \in Y \cap S^*$  have the Taylor series expansion  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ . Then  $-\frac{1}{5} \leq a_3 \leq 3$ .

From (10) and  $A_{2,3}(T)$  it follows that

$$4a_3 - a_2^2 = b_3 + 2b_2 + 1 \geq b_2^2 + 2b_2 \geq -1$$

and

$$4a_3 - a_2^2 = b_3 + 2b_2 + 1 \leq b_2^2 + 4 \leq 8.$$

Hence

**Theorem 10.** Let  $f \in Y \cap S^*$  have the Taylor series expansion  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ . Then  $-\frac{1}{4} \leq a_3 - \frac{1}{4}a_2^2 \leq 2$ .

The points of intersection of two parabolas from Theorem 9 coincide with two pairs of coefficients  $(a_2, a_3)$  of the functions  $f_1(z) = z$  and  $f_2(z) = \frac{z}{(1-z)^2}$ . From Theorem 10 it follows that the class  $Y \cap S^*$  is not a convex set because the set  $A_{2,3}(Y \cap S^*)$  is not convex.

Basing on Theorem 9 one can derive so called the Fekete-Szegő inequalities for  $Y \cap S^*$ .

**Theorem 11.** Let  $f \in Y \cap S^*$  be of the form  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ . Then

$$\begin{cases} \frac{1}{4\mu-5} & \mu \leq 1 \\ 3-4\mu & \mu \geq 1 \end{cases} \leq a_3 - \mu a_2^2 \leq \begin{cases} 3-4\mu & \mu \leq \frac{1}{2} \\ \frac{1}{4\mu-1} & \mu \geq \frac{1}{2} \end{cases}.$$

*Proof*

Assume that  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in Y \cap S^*$ . Let us denote by  $Q$  a function  $Q(a_2, a_3) = a_3 - \mu a_2^2$ . With a fixed  $\mu \in \mathbb{R}$  the function  $Q$  achieves its extremal value on the boundary of the set  $A_{2,3}(Y \cap S^*)$ .

Let us consider two functions

$$Q_1(x) = Q(x, \frac{1}{4}x^2 + x) = x^2 \left( \frac{1}{4} - \mu \right) + x$$

and

$$Q_2(x) = Q(x, \frac{5}{4}x^2 - x) = x^2 \left( \frac{5}{4} - \mu \right) - x.$$

For  $x \in [0, 2]$  the inequality

$$Q_1(x) \geq Q_2(x)$$

holds; hence

$$\begin{aligned} (12) \quad \max\{Q : f \in Y \cap S^*\} &= \max\{Q(x, y) : (x, y) \in A_{2,3}(Y \cap S^*)\} \\ &= \max\{Q_1(x) : x \in [0, 2]\} \end{aligned}$$

and

$$\begin{aligned} (13) \quad \min\{Q : f \in Y \cap S^*\} &= \min\{Q(x, y) : (x, y) \in A_{2,3}(Y \cap S^*)\} \\ &= \min\{Q_2(x) : x \in [0, 2]\}. \end{aligned}$$

The function  $Q_1$  for  $\mu \leq \frac{1}{2}$  is strictly increasing in  $[0, 2]$ ; thus  $\max\{Q_1(x) : x \in [0, 2]\} = Q_1(2)$ . For  $\mu > \frac{1}{2}$  the function  $Q_1$  increases in  $(0, x_1)$  and decreases in  $(x_1, 2)$ , where  $x_1 = \frac{2}{4\mu-1}$ . This results in  $\max\{Q_1(x) : x \in [0, 2]\} = Q_1(x_1)$ .

Similarly, the function  $Q_2$  for  $\mu < 1$  decreases in  $(0, x_2)$  and increases in  $(x_2, 2)$ , where  $x_2 = \frac{2}{5-4\mu}$ . Hence  $\min\{Q_2(x) : x \in [0, 2]\} = Q_2(x_2)$ . For  $\mu \geq 1$  the function  $Q_2$  is strictly decreasing in  $[0, 2]$ , so  $\min\{Q_2(x) : x \in [0, 2]\} = Q_2(2)$ . ■

Taking  $\mu = 0$  or  $\mu = \frac{1}{4}$  we obtain previously obtained results from Corollary 4 and from Theorem 10.

## References

- [1] Q. Deng, On circularly symmetric functions, Appl. Math. Lett. 23 No. 12 (2010) 1483-1488.
- [2] J.A. Jenkins, On circularly symmetric functions, Proc. Amer. Math. Soc. 6 (1955) 620-624.
- [3] R.J. Libera, Disk-like functions, J. Aust. Math. Soc. 11 (1970) 251-256.
- [4] M. Szapiel, W. Szapiel, Extreme points of convex sets (IV). Bounded typically real functions, Bull. Acad. Polon. Sci., Math. 30 (1982) 49-57.
- [5] P. Zaprawa, Distortion theorems for certain subclasses of typically real functions, Stud. Univ. Babes-Bolyai Math. 57 (2012) 301308.

**DOI: 10.7862/rf.2014.6**

**Leopold Koczan**

email: l.koczan@pollub.pl

**Paweł Zaprawa - corresponding author**

email: p.zaprawa@pollub.pl

Department of Mathematics

Lublin University of Technology

Nadbystrzycka 38D, 20-618 Lublin, Poland

*Received 25.06.2013, Accepted 14.08.2013*



# Properties of higher order differential polynomials generated by solutions of complex differential equations in the unit disc

*Zinelâabidine Latreuch, Benharrat Belaïdi*

ABSTRACT: The main purpose of this paper is to study the controllability of solutions of the differential equation

$$f^{(k)} + A_{k-1}(z) f^{(k-1)} + \dots + A_1(z) f' + A_0(z) f = 0.$$

In fact, we study the growth and oscillation of higher order differential polynomial with meromorphic coefficients in the unit disc  $\Delta = \{z : |z| < 1\}$  generated by solutions of the above  $k^{th}$  order differential equation.

AMS Subject Classification: *34M10, 30D35.*

Keywords and Phrases: *Iterated  $p$ -order, Linear differential equations, Iterated exponent of convergence of the sequence of distinct zeros, Unit disc.*

## 1 Introduction and main results

Throughout this paper, we assume that the reader is familiar with the fundamental results and the standard notations of the Nevanlinna's value distribution theory on the complex plane and in the unit disc  $\Delta = \{z : |z| < 1\}$  (see [13], [14], [18], [20]). We need to give some definitions and discussions. Firstly, let us give two definitions about the degree of small growth order of functions in  $\Delta$  as polynomials on the complex plane  $\mathbb{C}$ . There are many types of definitions of small growth order of functions in  $\Delta$  (see [10], [11]).

**Definition 1.1** ([10], [11]) Let  $f$  be a meromorphic function in  $\Delta$ , and

$$D(f) := \limsup_{r \rightarrow 1^-} \frac{T(r, f)}{\log \frac{1}{1-r}} = b.$$

If  $b < \infty$ , we say that  $f$  is of finite  $b$  degree (or is non-admissible). If  $b = \infty$ , we say that  $f$  is of infinite degree (or is admissible), both defined by characteristic function  $T(r, f)$ .

**Definition 1.2** ([10], [11]) Let  $f$  be an analytic function in  $\Delta$ , and

$$D_M(f) := \limsup_{r \rightarrow 1^-} \frac{\log^+ M(r, f)}{\log \frac{1}{1-r}} = a < \infty \text{ (or } a = \infty),$$

then we say that  $f$  is a function of finite  $a$  degree (or of infinite degree) defined by maximum modulus function  $M(r, f) = \max_{|z|=r} |f(z)|$ .

Now we give the definitions of iterated order and growth index to classify generally the functions of fast growth in  $\Delta$  as those in  $\mathbb{C}$  (see [4], [17], [18]). Let us define inductively, for  $r \in [0, 1)$ ,  $\exp_1 r = e^r$  and  $\exp_{p+1} r = \exp(\exp_p r)$ ,  $p \in \mathbb{N}$ . We also define for all  $r$  sufficiently large in  $(0, 1)$ ,  $\log_1 r = \log r$  and  $\log_{p+1} r = \log(\log_p r)$ ,  $p \in \mathbb{N}$ . Moreover, we denote by  $\exp_0 r = r$ ,  $\log_0 r = r$ ,  $\exp_{-1} r = \log_1 r$ ,  $\log_{-1} r = \exp_1 r$ .

**Definition 1.3** [5, 6] The iterated  $p$ -order of a meromorphic function  $f$  in  $\Delta$  is defined by

$$\rho_p(f) = \limsup_{r \rightarrow 1^-} \frac{\log_p^+ T(r, f)}{\log \frac{1}{1-r}} \quad (p \geq 1).$$

For an analytic function  $f$  in  $\Delta$ , we also define

$$\rho_{M,p}(f) = \limsup_{r \rightarrow 1^-} \frac{\log_{p+1}^+ M(r, f)}{\log \frac{1}{1-r}} \quad (p \geq 1).$$

**Remark 1.1** It follows by M. Tsuji in [25] that if  $f$  is an analytic function in  $\Delta$ , then

$$\rho_1(f) \leq \rho_{M,1}(f) \leq \rho_1(f) + 1.$$

However, it follows by Proposition 2.2.2 in [18]

$$\rho_{M,p}(f) = \rho_p(f), \quad (p \geq 2).$$

**Definition 1.4** [5] The growth index of the iterated order of a meromorphic function  $f(z)$  in  $\Delta$  is defined by

$$i(f) = \begin{cases} 0, & \text{if } f \text{ is non-admissible,} \\ \min \{j \in \mathbb{N}, \rho_j(f) < \infty\}, & \text{if } f \text{ is admissible,} \\ +\infty, & \text{if } \rho_j(f) = \infty \text{ for all } j \in \mathbb{N}. \end{cases}$$

For an analytic function  $f$  in  $\Delta$ , we also define

$$i_M(f) = \begin{cases} 0, & \text{if } f \text{ is non-admissible,} \\ \min \{j \in \mathbb{N}, \rho_{M,j}(f) < \infty\}, & \text{if } f \text{ is admissible,} \\ +\infty, & \text{if } \rho_{M,j}(f) = \infty \text{ for all } j \in \mathbb{N}. \end{cases}$$

**Definition 1.5** [3, 15, 20] The iterated  $p$ -type of a meromorphic function  $f$  of iterated  $p$ -order  $\rho$  ( $0 < \rho < \infty$ ) in  $\Delta$  is defined by

$$\tau_p(f) = \limsup_{r \rightarrow 1^-} (1-r)^{\rho_p(f)} \log_{p-1}^+ T(r, f).$$

**Definition 1.6** [7] Let  $f$  be a meromorphic function in  $\Delta$ . Then the iterated  $p$ -convergence exponent of the sequence of zeros of  $f(z)$  is defined by

$$\lambda_p(f) = \limsup_{r \rightarrow 1^-} \frac{\log_p^+ N\left(r, \frac{1}{f}\right)}{\log \frac{1}{1-r}},$$

where  $N\left(r, \frac{1}{f}\right)$  is the counting function of zeros of  $f(z)$  in  $\{z \in \mathbb{C} : |z| \leq r\}$ . Similarly, the iterated  $p$ -convergence exponent of the sequence of distinct zeros of  $f(z)$  is defined by

$$\bar{\lambda}_p(f) = \limsup_{r \rightarrow 1^-} \frac{\log_p^+ \bar{N}\left(r, \frac{1}{f}\right)}{\log \frac{1}{1-r}},$$

where  $\bar{N}\left(r, \frac{1}{f}\right)$  is the counting function of distinct zeros of  $f(z)$  in  $\{z \in \mathbb{C} : |z| \leq r\}$ .

**Definition 1.7** [7] The growth index of the convergence exponent of the sequence of the zeros of  $f(z)$  in  $\Delta$  is defined by

$$i_\lambda(f) = \begin{cases} 0, & \text{if } N\left(r, \frac{1}{f}\right) = O\left(\log \frac{1}{1-r}\right), \\ \min\{j \in \mathbb{N}, \lambda_j(f) < \infty\}, & \text{if some } j \in \mathbb{N} \text{ with } \lambda_j(f) < \infty, \\ +\infty, & \text{if } \lambda_j(f) = \infty \text{ for all } j \in \mathbb{N}. \end{cases}$$

Similarly, we can define the growth index of the convergence exponent of the sequence of distinct zeros  $i_{\bar{\lambda}}(f)$  of  $f(z)$  in  $\Delta$ .

Consider the complex differential equation

$$f^{(k)} + A_{k-1}(z) f^{(k-1)} + \dots + A_1(z) f' + A_0(z) f = 0 \tag{1.1}$$

and the  $k^{th}$  order differential polynomial

$$g_k = d_k f^{(k)} + d_{k-1} f^{(k-1)} + \dots + d_0 f, \tag{1.2}$$

where  $A_j$  ( $j = 0, 1, \dots, k-1$ ) and  $d_i$  ( $i = 0, 1, \dots, k$ ) are meromorphic functions in  $\Delta$ .

Let  $\mathcal{L}(\mathbf{G})$  denote a differential subfield of the field  $\mathcal{M}(\mathbf{G})$  of meromorphic functions in a domain  $\mathbf{G} \subset \mathbb{C}$ . If  $\mathbf{G} = \Delta$ , we simply denote  $\mathcal{L}$  instead of  $\mathcal{L}(\Delta)$ . Special case of such differential subfield

$$\mathcal{L}_{p+1, \rho} = \{g \text{ meromorphic: } \rho_{p+1}(g) < \rho\},$$

where  $\rho$  is a positive constant. In [7], T. B. Cao, H. Y. Xu and C. X. Zhu studied the complex oscillation of differential polynomial generated meromorphic solutions of second order linear differential equations with meromorphic coefficients and obtained the following results.

**Theorem A** [7] *Let  $A$  be an admissible meromorphic function of finite iterated order  $\rho_p(A) = \rho > 0$  ( $1 \leq p < \infty$ ) in the unit disc  $\Delta$  such that  $\delta(\infty, A) = \liminf_{r \rightarrow 1^-} \frac{m(r, A)}{T(r, A)} = \delta > 0$ , and let  $f$  be a non-zero meromorphic solution of the differential equation*

$$f'' + A(z)f = 0,$$

such that  $\delta(\infty, f) > 0$ . Moreover, let

$$P[f] = \sum_{j=0}^k p_j f^{(j)}$$

be a linear differential polynomial with coefficients  $p_j \in \mathcal{L}_{p+1, \rho}$ , assuming that at least one of the coefficients  $p_j$  does not vanish identically. If  $\varphi \in \mathcal{L}_{p+1, \rho}$  is a non-zero meromorphic function in  $\Delta$ , and neither  $P[f]$  nor  $P[f] - \varphi$  vanishes identically, then we have

$$i(f) = i_{\bar{\lambda}}(P[f] - \varphi) = p + 1$$

and

$$\bar{\lambda}_p(P[f] - \varphi) = \rho_{p+1}(f) = \rho_p(A) = \rho$$

if  $p > 1$ , while

$$\rho_p(A) \leq \bar{\lambda}_{p+1}(P[f] - \varphi) \leq \rho_{p+1}(f) \leq \rho_p(A) + 1$$

if  $p = 1$ .

**Remark 1.2** The idea of the proofs of Theorem A is borrowed from the paper of Laine, Rieppo [19] with the modifications reflecting the change from the complex plane  $\mathbb{C}$  to the unit disc  $\Delta$ .

Before we state our results, we define the sequence of meromorphic functions  $\alpha_{i,j}$  ( $j = 0, \dots, k-1$ ) in  $\Delta$  by

$$\alpha_{i,j} = \begin{cases} \alpha'_{i,j-1} + \alpha_{i-1,j-1} - A_i \alpha_{k-1,j-1}, & \text{for all } i = 1, \dots, k-1, \\ \alpha'_{0,j-1} - A_0 \alpha_{k-1,j-1}, & \text{for } i = 0 \end{cases} \quad (1.3)$$

and

$$\alpha_{i,0} = d_i - d_k A_i, \text{ for } i = 0, \dots, k-1 \quad (1.4)$$

we define also  $h_k$  by

$$h_k = \begin{vmatrix} \alpha_{0,0} & \alpha_{1,0} & \cdot & \cdot & \alpha_{k-1,0} \\ \alpha_{0,1} & \alpha_{1,1} & \cdot & \cdot & \alpha_{k-1,1} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \alpha_{0,k-1} & \alpha_{1,k-1} & \cdot & \cdot & \alpha_{k-1,k-1} \end{vmatrix}$$

and  $\psi_k(z)$  by

$$\psi_k(z) = C_0\varphi + C_1\varphi' + \cdots + C_{k-1}\varphi^{(k-1)},$$

where  $C_j$  ( $j = 0, \dots, k-1$ ) are finite iterated  $p$ -order meromorphic functions in  $\Delta$  depending on  $\alpha_{i,j}$ , and  $\varphi \not\equiv 0$  is a meromorphic function in  $\Delta$  with  $\rho_p(\varphi) < \infty$ .

The main purpose of this paper is to study the controllability of solutions of the differential equation (1.1). In the fact we study the growth and oscillation of higher order differential polynomial with meromorphic coefficients in the unit disc  $\Delta$  generated by solutions of equation (1.1).

**Theorem 1.1** *Let  $A_i(z)$  ( $i = 0, 1, \dots, k-1$ ) be meromorphic functions in  $\Delta$  of finite iterated  $p$ -order. Let  $d_j(z)$  ( $j = 0, 1, \dots, k$ ) be finite iterated  $p$ -order meromorphic functions in  $\Delta$  that are not all vanishing identically such that  $h \not\equiv 0$ . If  $f(z)$  is an infinite iterated  $p$ -order meromorphic solution in  $\Delta$  of (1.1) with  $\rho_{p+1}(f) = \rho$ , then the differential polynomial (1.2) satisfies*

$$\rho_p(g_k) = \rho_p(f) = \infty$$

and

$$\rho_{p+1}(g_k) = \rho_{p+1}(f) = \rho.$$

Furthermore, if  $f$  is a finite iterated  $p$ -order meromorphic solution in  $\Delta$  such that

$$\rho_p(f) > \max\{\rho_p(A_i) \ (i = 0, 1, \dots, k-1), \rho_p(d_j) \ (j = 0, 1, \dots, k)\}, \quad (1.5)$$

then

$$\rho_p(g_k) = \rho_p(f).$$

**Remark 1.3** In Theorem 1.1, if we do not have the condition  $h \not\equiv 0$ , then the conclusions of Theorem 1.1 cannot hold. For example, if we take  $d_i = d_k A_i$  ( $i = 0, \dots, k-1$ ), then  $h \equiv 0$ . It follows that  $g_k \equiv 0$  and  $\rho_p(g_k) = 0$ . So, if  $f(z)$  is an infinite iterated  $p$ -order meromorphic solution of (1.1), then  $\rho_p(g_k) = 0 < \rho_p(f) = \infty$ , and if  $f$  is a finite iterated  $p$ -order meromorphic solution of (1.1) such that (1.5) holds, then  $\rho_p(g_k) = 0 < \rho_p(f)$ .

**Theorem 1.2** *Under the hypotheses of Theorem 1.1, let  $\varphi(z) \not\equiv 0$  be a meromorphic function in  $\Delta$  with finite iterated  $p$ -order such that  $\psi_k(z)$  is not a solution of (1.1). If  $f(z)$  is an infinite iterated  $p$ -order meromorphic solution in  $\Delta$  of (1.1) with  $\rho_{p+1}(f) = \rho$ , then the differential polynomial (1.2) satisfies*

$$\bar{\lambda}_p(g_k - \varphi) = \lambda_p(g_k - \varphi) = \rho_p(f) = \infty$$

and

$$\bar{\lambda}_{p+1}(g_k - \varphi) = \lambda_{p+1}(g_k - \varphi) = \rho.$$

Furthermore, if  $f$  is a finite iterated  $p$ -order meromorphic solution in  $\Delta$  such that

$$\rho_p(f) > \max\{\rho_p(A_i) \ (i = 0, 1, \dots, k-1), \rho_p(d_j) \ (j = 0, 1, \dots, k), \rho_p(\varphi)\}, \quad (1.6)$$

then

$$\bar{\lambda}_p(g_k - \varphi) = \lambda_p(g_k - \varphi) = \rho_p(f).$$

From Theorems 1-2, we obtain the following corollaries which have been proved in [23].

**Corollary 1.1** [23] *Suppose that  $A(z)$  is admissible meromorphic function in  $\Delta$  such that  $i(A) = p$  ( $1 \leq p < \infty$ ) and  $\delta(\infty, A) = \delta > 0$ . Let  $d_j(z)$  ( $j = 0, 1, \dots, k$ ) be finite iterated  $p$ -order meromorphic functions in  $\Delta$  that are not all vanishing identically such that  $h \not\equiv 0$ , and let  $f$  be a nonzero meromorphic solution of*

$$f^{(k)} + A(z)f = 0. \quad (1.7)$$

*If  $\delta(\infty, f) > 0$ , then the differential polynomial  $g_k$  satisfies  $i(g_k) = p + 1$  and  $\rho_{p+1}(g_k) = \rho_{p+1}(f) = \rho_p(A)$  if  $p > 1$ , while*

$$\rho_p(A) \leq \rho_{p+1}(g_k) = \rho_{p+1}(f) \leq \rho_p(A) + 1$$

*if  $p = 1$ .*

**Corollary 1.2** [23] *Under the hypotheses of Corollary 1.1, let  $\varphi(z) \not\equiv 0$  be meromorphic function in  $\Delta$  with finite iterated  $p$ -order such that  $\psi_k(z) \not\equiv 0$ . Then the differential polynomial (1.2) satisfies*

$$\bar{\lambda}_{p+1}(g_k - \varphi) = \lambda_{p+1}(g_k - \varphi) = \rho_{p+1}(f) = \rho_p(A)$$

*if  $p > 1$ , while*

$$\rho_p(A) \leq \bar{\lambda}_{p+1}(g_k - \varphi) = \lambda_{p+1}(g_k - \varphi) = \rho_{p+1}(f) \leq \rho_p(A) + 1$$

*if  $p = 1$ .*

**Remark 1.4** The present article may be understood as an extension and improvement of the recent article of the authors [23] from equation (1.7) to equation (1.1). The method used in the proofs of our theorems is simple and quite different from the method used in the papers of Laine and Rieppo [19] and Cao, Xu and Zhu [7].

We consider now the differential equation

$$f'' + A_1(z)f' + A_0(z)f = 0, \quad (1.8)$$

where  $A_1(z), A_0(z)$  are analytic functions of finite iterated  $p$ -order in the unit disc  $\Delta$ . In the following we will give sufficient conditions on  $A_1$  and  $A_0$  which satisfied the results of Theorem 1.1 and Theorem 1.2 without the conditions "  $h_k \not\equiv 0$  " and "  $\psi_k(z)$  is not a solution of (1.1) " where  $k = 2$ .

**Corollary 1.3** *Let  $A_1(z), A_0(z) (\not\equiv 0)$  be analytic functions in  $\Delta$  such that  $\rho_p(A_0) = \rho$  ( $0 < \rho < \infty$ ),  $\tau_p(A_0) = \tau$  ( $0 < \tau < \infty$ ), and let  $\rho_p(A_1) < \rho_p(A_0)$  or  $\tau_p(A_1) <$*

$\tau_p(A_0)$  if  $\rho_p(A_0) = \rho_p(A_1)$ . Let  $d_2, d_1, d_0$  be analytic functions in  $\Delta$  such that at least one of  $d_2, d_1, d_0$  does not vanish identically with  $\max\{\rho_p(d_j) \ (j = 0, 1, 2)\} < \rho_p(A_0)$ . If  $f \not\equiv 0$  is a solution of (1.8), then the differential polynomial  $g_2 = d_2f'' + d_1f' + d_0f$  satisfies  $\rho_{p+1}(g_2) = \rho_{p+1}(f) = \rho_p(A_0)$  if  $p > 1$ , while

$$\rho_p(A_0) \leq \rho_{p+1}(g_2) = \rho_{p+1}(f) \leq \max\{\rho_M(A_j) \ (j = 0, 1)\}$$

if  $p = 1$ .

**Corollary 1.4** Let  $A_1(z), A_0(z) (\not\equiv 0)$  be analytic functions in  $\Delta$  such that  $\rho_p(A_0) = \rho$  ( $0 < \rho < \infty$ ),  $\tau_p(A_0) = \tau$  ( $0 < \tau < \infty$ ), and let  $\rho_p(A_1) < \rho_p(A_0)$  or  $2\tau_p(A_1) < \tau_p(A_0)$  if  $\rho_p(A_0) = \rho_p(A_1)$ . Let  $d_2, d_1, d_0$  be analytic functions in  $\Delta$  such that at least one of  $d_2, d_1, d_0$  does not vanish identically with  $\max\{\rho_p(d_j) \ (j = 0, 1, 2)\} < \rho_p(A_1)$ , and let  $\varphi(z) \not\equiv 0$  be analytic function in  $\Delta$  of finite iterated  $p$ -order such that  $\psi_2(z) \not\equiv 0$ . If  $f \not\equiv 0$  is a solution of (1.8), then the differential polynomial  $g_2 = d_2f'' + d_1f' + d_0f$  satisfies

$$\bar{\lambda}_{p+1}(g_2 - \varphi) = \lambda_{p+1}(g_2 - \varphi) = \rho_{p+1}(f) = \rho_p(A_0)$$

if  $p > 1$ , while

$$\rho_p(A_0) \leq \bar{\lambda}_{p+1}(g_2 - \varphi) = \lambda_{p+1}(g_2 - \varphi) = \rho_{p+1}(f) \leq \max\{\rho_M(A_j) \ (j = 0, 1)\}$$

if  $p = 1$ .

**Remark 1.5** For some papers related in the complex plane see [19, 22, 24] and in the unit disc see [7, 9, 12].

## 2 Auxiliary lemmas

**Lemma 2.1** [8] Let  $A_0, A_1, \dots, A_{k-1}, F \not\equiv 0$  be meromorphic functions in  $\Delta$ , and let  $f$  be a meromorphic solution of the differential equation

$$f^{(k)} + A_{k-1}(z)f^{(k-1)} + \dots + A_1(z)f' + A_0(z)f = F(z) \quad (2.1)$$

such that  $i(f) = p$  ( $1 \leq p < \infty$ ). If either

$$\max\{i(A_j) \ (j = 0, 1, \dots, k-1), i(F)\} < p$$

or

$$\max\{\rho_p(A_j) \ (j = 0, 1, \dots, k-1), \rho_p(F)\} < \rho_p(f),$$

then

$$i_{\bar{\lambda}}(f) = i_{\lambda}(f) = i(f) = p$$

and

$$\bar{\lambda}_p(f) = \lambda_p(f) = \rho_p(f).$$

Using the same arguments as in the proof of Lemma 2.1 (see, the proof of Lemma 2.5 in [8]), we easily obtain the following lemma.

**Lemma 2.2** *Let  $A_0, A_1, \dots, A_{k-1}, F \not\equiv 0$  be finite iterated  $p$ -order meromorphic functions in the unit disc  $\Delta$ . If  $f$  is a meromorphic solution with  $\rho_p(f) = \infty$  and  $\rho_{p+1}(f) = \rho < \infty$  of equation (2.1), then  $\bar{\lambda}_p(f) = \lambda_p(f) = \rho_p(f) = \infty$  and  $\bar{\lambda}_{p+1}(f) = \lambda_{p+1}(f) = \rho_{p+1}(f) = \rho$ .*

**Lemma 2.3** [5] *Let  $p \geq 1$  be an integer, and let  $A_0(z), \dots, A_{k-1}(z)$  be analytic functions in  $\Delta$  such that  $i(A_0) = p$ . If*

$$\max\{i(A_j) : j = 1, \dots, k-1\} < p$$

or

$$\max\{\rho_p(A_j) : j = 1, \dots, k-1\} < \rho_p(A_0),$$

then every solution  $f \not\equiv 0$  of equation (1.1) satisfies  $i(f) = p+1$  and  $\rho_p(f) = \infty$ ,  $\rho_p(A_0) \leq \rho_{p+1}(f) = \rho_{M,p+1}(f) \leq \max\{\rho_{M,p}(A_j) : j = 0, 1, \dots, k-1\}$ .

**Lemma 2.4** [3] *Let  $f$  and  $g$  be meromorphic functions in the unit disc  $\Delta$  such that  $0 < \rho_p(f), \rho_p(g) < \infty$  and  $0 < \tau_p(f), \tau_p(g) < \infty$ . Then we have*

(i) *If  $\rho_p(f) > \rho_p(g)$ , then we obtain*

$$\tau_p(f+g) = \tau_p(fg) = \tau_p(f).$$

(ii) *If  $\rho_p(f) = \rho_p(g)$  and  $\tau_p(f) \neq \tau_p(g)$ , then we get*

$$\rho_p(f+g) = \rho_p(fg) = \rho_p(f) = \rho_p(g).$$

**Lemma 2.5** ([14]) *Let  $f$  be a meromorphic function in the unit disc and let  $k \in \mathbb{N}$ . Then*

$$m\left(r, \frac{f^{(k)}}{f}\right) = S(r, f),$$

where  $S(r, f) = O\left(\log^+ T(r, f) + \log\left(\frac{1}{1-r}\right)\right)$ , possibly outside a set  $E_1 \subset [0, 1)$  with  $\int_{E_1} \frac{dr}{1-r} < \infty$ . If  $f$  is of finite order of growth, then

$$m\left(r, \frac{f^{(k)}}{f}\right) = O\left(\log\left(\frac{1}{1-r}\right)\right).$$

**Lemma 2.6** [2] *Let  $f$  be a meromorphic function in the unit disc for which  $i(f) = p \geq 1$  and  $\rho_p(f) = \beta < \infty$ , and let  $k \in \mathbb{N}$ . Then for any  $\varepsilon > 0$ ,*

$$m\left(r, \frac{f^{(k)}}{f}\right) = O\left(\exp_{p-2}\left(\log\frac{1}{1-r}\right)^{\beta+\varepsilon}\right)$$

for all  $r$  outside a set  $E_2 \subset [0, 1)$  with  $\int_{E_2} \frac{dr}{1-r} < \infty$ .



**Lemma 2.7** *Let  $f$  be a meromorphic function in  $\Delta$  with iterated order  $\rho_p(f) = \rho$  ( $0 < \rho < \infty$ ) and iterated type  $\tau_p(f) = \tau$  ( $0 < \tau < \infty$ ). Then for any given  $\beta < \tau$ , there exists a subset  $E_3$  of  $[0, 1)$  that has an infinite logarithmic measure such that  $\log_{p-1} T(r, f) > \beta \left(\frac{1}{1-r}\right)^\rho$  holds for all  $r \in E_3$ .*

*Proof.* When  $p = 1$ , the lemma is proved in [21]. Thus we assume  $p \geq 2$ . By definitions of iterated order and iterated type, there exists an increasing sequence  $\{r_m\}_{m=1}^\infty \subset [0, 1)$  ( $r_m \rightarrow 1^-$ ) satisfying  $\frac{1}{m} + (1 - \frac{1}{m})r_m < r_{m+1}$  and

$$\lim_{m \rightarrow \infty} \frac{\log_{p-1} T(r_m, f)}{\left(\frac{1}{1-r_m}\right)^\rho} = \tau_p(f).$$

Then there exists a positive integer  $m_0$  such that for all  $m > m_0$  and for any given  $0 < \varepsilon < \tau_p(f) - \beta$ , we have

$$\log_{p-1} T(r_m, f) > (\tau_p(f) - \beta) \left(\frac{1}{1-r_m}\right)^\rho. \tag{2.2}$$

For any given  $\beta < \tau_p(f) - \varepsilon$ , there exists a positive integer  $m_1$  such that for all  $m > m_1$  we have

$$\left(1 - \frac{1}{m}\right)^\rho > \frac{\beta}{\tau_p(f) - \varepsilon}. \tag{2.3}$$

Take  $m \geq m_2 = \max\{m_0, m_1\}$ . By (2.2) and (2.3), for any  $r \in [r_m, \frac{1}{m} + (1 - \frac{1}{m})r_m]$ , we have

$$\begin{aligned} \log_{p-1} T(r, f) &\geq \log_{p-1} T(r_m, f) > (\tau_p(f) - \beta) \left(\frac{1}{1-r_m}\right)^\rho \\ &\geq (\tau_p(f) - \beta) \left(1 - \frac{1}{m}\right)^\rho \left(\frac{1}{1-r}\right)^\rho > \beta \left(\frac{1}{1-r}\right)^\rho. \end{aligned}$$

Set  $E_3 = \cup_{m=m_2}^\infty [r_m, \frac{1}{m} + (1 - \frac{1}{m})r_m]$ , then there holds

$$m_l E_3 = \sum_{m=m_2}^\infty \int_{r_m}^{\frac{1}{m} + (1 - \frac{1}{m})r_m} \frac{dt}{1-t} = \sum_{m=m_2}^\infty \log \frac{m}{m-1} = \infty.$$

**Lemma 2.8** [16] *Let  $f$  be a solution of equation (1.1) where the coefficients  $A_j(z)$  ( $j = 0, \dots, k-1$ ) are analytic functions in the disc  $\Delta_R = \{z \in \mathbb{C} : |z| < R\}$ ,  $0 < R \leq \infty$ . Let  $n_c \in \{1, \dots, k\}$  be the number of nonzero coefficients  $A_j(z)$  ( $j = 0, \dots, k-1$ ), and let  $\theta \in [0, 2\pi[$  and  $\varepsilon > 0$ . If  $z_\theta = \nu e^{i\theta} \in \Delta_R$  is such that  $A_j(z_\theta) \neq 0$  for some  $j = 0, \dots, k-1$ , then for all  $\nu < r < R$ ,*

$$|f(re^{i\theta})| \leq C \exp \left( n_c \int_\nu^r \max_{j=0, \dots, k-1} |A_j(te^{i\theta})|^{\frac{1}{k-j}} dt \right), \tag{2.4}$$

where  $C > 0$  is a constant satisfying

$$C \leq (1 + \varepsilon) \max_{j=0, \dots, k-1} \left( \frac{|f^{(j)}(z_\theta)|}{(n_c)^j \max_{n=0, \dots, k-1} |A_n(z_\theta)|^{\frac{j}{k-n}}} \right).$$

**Lemma 2.9** [1, 14] *Let  $g : (0, 1) \rightarrow \mathbb{R}$  and  $h : (0, 1) \rightarrow \mathbb{R}$  be monotone increasing functions such that  $g(r) \leq h(r)$  holds outside of an exceptional set  $E_4 \subset [0, 1)$  for which  $\int_{E_4} \frac{dr}{1-r} < \infty$ . Then there exists a constant  $d \in (0, 1)$  such that if  $s(r) = 1 - d(1-r)$ , then  $g(r) \leq h(s(r))$  for all  $r \in [0, 1)$ .*

**Lemma 2.10** *Let  $A_1(z)$  and  $A_0(z)$  be analytic functions in  $\Delta$  such that  $\rho_p(A_0) = \rho$  ( $0 < \rho < \infty$ ),  $\tau_p(A_0) = \tau$  ( $0 < \tau < \infty$ ), and let  $\rho_p(A_1) < \rho_p(A_0)$  and  $\tau_p(A_1) < \tau_p(A_0)$  if  $\rho_p(A_1) = \rho_p(A_0)$ . If  $f \neq 0$  is a solution of (1.8) then  $\rho_p(f) = \infty$ ,  $\rho_{p+1}(f) = \rho_p(A_0)$  if  $p > 1$ , while*

$$\rho_p(f) = \infty, \quad \rho_p(A_0) \leq \rho_{p+1}(f) \leq \max\{\rho_M(A_j), (j = 0, 1)\}$$

if  $p = 1$ .

*Proof.* If  $\rho_p(A_1) < \rho_p(A_0)$  then the result can easily deduced by Lemma 2.3. We prove only the case when  $\rho_p(A_0) = \rho_p(A_1) = \rho$  and  $\tau_p(A_1) < \tau_p(A_0)$ . Since  $f \neq 0$ , then by (1.8) we have

$$A_0 = - \left( \frac{f''}{f} + A_1 \frac{f'}{f} \right). \quad (2.5)$$

Suppose that  $f$  is of finite  $p$ -iterated order, then by Lemma 2.6

$$T(r, A_0) \leq T(r, A_1) + O \left( \exp_{p-2} \left( \log \frac{1}{1-r} \right)^{\beta+\varepsilon} \right) \quad (\rho_p(f) = \beta < \infty) \quad (2.6)$$

which implies the contradiction

$$\tau_p(A_0) \leq \tau_p(A_1).$$

Hence  $\rho_p(f) = \infty$ . By using inequality (2.4), we have

$$\rho_{p+1}(f) \leq \max\{\rho_p(A_1), \rho_p(A_0)\} = \rho_p(A_0). \quad (2.7)$$

On the other hand, by Lemma 2.5

$$T(r, A_0) \leq T(r, A_1) + O \left( \log^+ T(r, f) + \log \frac{1}{1-r} \right) \quad (2.8)$$

holds possibly outside a set  $E_1 \subset [0, 1)$  with  $\int_{E_1} \frac{dr}{1-r} < \infty$ . By  $\tau_p(A_1) < \tau_p(A_0)$  we choose  $\alpha_0, \alpha_1$  satisfying  $\tau_p(A_1) < \alpha_1 < \alpha_0 < \tau_p(A_0)$  such that for  $r \rightarrow 1^-$ , we have

$$T(r, A_1) \leq \exp_{p-1} \left\{ \alpha_1 \left( \frac{1}{1-r} \right)^\rho \right\}. \quad (2.9)$$

By Lemma 2.7, there exists a subset  $E_2 \subset [0, 1)$  of infinite logarithmic measure such that

$$T(r, A_0) > \exp_{p-1} \left\{ \alpha_0 \left( \frac{1}{1-r} \right)^\rho \right\}. \quad (2.10)$$

By (2.8) – (2.10) we obtain for all  $r \in E_2 - E_1$

$$(1 - o(1)) \exp_{p-1} \left\{ \alpha_0 \left( \frac{1}{1-r} \right)^\rho \right\} \leq O \left( \log^+ T(r, f) + \log \frac{1}{1-r} \right). \quad (2.11)$$

By using (2.11) and Lemma 2.9, we obtain

$$\rho_{p+1}(f) \geq \rho_p(A_0). \quad (2.12)$$

From (2.7) and (2.12) we get  $\rho_p(f) = \infty$  and  $\rho_{p+1}(f) = \rho_p(A_0)$ .

### 3 Proof of the Theorems and the Corollaries

**Proof of Theorem 1.1** Suppose that  $f$  is an infinite iterated  $p$ -order meromorphic solution in  $\Delta$  of (1.1). By (1.1), we have

$$f^{(k)} = - \sum_{i=0}^{k-1} A_i f^{(i)} \quad (3.1)$$

which implies

$$\begin{aligned} g_k &= d_k f^{(k)} + d_{k-1} f^{(k-1)} + \dots + d_0 f \\ &= \sum_{i=0}^{k-1} (d_i - d_k A_i) f^{(i)}. \end{aligned} \quad (3.2)$$

We can write (3.2) as

$$g_k = \sum_{i=0}^{k-1} \alpha_{i,0} f^{(i)}, \quad (3.3)$$

where  $\alpha_{i,0}$  are defined in (1.4). Differentiating both sides of equation (3.3) and replacing  $f^{(k)}$  with  $f^{(k)} = - \sum_{i=0}^{k-1} A_i f^{(i)}$ , we obtain

$$\begin{aligned} g'_k &= \sum_{i=0}^{k-1} \alpha'_{i,0} f^{(i)} + \sum_{i=0}^{k-1} \alpha_{i,0} f^{(i+1)} = \sum_{i=0}^{k-1} \alpha'_{i,0} f^{(i)} + \sum_{i=1}^k \alpha_{i-1,0} f^{(i)} \\ &= \alpha'_{0,0} f + \sum_{i=1}^{k-1} \alpha'_{i,0} f^{(i)} + \sum_{i=1}^{k-1} \alpha_{i-1,0} f^{(i)} + \alpha_{k-1,0} f^{(k)} \\ &= \alpha'_{0,0} f + \sum_{i=1}^{k-1} \alpha'_{i,0} f^{(i)} + \sum_{i=1}^{k-1} \alpha_{i-1,0} f^{(i)} - \sum_{i=0}^{k-1} \alpha_{k-1,0} A_i f^{(i)} \end{aligned}$$

$$= (\alpha'_{0,0} - \alpha_{k-1,0}A_0) f + \sum_{i=1}^{k-1} (\alpha'_{i,0} + \alpha_{i-1,0} - \alpha_{k-1,0}A_i) f^{(i)}. \quad (3.4)$$

We can rewrite (3.4) as

$$g'_k = \sum_{i=0}^{k-1} \alpha_{i,1} f^{(i)}, \quad (3.5)$$

where

$$\alpha_{i,1} = \begin{cases} \alpha'_{i,0} + \alpha_{i-1,0} - \alpha_{k-1,0}A_i, & \text{for all } i = 1, \dots, k-1, \\ \alpha'_{0,0} - A_0\alpha_{k-1,0}, & \text{for } i = 0. \end{cases} \quad (3.6)$$

Differentiating both sides of equation (3.5) and replacing  $f^{(k)}$  with  $f^{(k)} = -\sum_{i=0}^{k-1} A_i f^{(i)}$ , we obtain

$$\begin{aligned} g''_k &= \sum_{i=0}^{k-1} \alpha'_{i,1} f^{(i)} + \sum_{i=0}^{k-1} \alpha_{i,1} f^{(i+1)} = \sum_{i=0}^{k-1} \alpha'_{i,1} f^{(i)} + \sum_{i=1}^k \alpha_{i-1,1} f^{(i)} \\ &= \alpha'_{0,1} f + \sum_{i=1}^{k-1} \alpha'_{i,1} f^{(i)} + \sum_{i=1}^{k-1} \alpha_{i-1,1} f^{(i)} + \alpha_{k-1,1} f^{(k)} \\ &= \alpha'_{0,1} f + \sum_{i=1}^{k-1} \alpha'_{i,1} f^{(i)} + \sum_{i=1}^{k-1} \alpha_{i-1,1} f^{(i)} - \sum_{i=0}^{k-1} A_i \alpha_{k-1,1} f^{(i)} \\ &= (\alpha'_{0,1} - \alpha_{k-1,1}A_0) f + \sum_{i=1}^{k-1} (\alpha'_{i,1} + \alpha_{i-1,1} - A_i \alpha_{k-1,1}) f^{(i)} \end{aligned} \quad (3.7)$$

which implies that

$$g''_k = \sum_{i=0}^{k-1} \alpha_{i,2} f^{(i)}, \quad (3.8)$$

where

$$\alpha_{i,2} = \begin{cases} \alpha'_{i,1} + \alpha_{i-1,1} - A_i \alpha_{k-1,1}, & \text{for all } i = 1, \dots, k-1, \\ \alpha'_{0,1} - A_0 \alpha_{k-1,1}, & \text{for } i = 0. \end{cases} \quad (3.9)$$

By using the same method as above we can easily deduce that

$$g_f^{(j)} = \sum_{i=0}^{k-1} \alpha_{i,j} f^{(i)}, \quad j = 0, 1, \dots, k-1, \quad (3.10)$$

where

$$\alpha_{i,j} = \begin{cases} \alpha'_{i,j-1} + \alpha_{i-1,j-1} - A_i \alpha_{k-1,j-1}, & \text{for all } i = 1, \dots, k-1, \\ \alpha'_{0,j-1} - A_0 \alpha_{k-1,j-1}, & \text{for } i = 0 \end{cases} \quad (3.11)$$

and

$$\alpha_{i,0} = d_i - d_k A_i, \text{ for all } i = 0, 1, \dots, k - 1. \quad (3.12)$$

By (3.3) – (3.12) we obtain the system of equations

$$\begin{cases} g_k = \alpha_{0,0}f + \alpha_{1,0}f' + \dots + \alpha_{k-1,0}f^{(k-1)}, \\ g'_k = \alpha_{0,1}f + \alpha_{1,1}f' + \dots + \alpha_{k-1,1}f^{(k-1)}, \\ g''_k = \alpha_{0,2}f + \alpha_{1,2}f' + \dots + \alpha_{k-1,2}f^{(k-1)}, \\ \dots \\ g_k^{(k-1)} = \alpha_{0,k-1}f + \alpha_{1,k-1}f' + \dots + \alpha_{k-1,k-1}f^{(k-1)}. \end{cases} \quad (3.13)$$

By Cramer's rule, and since  $h_k \neq 0$  we have

$$f = \frac{\begin{vmatrix} g_k & \alpha_{1,0} & \dots & \alpha_{k-1,0} \\ g'_k & \alpha_{1,1} & \dots & \alpha_{k-1,1} \\ \vdots & \vdots & \ddots & \vdots \\ g_k^{(k-1)} & \alpha_{1,k-1} & \dots & \alpha_{k-1,k-1} \end{vmatrix}}{h}. \quad (3.14)$$

Then

$$f = C_0 g_k + C_1 g'_k + \dots + C_{k-1} g_k^{(k-1)}, \quad (3.15)$$

where  $C_j$  are finite iterated  $p$ -order meromorphic functions in  $\Delta$  depending on  $\alpha_{i,j}$ , where  $\alpha_{i,j}$  is defined in (3.11).

If  $\rho_p(g_k) < +\infty$ , then by (3.15) we obtain  $\rho_p(f) < +\infty$ , and this is a contradiction. Hence  $\rho_p(g_k) = \rho_p(f) = +\infty$ .

Now, we prove that  $\rho_{p+1}(g_k) = \rho_{p+1}(f) = \rho$ . By (3.2), we get  $\rho_{p+1}(g_k) \leq \rho_{p+1}(f)$  and by (3.15) we have  $\rho_{p+1}(f) \leq \rho_{p+1}(g_k)$ . This yield  $\rho_{p+1}(g_k) = \rho_{p+1}(f) = \rho$ .

Furthermore, if  $f$  is a finite iterated  $p$ -order meromorphic solution in  $\Delta$  of equation (1.1) such that

$$\rho_p(f) > \max \{ \rho_p(A_i) \ (i = 0, \dots, k - 1), \rho_p(d_j) \ (j = 0, 1, \dots, k) \}, \quad (3.16)$$

then

$$\rho_p(f) > \max \{ \rho_p(\alpha_{i,j}) : i = 0, \dots, k - 1, j = 0, \dots, k - 1 \}. \quad (3.17)$$

By (3.2) and (3.16) we have  $\rho_p(g_k) \leq \rho_p(f)$ . Now, we prove  $\rho_p(g_k) = \rho_p(f)$ . If  $\rho_p(g_k) < \rho_p(f)$ , then by (3.15) and (3.17) we get

$$\rho_p(f) \leq \max \{ \rho_p(C_j) \ (j = 0, \dots, k - 1), \rho_p(g_k) \} < \rho_p(f)$$

and this is a contradiction. Hence  $\rho_p(g_k) = \rho_p(f)$ .

**Remark 3.1** From (3.13), it follows that the condition  $h \neq 0$  is equivalent to the condition  $g_k, g'_k, \dots, g_k^{(k-1)}$  are linearly independent over the field of meromorphic functions of finite iterated  $p$ -order.

**Proof of Theorem 1.2** Suppose that  $f$  is an infinite iterated  $p$ -order meromorphic solution in  $\Delta$  of equation (1.1) with  $\rho_{p+1}(f) = \rho$ . Set  $w(z) = g_k - \varphi$ . Since  $\rho_p(\varphi) < \infty$ , then by Theorem 1.1 we have  $\rho_p(w) = \rho_p(g_k) = \infty$  and  $\rho_{p+1}(w) = \rho_{p+1}(g_k) = \rho$ . To prove  $\bar{\lambda}_p(g_k - \varphi) = \lambda_p(g_k - \varphi) = \infty$  and  $\bar{\lambda}_{p+1}(g_k - \varphi) = \lambda_{p+1}(g_k - \varphi) = \rho$  we need to prove  $\bar{\lambda}_p(w) = \lambda_p(w) = \infty$  and  $\bar{\lambda}_{p+1}(w) = \lambda_{p+1}(w) = \rho$ . By  $g_k = w + \varphi$ , and using (3.15), we get

$$f = C_0 w + C_1 w' + \cdots + C_{k-1} w^{(k-1)} + \psi_k(z), \quad (3.18)$$

where

$$\psi_k(z) = C_0 \varphi + C_1 \varphi' + \cdots + C_{k-1} \varphi^{(k-1)}. \quad (3.19)$$

Substituting (3.18) into (1.1), we obtain

$$C_{k-1} w^{(2k-1)} + \sum_{j=0}^{2k-2} \phi_j w^{(j)} = - \left( \psi_k^{(k)} + A_{k-1}(z) \psi_k^{(k-1)} + \cdots + A(z) \psi_k \right) = H, \quad (3.20)$$

where  $C_{k-1}, \phi_j$  ( $j = 0, \dots, 2k-1$ ) are meromorphic functions in  $\Delta$  with finite iterated  $p$ -order. Since  $\psi_k(z)$  is not a solution of (1.1), it follows that  $H \not\equiv 0$ . Then by Lemma 2.2, we obtain  $\bar{\lambda}_p(w) = \lambda_p(w) = \infty$  and  $\bar{\lambda}_{p+1}(w) = \lambda_{p+1}(w) = \rho$ , i. e.,  $\bar{\lambda}_p(g_k - \varphi) = \lambda_p(g_k - \varphi) = \infty$  and  $\bar{\lambda}_{p+1}(g_k - \varphi) = \lambda_{p+1}(g_k - \varphi) = \rho$ .

Suppose that  $f$  is a finite iterated  $p$ -order meromorphic solution in  $\Delta$  of equation (1.1) such that (1.6) holds. Set  $w(z) = g_k - \varphi$ . Since  $\rho_p(\varphi) < \rho_p(f)$ , then by Theorem 1.1 we have  $\rho_p(w) = \rho_p(g_k) = \rho_p(f)$ . To prove  $\bar{\lambda}_p(g_k - \varphi) = \lambda_p(g_k - \varphi) = \rho_p(f)$  we need to prove  $\bar{\lambda}_p(w) = \lambda_p(w) = \rho_p(f)$ . Using the same reasoning as above, we get

$$C_{k-1} w^{(2k-1)} + \sum_{j=0}^{2k-2} \phi_j w^{(j)} = - \left( \psi_k^{(k)} + A_{k-1}(z) \psi_k^{(k-1)} + \cdots + A(z) \psi_k \right) = F,$$

where  $C_{k-1}, \phi_j$  ( $j = 0, \dots, 2k-1$ ) are meromorphic functions in  $\Delta$  with finite iterated  $p$ -order  $\rho_p(C_{k-1}) < \rho_p(f)$ ,  $\rho_p(\phi_j) < \rho_p(f)$  ( $j = 0, \dots, 2k-1$ ), and

$$\psi_k(z) = C_0 \varphi + C_1 \varphi' + \cdots + C_{k-1} \varphi^{(k-1)}, \quad \rho_p(F) < \rho_p(f).$$

Since  $\psi_k(z)$  is not a solution of (1.1), it follows that  $F \not\equiv 0$ . Then by Lemma 2.1, we obtain  $\bar{\lambda}_p(w) = \lambda_p(w) = \rho_p(f)$ , i. e.,  $\bar{\lambda}_p(g_k - \varphi) = \lambda_p(g_k - \varphi) = \rho_p(f)$ .

**Proof of Corollary 1.3** Suppose that  $f$  is a nontrivial solution of (1.8). Then by Lemma 2.3, we have

$$\rho_p(A_0) \leq \rho_{p+1}(f) \leq \max \{ \rho_{M,p}(A_j) \quad (j = 0, 1) \} \quad (p \geq 1).$$

By the same reasoning as before we obtain that

$$\begin{cases} g_2 = \alpha_{0,0} f + \alpha_{1,0} f', \\ g'_2 = \alpha_{0,1} f + \alpha_{1,1} f', \end{cases}$$

where

$$\alpha_{0,0} = d_0 - d_2 A_0, \quad \alpha_{1,1} = d_2 A_1^2 - (d_2 A_1)' - d_1 A_1 - d_2 A_0 + d_0 + d_1'$$

and

$$\alpha_{0,1} = d_2 A_0 A_1 - (d_2 A_0)' - d_1 A_0 + d_0', \quad \alpha_{1,0} = d_1 - d_2 A_1.$$

First, we suppose that  $d_2 \neq 0$ . We have

$$\begin{aligned} h_2 &= \begin{vmatrix} \alpha_{1,0} & \alpha_{0,0} \\ \alpha_{1,1} & \alpha_{0,1} \end{vmatrix} \\ &= -d_2^2 A_0^2 - d_0 d_2 A_1^2 + (-d_2 d_1 + d_1' d_2 + 2d_0 d_2 - d_1^2) A_0 \\ &\quad + (d_1' d_0 - d_2 d_0' + d_0 d_1) A_1 + d_1 d_2 A_0 A_1 - d_1 d_2 A_0' + d_0 d_2 A_1' \\ &\quad + d_2^2 A_0' A_1 - d_2^2 A_0 A_1' + d_0' d_1 - d_0 d_1' - d_0^2. \end{aligned}$$

By  $d_2 \neq 0$ ,  $A_0 \neq 0$  and Lemma 2.4, we have  $\rho_p(h) = \rho_p(A_0)$ . Hence  $h \neq 0$ . Now suppose  $d_2 \equiv 0$ ,  $d_1 \neq 0$  or  $d_2 \equiv 0$ ,  $d_1 \equiv 0$  and  $d_0 \neq 0$ , then by using a similar reasoning as above we get  $h \neq 0$ , and we obtain

$$f = \frac{\alpha_{1,0} g_f' - \alpha_{1,1} g_f}{h_2}. \quad (3.21)$$

It is clear that  $\rho_p(g_2) \leq \rho_p(f)$  ( $\rho_{p+1}(g_2) \leq \rho_{p+1}(f)$ ) and by (3.21) we have  $\rho_p(f) \leq \rho_p(g_2)$  ( $\rho_{p+1}(f) \leq \rho_{p+1}(g_2)$ ). Hence  $\rho_p(g_2) = \rho_p(f)$  ( $\rho_{p+1}(g_2) = \rho_{p+1}(f)$ ).

**Proof of Corollary 1.4** Set  $w(z) = d_2 f'' + d_1 f' + d_0 f - \varphi$ . Then, by  $\rho_p(\varphi) < \infty$ , we have  $\rho_p(w) = \rho_p(g_2) = \rho_p(f)$  and  $\rho_{p+1}(w) = \rho_{p+1}(g_2) = \rho_{p+1}(f)$ . In order to prove  $\bar{\lambda}_{p+1}(g_2 - \varphi) = \lambda_{p+1}(g_2 - \varphi) = \rho_{p+1}(f)$ , we need to prove only  $\bar{\lambda}_{p+1}(w) = \lambda_{p+1}(w) = \rho_{p+1}(f)$ . Using  $g_2 = w + \varphi$ , we get from (3.21)

$$f = \frac{\alpha_{1,0} w' - \alpha_{1,1} w}{h_2} + \psi_2, \quad (3.22)$$

where

$$\psi_2(z) = \frac{\alpha_{1,0} \varphi' - \alpha_{1,1} \varphi}{h_2}. \quad (3.23)$$

Substituting (3.22) into equation (1.8), we obtain

$$\begin{aligned} &\frac{\alpha_{1,0}}{h_2} w''' + \phi_2 w'' + \phi_1 w' + \phi_0 w \\ &= -\left(\psi_2'' + A_1(z) \psi_2' + A_0(z) \psi_2\right) = A, \end{aligned} \quad (3.24)$$

where  $\phi_j$  ( $j = 0, 1, 2$ ) are meromorphic functions in  $\Delta$  with  $\rho_p(\phi_j) < \infty$  ( $j = 0, 1, 2$ ). First, we prove that  $\psi_2 \neq 0$ . Suppose that  $\psi_2 \equiv 0$ , then by (3.23) we obtain

$$\alpha_{1,1} = \alpha_{1,0} \frac{\varphi'}{\varphi}. \quad (3.25)$$

It follows that by using Lemma 2.6

$$m(r, \alpha_{1,1}) \leq m(r, \alpha_{1,0}) + O\left(\exp_{p-2}\left(\log\frac{1}{1-r}\right)^{\beta+\varepsilon}\right), \quad \rho_p(\varphi) = \beta < \infty. \quad (3.26)$$

(i) If  $d_2 \not\equiv 0$ , then by using Lemma 2.4 we obtain the contradiction

$$\begin{cases} \rho_p(A_0) \leq \rho_p(A_1), & \text{if } \rho_p(A_0) > \rho_p(A_1), \\ \tau_p(A_0) \leq \tau_p(A_1), & \text{if } \rho_p(A_0) = \rho_p(A_1). \end{cases}$$

(ii) If  $d_2 \equiv 0$  and  $d_1 \not\equiv 0$ , we obtain the contradiction

$$\rho_p(A_1) \leq \rho_p(d_1).$$

(iii) If  $d_2 = d_1 \equiv 0$  and  $d_0 \not\equiv 0$ , we have by (3.25)

$$d_0 = \alpha_{1,1} = \alpha_{1,0} \frac{\varphi'}{\varphi} = 0 \times \frac{\varphi'}{\varphi} \equiv 0,$$

which is a contradiction. Hence  $\psi_2 \not\equiv 0$ . It is clear now that  $\psi_2 \not\equiv 0$  cannot be a solution of (1.8) because  $\rho_p(\psi_2) < \infty$ . Then, by Lemma 2.1, we obtain  $\bar{\lambda}_{p+1}(g_2 - \varphi) = \lambda_{p+1}(g_2 - \varphi) = \rho_{p+1}(f)$ , i. e.,  $\bar{\lambda}_{p+1}(w) = \lambda_{p+1}(w) = \rho_{p+1}(f)$ .

## References

- [1] S. Bank, *General theorem concerning the growth of solutions of first-order algebraic differential equations*, Compositio Math. 25 (1972), 61-70.
- [2] B. Belaïdi, *Oscillation of fast growing solutions of linear differential equations in the unit disc*, Acta Univ. Sapientiae Math. 2 (2010), no. 1, 25-38.
- [3] B. Belaïdi, A. El Farissi, *Fixed points and iterated order of differential polynomial generated by solutions of linear differential equations in the unit disc*, J. Adv. Res. Pure Math. 3 (2011), no. 1, 161-172.
- [4] L. G. Bernal, *On growth  $k$ -order of solutions of a complex homogeneous linear differential equation*, Proc. Amer. Math. Soc. 101 (1987), no. 2, 317-322.
- [5] T. B. Cao and H. X. Yi, *The growth of solutions of linear differential equations with coefficients of iterated order in the unit disc*, J. Math. Anal. Appl. 319 (2006), no. 1, 278-294.
- [6] T. B. Cao, *The growth, oscillation and fixed points of solutions of complex linear differential equations in the unit disc*, J. Math. Anal. Appl. 352 (2009), no. 2, 739-748.
- [7] T. B. Cao, H. Y. Xu and C. X. Zhu, *On the complex oscillation of differential polynomials generated by meromorphic solutions of differential equations in the unit disc*, Proc. Indian Acad. Sci. Math. Sci. 120 (2010), no. 4, 481-493.



- [8] T. B. Cao and Z. S. Deng, *Solutions of non-homogeneous linear differential equations in the unit disc*, Ann. Polo. Math. 97(2010), no. 1, 51-61.
- [9] T. B. Cao, L. M. Li, J. Tu and H. Y. Xu, *Complex oscillation of differential polynomials generated by analytic solutions of differential equations in the unit disc*, Math. Commun. 16 (2011), no. 1, 205-214.
- [10] Z. X. Chen and K. H. Shon, *The growth of solutions of differential equations with coefficients of small growth in the disc*, J. Math. Anal. Appl. 297 (2004), no. 1, 285-304.
- [11] I. E. Chyzykov, G. G. Gundersen and J. Heittokangas, *Linear differential equations and logarithmic derivative estimates*, Proc. London Math. Soc. (3) 86 (2003), no. 3, 735-754.
- [12] A. El Farissi, B. Belaïdi and Z. Latreuch, *Growth and oscillation of differential polynomials in the unit disc*, Electron. J. Diff. Equ., Vol. 2010(2010), No. 87, 1-7.
- [13] W. K. Hayman, *Meromorphic functions*, Oxford Mathematical Monographs Clarendon Press, Oxford, 1964.
- [14] J. Heittokangas, *On complex differential equations in the unit disc*, Ann. Acad. Sci. Fenn. Math. Diss. 122 (2000), 1-54.
- [15] J. Heittokangas, R. Korhonen and J. Rättyä, *Fast growing solutions of linear differential equations in the unit disc*, Results Math. 49 (2006), no. 3-4, 265-278.
- [16] J. Heittokangas, R. Korhonen and J. Rättyä, *Growth estimates for solutions of linear complex differential equations*, Ann. Acad. Sci. Fenn. Math. 29 (2004), no. 1, 233-246.
- [17] L. Kinnunen, *Linear differential equations with solutions of finite iterated order*, Southeast Asian Bull. Math. 22 (1998), no. 4, 385-405.
- [18] I. Laine, *Nevanlinna theory and complex differential equations*, de Gruyter Studies in Mathematics, 15. Walter de Gruyter & Co., Berlin-New York, 1993.
- [19] I. Laine and J. Rieppo, *Differential polynomials generated by linear differential equations*, Complex Var. Theory Appl. 49 (2004), no. 12, 897-911.
- [20] I. Laine, *Complex differential equations*, Handbook of differential equations: ordinary differential equations. Vol. IV, 269-363, Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam, 2008.
- [21] Z. Latreuch, B. Belaïdi and Abdallah El Farissi, *Complex oscillation of differential polynomials in the unit disc*, Period. Math. Hungar. 66 (2013), no. 1, 45-60.
- [22] Z. Latreuch and B. Belaïdi, *Growth and oscillation of differential polynomials generated by complex differential equations*, Electron. J. Diff. Equ., Vol. 2013 (2013), No. 16, 1-14.

- [23] Z. Latreuch and B. Belaïdi, *Properties of solutions of complex differential equations in the unit disc*, International Journal of Analysis and Applications, Volume 4, Number 2 (2014), 159-173.
- [24] Z. Latreuch and B. Belaïdi, *Further results on the properties of differential polynomials generated by solutions of complex differential equations*, Submitted.
- [25] M. Tsuji, *Potential Theory in Modern Function Theory*, Chelsea, New York, (1975), reprint of the 1959 edition.

**DOI: 10.7862/rf.2014.7**

**Zinelâabidine Latreuch**

email: z.latreuch@gmail.com

**Benharrat Belaïdi – Corresponding author**

email: belaidi@univ-mosta.dz

Department of Mathematics  
Laboratory of Pure and Applied Mathematics  
University of Mostaganem (UMAB)  
B. P. 227 Mostaganem-(Algeria)

*Received 10.09.2013, Accepted 20.11.2013*

# On efficient evaluation of integrals entering boundary equations of 3D potential and elasticity theory

*Liliana Rybarska-Rusinek, Dawid Jaworski,  
Aleksandr Linkov*

ABSTRACT: The paper presents recurrent formulae for efficient evaluation of all the integrals needed for solving static 3D potential and elasticity problems by the boundary elements method. The power-type asymptoties for the density at edges of the boundary are accounted for explicitly.

AMS Subject Classification: *65M38, 33E05*

Keywords and Phrases: *potential and elasticity problems, singular and hypersingular boundary integral equations, boundary element method, higher order approximations, edge element, elliptic integrals*

## 1 Introduction

The purpose of the paper is to suggest an efficient general method for evaluation of the influence coefficients of the 3D boundary element method accounting for both smooth behaviour of the densities at internal parts of the boundary and power-type asymptotic behaviour near edges of the boundary.

Inspection of the boundary integrals equations of static 3D potential and elasticity theory [4] shows that it is sufficient to consider the function

$$\int_{S^q} \frac{f(y)}{R} dS_y, \quad (1.1)$$

and its spatial derivatives  $\partial/\partial x_i$ ,  $\partial^2/\partial x_i \partial x_j$ ,  $\partial^3/\partial x_i \partial x_j \partial x_k$ .

Herein,  $S^q$  is the surface of a boundary element;  $f(y)$  is a function to be properly approximated on the element;  $R = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + (x_3 - y_3)^2}$ , where  $x_1$ ,  $x_2$ ,  $x_3$  and  $y_1$ ,  $y_2$ ,  $y_3$  are global coordinates of the field and integration point, respectively.

## 2 Approximation of the boundary and density

We shall assume that, as usual (e.g. [1]), a curvilinear, in general, surface element is transformed into a plane element. The global coordinates are transformed to the local Cartesian coordinates of the plane element with the local axes  $y'_2, y'_3$  in the element plane; the origin  $O'$  is in the plane of the transformed element. Besides, we assume that the entries of Jacobian matrix, its determinant and the expression for  $R$  are expanded into power series in  $x'_i - y'_i$  and truncated. From now on, to simplify notation, we shall drop the prime in the transformed coordinates and refer (1.1) to a plane element in its local coordinates. Then  $y_1 = 0$  and the function  $f(y)$  is the product of the density depending on the local coordinates  $y_2, y_3$  and powers of  $y_2$  and  $y_3$ , which result from the truncated expressions mentioned.

Furthermore, we assume the plane element to be a trapezoid. This type of boundary elements includes as particular cases commonly used triangular, parallelogram, rectangular and square elements. Without loss of generality, we direct the  $y_2$ -axis along the trapezoid base, the  $y_3$ -axis orthogonal to it and we locate the origin in the lower left apex of the trapezoid (Fig. 1). For an edge element, we choose its edge as the base of the trapezoid. Then if the density near the edge has the power-type behaviour, it is described by the factor  $y_3^\alpha$  with  $0 \leq \alpha < 1$ . The general approximation of the function  $f(y)$  in (1.1) is of the form:




Figure 1: Trapezoidal element in local coordinates,  $b_b = 0$

$$f(y) = y_3^\alpha \sum_{k+l=0}^{m_p} c_{kl} y_2^{k+s} y_3^{l+q}, \quad 0 \leq \alpha < 1, \quad (2.1)$$

where  $m_p$  is the degree of a polynomial approximating the density,  $c_{kl}$  are coefficients of approximation,  $s$  and  $q$  are degrees arising from the coordinate transformation (for initially plane parts of the boundary  $s = q = 0$ ).

The two most important cases are: (i)  $\alpha = 0$  what corresponds to smooth behaviour of the density, and (ii)  $\alpha = 1/2$  what corresponds to square-root asymptotics typical for problem of linear fracture mechanics. Still, other exponents  $\alpha$  may arise

in approximations. For instance,  $\alpha = 2/3$  for fracturing impermeable rock by a Newtonian fluid. Therefore, it is reasonable to specify a particular value of  $\alpha$  at the end of the discussion.

Using (2.1) in (1.1) with  $S^q$  being the plane trapezoid of the height  $h$  (Fig. 1) implies that it is sufficient to consider the main integrals of the form:

$$A_\alpha^{kl}(x_1, x_2, x_3) = \int_0^h \left( y_3^{l+\alpha} \int_{y_2=a_b y_3+b_b}^{y_2=a_f y_3+b_f} \frac{(x_2 - y_2)^k}{R} dy_2 \right) dy_3. \quad (2.2)$$

and their partial derivatives  $\partial/\partial x_i$ ,  $\partial^2/\partial x_i \partial x_j$ ,  $\partial^3/\partial x_i \partial x_j \partial x_k$ .

### 3 Evaluation of the main integrals

The integrals (2.2) are evaluated recurrently by using starting integrals for  $k = 0$  and  $k = 1$ :

$$A_\alpha^{0j}(x_1, x_2, x_3) = - \int_0^h \left[ y_3^{j+\alpha} \ln[(x_2 - b_\xi - a_\xi y_3) + R_\xi] \right]_{\xi=b}^{\xi=f} dy_3, \quad (3.1)$$

$$A_\alpha^{1j}(x_1, x_2, x_3) = - \int_0^h \left[ y_3^{j+\alpha} R_\xi \right]_{\xi=b}^{\xi=f} dy_3, \quad (3.2)$$

where where  $R_\xi = \sqrt{x_1^2 + (x_2 - b_\xi - a_\xi y_3)^2 + (x_3 - y_3)^2}$ ; the symbol  $[ ]_{x=a}^{x=b}$  means the double substitution:  $[f(x)]_{x=a}^{x=b} = f(b) - f(a)$ .

For  $k \geq 2$ , the recurrent equations are:

$$A_\alpha^{kl} = -\frac{1}{k} \left( \int_0^h y_3^{\alpha+l} \left[ (x_2 - b_\xi - a_\xi y_3)^{k-1} R_\xi \right]_{\xi=b}^{\xi=f} dy_3 + \right. \\ \left. + (k-1)(x_1^2 + x_3^2) A_\alpha^{(k-2)l} - 2x_3(k-1) A_\alpha^{(k-2)(l+1)} + (k-1) A_\alpha^{(k-2)(l+2)} \right). \quad (3.3)$$

Note that the integral  $A_\alpha^{1j}$  in (3.2) is a particular case of the integrals on the r.h.s. of (3.3) when  $k = 1$ . Therefore, it remains to consider the integral on the r.h.s. of the (3.3) and the integral  $A_\alpha^{0j}$  defined by (3.1). For both of them, an analysis shows that they are promptly expressed as linear combinations of three standard terms:

$$\left[ \left[ y_3^{l+1+\alpha} \ln[(x_2 - b_\xi - a_\xi y_3) + R_\xi] \right]_{\xi=b}^{\xi=f} \right]_{y_3=0}^{y_3=h}, \quad (3.4)$$

$$\left[ u_s^\xi \int_0^h \frac{y_3^\alpha y_3^s}{R_\xi} dy_3 \right]_{\xi=b}^{\xi=f}, \quad (3.5)$$

$$\left[ \int_0^h \frac{y_3^\alpha (\tilde{A}y_3 + \tilde{B})}{R_0^2 R_\xi} dy_3 \right]_{\xi=b}^{\xi=f}, \quad (3.6)$$

where  $u_s^\xi$ ,  $\tilde{A}$  and  $\tilde{B}$  are known coefficients depending on  $\xi$ ,  $R_0^2 = x_1^2 + (x_3 - y_3)^2$ .

From (3.4)–(3.6) it follows that the problem is reduced to calculation of the integrals (3.5), (3.6) and their partial derivatives of the first, second and third order. Differentiation of (3.4), being trivial, we focus on the derivatives of the integrals (3.5) and (3.6).

## 4 Main integrals defining the first, second and third derivatives of standard terms

Evaluation of the first, second and third derivatives of the standard term (3.5) shows that it results in two new standard terms:

$$\left[ v_i^\xi \int_0^h \frac{y_3^\alpha}{(y_3 + z_\xi)^i R_\xi} dy_3 \right]_{\xi=b}^{\xi=f}, \quad \left[ \bar{v}_i^\xi \int_0^h \frac{y_3^\alpha}{(y_3 + \bar{z}_\xi)^i R_\xi} dy_3 \right]_{\xi=b}^{\xi=f}, \quad (4.1)$$

where  $v_i^\xi$  are known, in general complex, coefficients ( $i = 1, 2, 3$ );  $z_\xi$  is the complex root of the polynomial  $R_\xi^2$ , so that  $(1 + a_\xi^2)(y_3 + z_\xi)(y_3 + \bar{z}_\xi) = R_\xi^2$ ; the overbar denotes complex conjugation.

Similar analysis of the partial derivatives of the standard term (3.6) also yields two new standard terms:

$$\left[ w_j^\xi \int_0^h \frac{y_3^\alpha dy_3}{(y_3 + z_0)^j R_\xi} \right]_{\xi=b}^{\xi=f}, \quad \left[ \bar{w}_j^\xi \int_0^h \frac{y_3^\alpha dy_3}{(y_3 + \bar{z}_0)^j R_\xi} \right]_{\xi=b}^{\xi=f}, \quad (4.2)$$

where  $w_j^\xi$  are known, in general complex, coefficients;  $z_0 = -x_3 + ix_1$  is the root of  $R_0^2$ ,  $j = 1, 2, 3, 4$ , when  $x_1 \neq 0$ ; in the case  $x_1 = 0$  we have  $z_0 = \bar{z}_0 = -x_3$  and then  $j = 1, 2, \dots, 8$ . Actually (4.2) are particular cases of (4.1) when the root  $z_\xi$  of  $R_\xi^2$  is changed to the root  $z_0$  of  $R_0^2$ .

Noting that the second expression in (4.1) and (4.2) are the conjugated first ones, we come to the conclusion that the problem is reduced to evaluation of three types of integrals, at most:

$$\left[ u_\xi \int_0^h \frac{y_3^\alpha y_3^s}{R_\xi} dy_3 \right]_{\xi=b}^{\xi=f}, \quad \left[ v_\xi^i \int_0^h \frac{y_3^\alpha}{(y_3 + z_\xi)^i R_\xi} dy_3 \right]_{\xi=b}^{\xi=f}, \quad \left[ w_\xi^j \int_0^h \frac{y_3^\alpha dy_3}{(y_3 + z_0)^j R_\xi} \right]_{\xi=b}^{\xi=f}, \quad (4.3)$$

where  $i = 1, 2, 3$ ,  $j = 1, 2, 3, 4$  for  $x_1 \neq 0$  and  $j = 1, 2, \dots, 8$  for  $x_1 = 0$ . Emphasise that when representing the exponent  $\alpha$  as a proper rational fracture  $\alpha = n/m$ , ( $n < m$ ),

the integrals (4.3) may be evaluated recurrently. Below we give the explicit formulae for the cases most important for application:  $\alpha = 0$  and  $\alpha = 1/2$ . Before presenting them, we distinguish three cases which suggest simplifications.

**(i) Differentiation with respect to  $x_2$ .** In this case, we may avoid using the recursive equation (3.3) by the method suggested in the paper [5]. Specifically, by the relation  $\partial A^{kl}/\partial x_2 = -\partial A^{kl}/\partial y_2$  we obtain

$$\frac{\partial A^{kl}}{\partial x_2} = - \int_0^h \left[ \frac{y_3^{l+\alpha} (x_2 - b_\xi - a_\xi y_3)^k}{R_\xi} \right]_{\xi=b}^{\xi=f} dy_3. \quad (4.4)$$

This shows that differentiation with respect to  $x_2$  immediately leads to arithmetic operations with the expressions (4.1).

**(ii) Differentiation with respect to  $x_1$ .** By differentiating equation (3.3) with respect to  $x_1$ , we obtain:

$$\begin{aligned} \frac{\partial A_\alpha^{kl}}{\partial x_1} = & -\frac{1}{k} \left[ \int_0^h \frac{x_1 y_3^{\alpha+l} (x_2 - b_\xi - a_\xi y_3)^{k-1}}{R_\xi} dy_3 \right]_{\xi=b}^{\xi=f} + \\ & -\frac{k-1}{k} \left( 2x_1 A_\alpha^{(k-2)l} + (x_1^2 + x_3^2) \frac{\partial A_\alpha^{(k-2)l}}{\partial x_1} - 2x_3 \frac{\partial A_\alpha^{(k-2)(l+1)}}{\partial x_1} + \frac{\partial A_\alpha^{(k-2)(l+2)}}{\partial x_1} \right). \end{aligned} \quad (4.5)$$

The derivative of the starting integral  $\frac{\partial A_\alpha^{1l}}{\partial x_1}$  has the form of the first integral on the right hand side of the formula (4.5). Thus it is enough to consider the derivative of the starting integral  $A_\alpha^{0l}$ .

$$\frac{\partial A_\alpha^{0l}}{\partial x_1} = x_1 \int_0^h \left[ \frac{y_3^{l+\alpha} (x_2 - b_\xi - a_\xi y_3)}{R_0^2 R_\xi} \right]_{\xi=b}^{\xi=f} dy_3 - x_1 \int_0^h \left[ \frac{y_3^{l+\alpha}}{R_0^2} \right]_{\xi=b}^{\xi=f} dy_3. \quad (4.6)$$

The first integral after decomposition into a sum of real partial fractions is evaluated by arithmetic operations with the integrals (3.5) and (3.6). The second integral does not depend on  $\xi$ , therefore it is zero. We see that evaluation of partial derivatives, containing differentiation with respect to  $x_1$ , is reduced to evaluation of expressions of the forms (4.1) for  $i = 1, 2$  and (4.2) for  $j = 1, 2, 3$ .

**(iii) Double differentiation with respect to  $x_3$ .** Since the function  $1/R$  satisfies the Laplace equation when  $R \neq 0$ , we may avoid repeated differentiation with respect to  $x_3$  by using the equation

$$\frac{\partial^2 A_\alpha^{kl}}{\partial x_3^2} = - \left( \frac{\partial^2 A_\alpha^{kl}}{\partial x_1^2} + \frac{\partial^2 A_\alpha^{kl}}{\partial x_2^2} \right). \quad (4.7)$$

Then simplifications of points (i) and (ii) become available.

## 5 Case of smooth density ( $\alpha = 0$ )

In this case, all the integrals are evaluated analytically. Specifically, the integrals (3.5), (4.1) and (4.2) become respectively:

$$I_s = \int_0^h \frac{y_3^s}{\sqrt{(y_3 + z_\xi)(y_3 + \bar{z}_\xi)}} dy_3, \quad J_s = \int_0^h \frac{1}{(y_3 + z_\xi)^s \sqrt{(y_3 + z_\xi)(y_3 + \bar{z}_\xi)}} dy_3,$$

$$\text{and } K_s = \int_0^h \frac{dy_3}{(y_3 + z_0)^s \sqrt{(y_3 + z_\xi)(y_3 + \bar{z}_\xi)}}. \quad (5.1)$$

Each of them is evaluated recurrently, with starting expressions:

$$I_0 = J_0 = K_0 = 2 \left[ \ln \left( \sqrt{y_3 + z_\xi} + \sqrt{y_3 + \bar{z}_\xi} \right) \right]_{y_3=0}^{y_3=h},$$

$$I_1 = \left[ \sqrt{(y_3 + z_\xi)(y_3 + \bar{z}_\xi)} \right]_{y_3=0}^{y_3=h} - \operatorname{Re}(z_\xi) I_0, \quad (5.2)$$

$$K_1 = \frac{2 \left[ \arctan \left( \frac{\sqrt{y_3 + z_\xi} \sqrt{\bar{z}_\xi - z_0}}{\sqrt{y_3 + \bar{z}_\xi} \sqrt{z_0 - z_\xi}} \right) \right]_{y_3=0}^{y_3=h}}{\sqrt{z_0 - z_\xi} \sqrt{\bar{z}_\xi - z_0}}.$$

The recursive formulas are:

$$I_s = \frac{1}{s} \left( \left[ y_3^{s-1} \sqrt{(y_3 + z_\xi)(y_3 + \bar{z}_\xi)} \right]_{y_3=0}^{y_3=h} - (2s-1) \operatorname{Re}(z_\xi) I_{s-1} - (s-1) |z_\xi|^2 I_{s-2} \right), \quad (5.3)$$

$$J_s = \frac{1}{(s - \frac{1}{2})(z_\xi - \bar{z}_\xi)} \left( \left[ \frac{\sqrt{y_3 + \bar{z}_\xi}}{(y_3 + z_\xi)^{s-\frac{1}{2}}} \right]_{y_3=0}^{y_3=h} + (s-1) J_{s-1} \right), \quad (5.4)$$

$$K_s = \frac{1}{(s-1)(z_\xi - z_0)(\bar{z}_\xi - z_0)} \left( - \left[ \frac{\sqrt{(y_3 + z_\xi)(y_3 + \bar{z}_\xi)}}{(y_3 + z_0)^{s-1}} \right]_{y_3=0}^{y_3=h} + \left( s - \frac{3}{2} \right) (2z_0 - z_\xi - \bar{z}_\xi) K_{s-1} - (s-2) K_{s-2} \right). \quad (5.5)$$

Note that in the considered case, the representation of the trapezoid as a sum of right triangles and a rectangle, allows us to use also the efficient method suggested in [5].



## 6 The case of the density with square-root asymptotics near the element edge ( $\alpha = 1/2$ )

In this case, the starting integrals for evaluation of the integrals

$$I_s = \int_0^h \frac{y_3^s q dy_3}{\sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)}}, \quad J_s = \int_0^h \frac{dy_3}{(y_3 + z_\xi)^s \sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)}},$$

$$K_s = \int_0^h \frac{dy_3}{\sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)} (y_3 + z_0)^s},$$

are:

$$I_0 = J_0 = K_0 = \int_0^h \frac{dy_3}{\sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)}}, \quad (6.1)$$

$$I_1 = \int_0^h \frac{y_3}{\sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)}} dy_3, \quad (6.2)$$

$$J_1 = \int_0^h \frac{dy_3}{(y_3 + z_\xi) \sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)}}, \quad (6.3)$$

$$K_1 = \int_0^h \frac{dy_3}{\sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)} (y_3 + z_0)}, \quad (6.4)$$

$$K_2 = \int_0^h \frac{dy_3}{\sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)} (y_3 + z_0)^2}. \quad (6.5)$$

The recursive formulae are:

$$I_s = \frac{1}{(2s-1)} \left( 2 \left[ y_3^{s-2} \sqrt{y_3 (y_3 + z_\xi) (y_3 + \bar{z}_\xi)} \right]_{y_3=0}^{y_3=h} + \right. \\ \left. - 4 \operatorname{Re}(z_\xi) (s-1) I_{s-1} - |z_\xi|^2 (2s-3) I_{s-2} \right), \quad (6.6)$$

$$J_s = \frac{1}{(s-\frac{1}{2})(\bar{z}_\xi - z_\xi) z_\xi} \left[ \left[ \frac{\sqrt{(y_3 + \bar{z}_\xi) y_3}}{(y_3 + z_\xi)^{s-\frac{1}{2}}} \right]_{y_3=0}^{y_3=h} + \right. \\ \left. + (s-1)(\bar{z}_\xi - 2z_\xi) J_{s-1} + \left( s - \frac{3}{2} \right) J_{s-2} \right], \quad (6.7)$$

$$\begin{aligned}
K_s = & \frac{1}{2(s-1)(z_0 - z_\xi)(z_0 - \bar{z}_\xi)z_0} \left( \left[ \frac{2\sqrt{y_3(y_3 + z_\xi)(y_3 + \bar{z}_\xi)}}{(y_3 + z_0)^{s-1}} \right]_{y_3=0}^{y_3=h} + \right. \\
& \left. + (2s-5)K_{s-3} - 2(s-2)(3z_0 - \bar{z}_\xi - z_\xi)K_{s-2} \right) + \\
& + \frac{(2s-3)}{2(s-1)} \left( \frac{1}{z_0} + \frac{1}{z_0 - z_\xi} + \frac{1}{z_0 - \bar{z}_\xi} \right) K_{s-1}. \tag{6.8}
\end{aligned}$$

**Remark 6.1** For  $z_0 = 0$  (i.e.  $x_1 = 0, x_3 = 0$ ),

$$K_s = \int_0^h \frac{dy_3}{\sqrt{y_3(y_3 + z_\xi)(y_3 + \bar{z}_\xi)}(y_3)^s},$$

and the recursive formula becomes:

$$\begin{aligned}
K_s = & -\frac{1}{(s - \frac{1}{2})|z_\xi|^2} \left( \left[ y_3^{-(s-\frac{1}{2})} \sqrt{(y_3 + z_\xi)(y_3 + \bar{z}_\xi)} \right]_{y_3=0}^{y_3=h} + \right. \\
& \left. + \left( s - \frac{3}{2} \right) K_{s-2} + 2 \operatorname{Re}(z_\xi)(s-1)K_{s-1} \right), \tag{6.9}
\end{aligned}$$

with starting integrals:

$$K_{-1} = I_1, \quad K_0 = I_0. \tag{6.10}$$

Evaluation of the starting elliptic integrals  $I_0, I_1, J_1, K_1$  and  $K_2$  is efficiently performed by proper adjusting the Carlson algorithms as explained in the next section.

## 7 Efficient evaluation of standard elliptic integrals for problems involving cracks ( $\alpha = 1/2$ )

The conventional methods of evaluation the elliptic integrals employ Gauss and Landen transformations [6]. They converge quadratically and work well for elliptic integrals of the first and second kind. However, as emphasised in [6] and confirmed by our experience, they suffer from lost of significant digits for the integrals of the third kind needed for our purpose. In contrast, the Carlson algorithm provides a unified method for all the three kinds of integrals with extremely high efficiency. To use this algorithm, we introduce the new variable  $t$  defined by equation:

$$y_3 = \frac{1}{t + \frac{1}{h}}. \tag{7.1}$$

Then the starting integrals become:

$$I_0 = \frac{1}{|z_\xi|} \int_0^\infty \frac{dt}{\sqrt{\left(t + \frac{1}{h}\right) \left(t + \frac{1}{h} + \frac{1}{z_\xi}\right) \left(t + \frac{1}{h} + \frac{1}{\bar{z}_\xi}\right)}}, \quad (7.2)$$

$$I_1 = \frac{1}{|z_\xi|} \int_0^\infty \frac{dt}{\left(t + \frac{1}{h}\right)^{\frac{3}{2}} \sqrt{\left(t + \frac{1}{h} + \frac{1}{z_\xi}\right) \left(t + \frac{1}{h} + \frac{1}{\bar{z}_\xi}\right)}}, \quad (7.3)$$

$$J_1 = \frac{1}{z_\xi |z_\xi|} \int_0^\infty \frac{\left(t + \frac{1}{h}\right) dt}{\sqrt{\left(t + \frac{1}{h}\right) \left(t + \frac{1}{h} + \frac{1}{z_\xi}\right) \left(t + \frac{1}{h} + \frac{1}{\bar{z}_\xi}\right) \left(t + \frac{1}{h} + \frac{1}{z_0}\right)}}, \quad (7.4)$$

$$K_1 = \frac{1}{|z_\xi| z_0} \int_0^\infty \frac{\left(t + \frac{1}{h}\right) dt}{\sqrt{\left(t + \frac{1}{h}\right) \left(t + \frac{1}{h} + \frac{1}{z_\xi}\right) \left(t + \frac{1}{h} + \frac{1}{\bar{z}_\xi}\right) \left(t + \frac{1}{h} + \frac{1}{z_0}\right)}}, \quad (7.5)$$

$$K_2 = \frac{1}{|z_\xi| z_0^2} \int_0^\infty \frac{\left(t + \frac{1}{h}\right)^2 dt}{\sqrt{\left(t + \frac{1}{h}\right) \left(t + \frac{1}{h} + \frac{1}{z_\xi}\right) \left(t + \frac{1}{h} + \frac{1}{\bar{z}_\xi}\right) \left(t + \frac{1}{h} + \frac{1}{z_0}\right)^2}}. \quad (7.6)$$

They are promptly expressed in terms of Carlson integrals  $R_F$ ,  $R_D$  and  $R_J$  of the first, second and third kind, respectively, defined as:

$$\begin{aligned} R_F(x, y, z) &= \frac{1}{2} \int_0^\infty [(t+x)(t+y)(t+z)]^{-\frac{1}{2}} dt, \\ R_D(z, y, z) = R_J(x, y, z, z) &= \frac{3}{2} \int_0^\infty [(t+x)(t+y)]^{-\frac{1}{2}} (t+z)^{-\frac{3}{2}} dt, \\ R_J(x, y, z, p) &= \frac{3}{2} \int_0^\infty [(t+x)(t+y)(t+z)]^{-\frac{1}{2}} (t+p)^{-1} dt. \end{aligned} \quad (7.7)$$

In terms of the Carlson integrals, the starting integrals are:

$$I_0 = \frac{2}{|z_\xi|} R_F\left(\frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{\bar{z}_\xi}\right), \quad (7.8)$$

$$I_1 = \frac{2}{3|z_\xi|} R_D\left(\frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{\bar{z}_\xi}, \frac{1}{h}\right), \quad (7.9)$$

$$J_1 = \frac{I_0}{z_\xi} - \frac{2}{3z_\xi^2 |z_\xi|} R_D\left(\frac{1}{h}, \frac{1}{h} + \frac{1}{\bar{z}_\xi}, \frac{1}{h} + \frac{1}{z_\xi}\right), \quad (7.10)$$

$$K_1 = \frac{I_0}{z_0} - \frac{2}{3|z_\xi| z_0^2} R_J\left(\frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{\bar{z}_\xi}, \frac{1}{h} + \frac{1}{z_0}\right), \quad (7.11)$$

$$\begin{aligned}
K_2 = & -\frac{I_0}{2z_0^2} \left( 1 + \frac{z_\xi}{z_0 - z_\xi} + \frac{\bar{z}_\xi}{z_0 - \bar{z}_\xi} \right) - \frac{I_1}{2z_0^3} + \\
& + \frac{K_1}{2z_0} \left( 3 + \frac{z_\xi}{z_0 - z_\xi} + \frac{\bar{z}_\xi}{z_0 - \bar{z}_\xi} \right) + \frac{1}{|z_\xi| z_0^4} \frac{\sqrt{h}}{\left( \frac{1}{h} + \frac{1}{z_0} \right) \left| \frac{1}{h} + \frac{1}{z_\xi} \right|} + \\
& - \frac{1}{3|z_\xi| z_0^3} \left( \frac{z_\xi}{z_\xi - z_0} R_D \left( \frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_\xi} \right) + \frac{\bar{z}_\xi}{\bar{z}_\xi - z_0} R_D \left( \frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_\xi} \right) \right). \tag{7.12}
\end{aligned}$$

Finally, we need to evaluate five Carlson integrals only:

$$\begin{aligned}
R_D \left( \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} \right), & \quad R_D \left( \frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_\xi} \right), \\
R_D \left( \frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_\xi} \right), & \quad R_F \left( \frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_\xi} \right), \\
R_J \left( \frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_0} \right). & \tag{7.13}
\end{aligned}$$

The integrals  $R_D$  and  $R_F$  are evaluated very fast and accurately by algorithms presented by Carlson in the paper [3]. The same also refers to the integrals  $R_J$  when its last argument is not a negative real number. The case, when the last argument  $\left( \frac{1}{h} + \frac{1}{z_0} \right)$  of the integral  $R_J$  is a real negative number, is special. It occurs when the field point is within the strip  $x_1 = 0, 0 < x_3 < h$ . Then the integral  $R_J$  is a singular real Cauchy integral:

$$R_J \left( \frac{1}{h}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_\xi}, \frac{1}{h} + \frac{1}{z_0} \right) = \int_0^\infty \frac{dt}{\sqrt{t + \frac{1}{h}} \sqrt{f + gt + t^2} \left( t + \frac{1}{h} + \frac{1}{z_0} \right)}, \tag{7.14}$$

where  $f = |z_\xi|^2, g = 2 \operatorname{Re} z_\xi$ . In the paper [2] Carlson provides equations serving for efficient of this integral:

$$\begin{aligned}
R_J = & \frac{2c_{11}}{3c_{44}} \left[ -4x_3 \left( c_{14}^2 + \sqrt{c_{11}^2 c_{44}^2} \right) R_J(M^2, L_-^2, L_+^2, W_+^2) + \right. \\
& \left. -6R_F(M^2, L_-^2, L_+^2) + 3R_C(U^2, W^2) - 2R_C(P^2, Q^2) \right], \tag{7.15}
\end{aligned}$$

where

$$\begin{aligned}
c_{11}^2 &= 2 \left( f - \frac{g}{h} + \frac{1}{h^2} \right), \\
c_{14}^2 &= 2 \left( f - g \left( \frac{1}{h} + \frac{1}{2z_0} \right) + \frac{1}{h} \left( \frac{1}{h} + \frac{1}{z_0} \right) \right), \\
c_{44}^2 &= 2 \left( f - g \left( \frac{1}{h} + \frac{1}{z_0} \right) + \left( \frac{1}{h} + \frac{1}{z_0} \right)^2 \right), \tag{7.16}
\end{aligned}$$

and

$$\begin{aligned}
 M^2 &= 2\sqrt{f} + g, \\
 U^2 &= \frac{1}{h}, \\
 W^2 &= U^2 + \frac{1}{2}z_0c_{11}^2, \\
 W_+^2 &= M^2 + z_0(c_{14}^2 + c_{11}c_{44}), \\
 L_{\pm}^2 &= M^2 + \left(\frac{2}{h} - g\right) \pm c_{11}\sqrt{2}, \\
 Q^2 &= W^2 \left(1 + \frac{h}{z_0}\right), \\
 P^2 &= Q^2 - \frac{1}{2}z_0c_{44}^2, \\
 R_C(a, b) &= R_F(a, b, b).
 \end{aligned} \tag{7.17}$$

With using these equations, evaluation of the elliptic integrals, needed for problems involving cracks, becomes extremely efficient. Our experience shows that calculations of influence coefficients for square-root edge elements ( $\alpha = 1/2$ ) are performed as accurate and fast as those for ordinary elements ( $\alpha = 0$ ).

We believe that similar, highly efficient algorithms may be developed for any proper fraction  $\alpha = m/n$  ( $m < n$ ).

#### Acknowledgement

Authors gratefully acknowledge the support of the European Research Agency (FP7-PEOPLE-2009-IAPP Marie Curie IAPP transfer of knowledge programme, Project Reference #251475).

## References

- [1] Banerjee P. K., Butterfield R., Boundary element methods in engineering science, McGrawHill Book Co., UK, 1981.
- [2] Carlson B. C., A table of elliptic integrals: one quadratic factor, Mathematics of computation, 56, 1991, pp. 267-280.
- [3] Carlson B. C., Numerical computation of real or complex elliptic integrals, Numerical Algorithms, 10, 1995, pp. 13-26.
- [4] Linkov A. M., Boundary integral equations in elasticity theory, Kluwer Academic Publishers, Dordrecht-Boston-London, 2002.
- [5] Linkov A. M., Zoubkov V. V. and Kheib M. A., A method of solving three-dimensional problems of seam working and geological faults, J. Min. Sci., vol. 33 (4), 1997, pp. 3-18.
- [6] Zill D. G., Carlson B. C., Symmetric elliptic integrals of the third kind, Mathematics of computation, 24, 1970, pp. 199-214.

DOI: 10.7862/rf.2014.8

**Liliana Rybarska-Rusinek - corresponding author**

email: rybarska@prz.edu.pl

**Dawid Jaworski**

email: dawidj.poczta@gmail.com

**Aleksandr Linkov**

email: linkoval@prz.edu.pl

Department of Mathematics,  
Rzeszów University of Technology,  
al. Powstańców Warszawy 12,  
35-959 Rzeszów, Poland

*Received 10.10.2013, Accepted 18.12.2013*

## On differential sandwich theorems of analytic functions defined by certain generalized linear operator

*T. M. Seoudy, M. K. Aouf*

ABSTRACT: In this paper, we obtain some applications of first order differential subordination and superordination results involving certain linear operator and other linear operators for certain normalized analytic functions. Some of our results improve and generalize previously known results.

AMS Subject Classification: *30C45*

Keywords and Phrases: *Analytic function, Hadamard product, differential subordination, superordination, linear operator.*

### 1. Introduction

Let  $H(U)$  be the class of analytic functions in the open unit disk  $U = \{z \in \mathbb{C} : |z| < 1\}$  and let  $H[a, k]$  be the subclass of  $H(U)$  consisting of functions of the form:

$$f(z) = a + a_k z^k + a_{k+1} z^{k+1} \dots \quad (a \in C). \quad (1.1)$$

For simplicity  $H[a] = H[a, 1]$ . Also, let  $\mathcal{A}$  be the subclass of  $H(U)$  consisting of functions of the form:

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k. \quad (1.2)$$

If  $f, g \in H(U)$ , we say that  $f$  is subordinate to  $g$  or  $f$  is superordinate to  $g$ , written  $f(z) \prec g(z)$  if there exists a Schwarz function  $\omega$ , which (by definition) is analytic in  $U$  with  $\omega(0) = 0$  and  $|\omega(z)| < 1$  for all  $z \in U$ , such that  $f(z) = g(\omega(z))$ ,  $z \in U$ . Furthermore, if the function  $g$  is univalent in  $U$ , then we have the following equivalence, (cf., e.g., [6], [16] and [17]):

$$f(z) \prec g(z) \Leftrightarrow f(0) = g(0) \text{ and } f(U) \subset g(U).$$

Let  $\phi : \mathbb{C}^2 \times U \rightarrow \mathbb{C}$  and  $h(z)$  be univalent in  $U$ . If  $p(z)$  is analytic in  $U$  and satisfies the first order differential subordination:

$$\phi(p(z), zp'(z); z) \prec h(z), \quad (1.3)$$

then  $p(z)$  is a solution of the differential subordination (1.3). The univalent function  $q(z)$  is called a dominant of the solutions of the differential subordination (1.3) if  $p(z) \prec q(z)$  for all  $p(z)$  satisfying (1.3). A univalent dominant  $\tilde{q}$  that satisfies  $\tilde{q} \prec q$  for all dominants of (1.3) is called the best dominant. If  $p(z)$  and  $\phi(p(z), zp'(z); z)$  are univalent in  $U$  and if  $p(z)$  satisfies first order differential superordination:

$$h(z) \prec \phi(p(z), zp'(z); z), \quad (1.4)$$

then  $p(z)$  is a solution of the differential superordination (1.4). An analytic function  $q(z)$  is called a subordinant of the solutions of the differential superordination (1.4) if  $q(z) \prec p(z)$  for all  $p(z)$  satisfying (1.4). A univalent subordinant  $\tilde{q}$  that satisfies  $q \prec \tilde{q}$  for all subordinants of (1.4) is called the best subordinant. Using the results of Miller and Mocanu [17], Bulboaca [5] considered certain classes of first order differential subordinations as well as superordination-preserving integral operators [6]. Ali et al. [1], have used the results of Bulboaca [5] to obtain sufficient conditions for normalized analytic functions to satisfy:

$$q_1(z) \prec \frac{zf'(z)}{f(z)} \prec q_2(z),$$

where  $q_1$  and  $q_2$  are given univalent functions in  $U$  with  $q_1(0) = q_2(0) = 1$ . Also, Tuneski [25] obtained a sufficient condition for starlikeness of  $f$  in terms of the quantity  $\frac{f''(z)f(z)}{(f'(z))^2}$ . Recently, Shanmugam et al. [24] obtained sufficient conditions for the normalized analytic function  $f$  to satisfy

$$q_1(z) \prec \frac{f(z)}{zf'(z)} \prec q_2(z)$$

and

$$q_1(z) \prec \frac{z^2 f'(z)}{\{f(z)\}^2} \prec q_2(z).$$

They [24] also obtained results for functions defined by using Carlson-Shaffer operator [7], Ruscheweyh derivative [20] and Sălăgean operator [22].

For functions  $f$  given by (1.1) and  $g \in \mathcal{A}$  given by

$$g(z) = z + \sum_{k=2}^{\infty} b_k z^k, \quad (1.5)$$

the Hadamard product (or convolution) of  $f$  and  $g$  is defined by

$$(f * g)(z) = z + \sum_{k=2}^{\infty} a_k b_k z^k = (g * f)(z).$$



For functions  $f, g \in \mathcal{A}$ , we define the linear operator  $D_\lambda^n : \mathcal{A} \rightarrow \mathcal{A}$  ( $\lambda \geq 0, l \geq 0; n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, \mathbb{N} = \{1, 2, \dots\}$ ) by:

$$D_{\lambda,l}^0(f * g)(z) = (f * g)(z),$$

$$\begin{aligned} D_{\lambda,l}^1(f * g)(z) &= D_{\lambda,l}(f * g)(z) = (1 - \lambda)(f * g)(z) + \frac{\lambda}{(l + 1)z^{l-1}}(z^l(f * g)(z))', \\ &= z + \sum_{k=2}^{\infty} \left[ \frac{l + 1 + \lambda(k - 1)}{l + 1} \right] a_k b_k z^k \quad (\lambda \geq 0; l \geq 0), \end{aligned} \quad (1.6)$$

and ( in general )

$$\begin{aligned} D_{\lambda,l}^n(f * g)(z) &= D_{\lambda,l}(D_{\lambda,l}^{n-1}(f * g)(z)) \\ &= z + \sum_{k=2}^{\infty} \left[ \frac{l + 1 + \lambda(k - 1)}{l + 1} \right]^n a_k b_k z^k \end{aligned} \quad (1.7)$$

$$(\lambda \geq 0; l \geq 0; n \in \mathbb{N}_0).$$

From (1.7), we can easily deduce that

$$\lambda z (D_{\lambda,l}^n(f * g)(z))' = (l + 1) D_{\lambda,l}^{n+1}(f * g)(z) - (l + 1 - \lambda) D_{\lambda,l}^n(f * g)(z) \quad (1.8)$$

$$(\lambda > 0; l \geq 0; n \in \mathbb{N}_0).$$

We observe that the linear operator  $D_{\lambda,l}^n(f * g)(z)$  reduces to several interesting many other linear operators considered earlier for different choices of  $n, \lambda, l$  and the function  $g$  :

(i)  $D_{\lambda,0}^n(f * g)(z) = D_\lambda^n(f * g)(z)$ , where  $D_\lambda^n(f * g)(z)$  is linear operator which was defined by Aouf and Mostafa [3];

(ii) For  $g(z) = \frac{z}{1-z}$ , we have  $D_{\lambda,l}^n(f * g)(z) = I(n, \lambda, l)f(z)$ , where  $I(n, \lambda, l)$  is the generalized multiplier transformation which was introduced and studied by Cătaş et al. [8];

(iii) For  $\lambda = 1$  and  $g(z) = \frac{z}{1-z}$ , we see that  $D_{1,l}^n(f * g)(z) = I(n, l)f(z)$ , where  $I(n, l)f(z)$  is the multiplier transformation (see [9]);

(iv) For  $l = 0$  and  $g(z) = \frac{z}{1-z}$ , we see that  $D_{\lambda,0}^n(f * g)(z) = D_\lambda^n f(z)$  where  $D_\lambda^n$  is the generalized Sălăgean operator ( or Al-Oboudi operator [2] ) which yield Sălăgean operator  $D^n$  for  $\lambda = 1$  introduced and studied by Sălăgean [22];

(v) For  $l = 0$  and

$$g(z) = z + \sum_{k=2}^{\infty} \Gamma_k [a_1; b_1] z^k, \quad (1.9)$$

$$\Gamma_k [a_1; b_1] = \frac{(a_1)_{k-1} \dots (a_q)_{k-1}}{(b_1)_{k-1} \dots (b_s)_{k-1} (1)_{k-1}} \quad (1.10)$$

$$(a_i \in \mathbb{C}; i = 1, \dots, q; b_j \in \mathbb{C} \setminus \mathbb{Z}_0^- = \{0, -1, -2, \dots\}; j = 1, \dots, s; q \leq s + 1; q, s \in \mathbb{N}_0),$$

where

$$(x)_k = \begin{cases} 1 & (k = 0; x \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}) \\ x(x+1)\dots(x+k-1) & (k \in \mathbb{N}; x \in \mathbb{C}), \end{cases}$$

we have  $D_{\lambda,0}^n(f * g)(z) = D_{\lambda}^n(a_1, b_1)f(z)$ , where  $D_{\lambda}^n(a_1, b_1)$  is the linear operator which was introduced and studied by Selvaraj and Karthikeyan [23]. The operator  $D_{\lambda}^n(a_1, b_1)f(z)$ , contains in turn many interesting operators such as, Dziok-Srivastava operator [10] ( see also [11]), Hohlov linear operator (see [13]), the Carlson-Shaffer linear operator (see [7] and [21] ), the Ruscheweyh derivative operator (see [20]), the Bernardi-Libera-Livingston operator ( see [4], [14] and [15]) and Owa-Srivastava fractional derivative operator (see [19]);

(iv) For  $g(z)$  of the form (1.9), we obtain

$$D_{\lambda,l}^n(f * g)(z) = I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z) = z + \sum_{k=2}^{\infty} \left[ \frac{l+1+\lambda(k-1)}{l+1} \right]^n \Gamma_k[a_1; b_1] z^k, \quad (1.11)$$

where the operator  $I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)$  is introduced and studied by El-Ashwah and Aouf [12].

In this paper, we will derive several subordination and superordination results and sandwich results involving the operator  $D_{\lambda,l}^n(f * g)$  and some of its special chooses of  $n, l, \lambda$  and the function  $g(z)$ .

## 2. Definitions and Preliminaries

In order to prove our subordinations and superordinations, we need the following definition and lemmas.

**Definition 1** [17]. Denote by  $Q$ , the set of all functions  $f$  that are analytic and injective on  $U \setminus E(f)$ , where

$$E(f) = \left\{ \zeta \in \partial U : \lim_{z \rightarrow \zeta} f(z) = \infty \right\},$$

and are such that  $f'(\zeta) \neq 0$  for  $\zeta \in \partial U \setminus E(f)$ .

**Lemma 1** [24]. Let  $q(z)$  be univalent in  $U$  with  $q(0) = 1$ . Let  $\alpha \in \mathbb{C}; \gamma \in \mathbb{C}^*$ , further assume that

$$\Re \left\{ 1 + \frac{zq''(z)}{q'(z)} \right\} > \max \left\{ 0, -\Re \left( \frac{\alpha}{\gamma} \right) \right\}. \quad (2.1)$$

If  $p(z)$  is analytic in  $U$ , and

$$\alpha p(z) + \gamma zp'(z) \prec \alpha q(z) + \gamma zq'(z),$$

then  $p(z) \prec q(z)$  and  $q(z)$  is the best dominant.

**Lemma 2** [24]. Let  $q(z)$  be convex univalent in  $U$ ,  $q(0) = 1$ . Let  $\alpha \in \mathbb{C}$ ;  $\gamma \in \mathbb{C}^*$  and  $\Re\left(\frac{\alpha}{\gamma}\right) > 0$ . If  $p(z) \in H[q(0), 1] \cap Q$ ,  $\alpha p(z) + \gamma zp'(z)$  is univalent in  $U$  and

$$\alpha q(z) + \gamma zq'(z) \prec \alpha p(z) + \gamma zp'(z),$$

then  $q(z) \prec p(z)$  and  $q(z)$  is the best subordinant.

### 3. Sandwich Results

Unless otherwise mentioned, we assume throughout this paper that  $l \geq 0, \lambda > 0, n \in \mathbb{N}_0$  and  $g(z)$  is given by (1.5).

**Theorem 1.** Let  $q(z)$  be univalent in  $U$  with  $q(0) = 1$ , and  $\gamma \in \mathbb{C}^*$ . Further, assume that

$$\Re\left\{1 + \frac{zq''(z)}{q'(z)}\right\} > \max\left\{0, -\Re\left(\frac{1}{\gamma}\right)\right\}. \tag{3.1}$$

If  $f, g \in \mathcal{A}$  satisfy the following subordination condition:

$$\begin{aligned} \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{1 - \frac{D_{\lambda,l}^n(f * g)(z)D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2}\right\} \\ \prec q(z) + \gamma zq'(z), \end{aligned} \tag{3.2}$$

then

$$\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} \prec q(z)$$

and  $q(z)$  is the best dominant.

**Proof.** Define a function  $p(z)$  by

$$p(z) = \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} \quad (z \in U). \tag{3.3}$$

Then the function  $p(z)$  is analytic in  $U$  and  $p(0) = 1$ . Therefore, differentiating (3.3) logarithmically with respect to  $z$  and using the identity (1.8) in the resulting equation, we have

$$\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{1 - \frac{D_{\lambda,l}^n(f * g)(z)D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2}\right\} = p(z) + \gamma zp'(z),$$

that is,

$$p(z) + \gamma zp'(z) \prec q(z) + \gamma zq'(z).$$

Therefore, Theorem 1 now follows by applying Lemma 1. ■

Putting  $q(z) = \frac{1+Az}{1+Bz}$  ( $-1 \leq B < A \leq 1$ ) in Theorem 1, we obtain the following corollary.

**Corollary 1.** Let  $\gamma \in \mathbb{C}^*$  and

$$\Re \left\{ \frac{1-Bz}{1+Bz} \right\} > \max \left\{ 0, -\Re \left( \frac{1}{\gamma} \right) \right\}.$$

If  $f, g \in \mathcal{A}$  satisfy the following subordination condition:

$$\begin{aligned} \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z)D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\} \\ \prec \frac{1+Az}{1+Bz} + \gamma \frac{(A-B)z}{(1+Bz)^2}, \end{aligned}$$

then

$$\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} \prec \frac{1+Az}{1+Bz}$$

and the function  $\frac{1+Az}{1+Bz}$  is the best dominant.

Taking  $g(z) = \frac{z}{1-z}$  in Theorem 1, we obtain the following subordination result for the generalized multiplier transformation  $I(n, \lambda, l)$ .

**Corollary 2.** Let  $q(z)$  be univalent in  $U$  with  $q(0) = 1$ , and  $\gamma \in \mathbb{C}^*$ . Further assume that (3.1) holds. If  $f \in \mathcal{A}$  satisfies the following subordination condition:

$$\begin{aligned} \frac{I(n, \lambda, l)f(z)}{I(n+1, \lambda, l)f(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{I(n, \lambda, l)f(z)I(n+2, \lambda, l)f(z)}{[I(n+1, \lambda, l)f(z)]^2} \right\} \\ \prec q(z) + \gamma z q'(z), \end{aligned}$$

then

$$\frac{I(n, \lambda, l)f(z)}{I(n+1, \lambda, l)f(z)} \prec q(z)$$

and  $q(z)$  is the best dominant.

Taking  $g(z)$  of the form (1.9) in Theorem 1, we obtain the following subordination result for the operator  $I_{q,s,\lambda}^{n,l}(a_1; b_1)$ .

**Corollary 3.** Let  $q(z)$  be univalent in  $U$  with  $q(0) = 1$ , and  $\gamma \in \mathbb{C}^*$ . Further assume that (3.1) holds. If  $f \in \mathcal{A}$  satisfies the following subordination condition:

$$\begin{aligned} \frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)}{I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)I_{q,s,\lambda}^{n+2,l}(a_1, b_1)f(z)}{[I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)]^2} \right\} \\ \prec q(z) + \gamma z q'(z), \end{aligned}$$

then

$$\frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)}{I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)} \prec q(z)$$

and  $q(z)$  is the best dominant.

Taking  $l = 0, \lambda = 1$  and  $g(z) = \frac{z}{1-z}$  in Theorem 1, we obtain the following subordination result for Sălăgean operator which improves the result of Shanmugam et al. [24, Theorem 5.1] and obtained by Nechita [18].

**Corollary 4** [18, Corollary 7]. Let  $q(z)$  be univalent in  $U$  with  $q(0) = 1$ , and  $\gamma \in \mathbb{C}^*$ . Further assume that (3.1) holds. If  $f \in \mathcal{A}$  satisfies the following subordination condition:

$$\frac{D^n f(z)}{D^{n+1} f(z)} + \gamma \left\{ 1 - \frac{D^n f(z) D^{n+2} f(z)}{[D^{n+1} f(z)]^2} \right\} \prec q(z) + \gamma z q'(z),$$

then

$$\frac{D^n f(z)}{D^{n+1} f(z)} \prec q(z)$$

and  $q(z)$  is the best dominant.

Now, by appealing to Lemma 2 it can be easily prove the following theorem.

**Theorem 2.** Let  $q(z)$  be convex univalent in  $U$  with  $q(0) = 1$ . Let  $\gamma \in \mathbb{C}$  with  $\Re(\bar{\gamma}) > 0$ . If  $f, g \in \mathcal{A}$  such that  $\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} \in H[q(0), 1] \cap Q$ ,

$$\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z) \cdot D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\}$$

is univalent in  $U$ , and the following superordination condition

$$q(z) + \gamma z q'(z) \prec \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z) \cdot D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\}$$

holds, then

$$q(z) \prec \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)}$$

and  $q(z)$  is the best subdominant.

Taking  $q(z) = \frac{1+Bz}{1+Bz}$  ( $-1 \leq B < A \leq 1$ ) in Theorem 2, we obtain the following corollary.

**Corollary 5.** Let  $\gamma \in \mathbb{C}$  with  $\Re(\bar{\gamma}) > 0$ . If  $f, g \in \mathcal{A}$  such that  $\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} \in H[q(0), 1] \cap Q$ ,

$$\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z) D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\}$$

is univalent in  $U$ , and the following superordination condition

$$\frac{1 + Az}{1 + Bz} + \gamma \frac{(A - B)z}{(1 + Bz)^2} \prec \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z)D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\}$$

holds, then

$$\frac{1 + Az}{1 + Bz} \prec \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)}$$

and  $q(z)$  is the best subordinated.

Taking  $g(z) = \frac{z}{1-z}$  in Theorem 2, we obtain the following superordination result for the generalized multiplier transformation  $I(n, \lambda, l)$ .

**Corollary 6.** Let  $q(z)$  be convex univalent in  $U$  with  $q(0) = 1$ . Let  $\gamma \in \mathbb{C}$  with  $\Re(\bar{\gamma}) > 0$ . If  $f, g \in \mathcal{A}$  such that  $\frac{I(n, \lambda, l)f(z)}{I(n+1, \lambda, l)f(z)} \in H[q(0), 1] \cap Q$ ,

$$\frac{I(n, \lambda, l)f(z)}{I(n+1, \lambda, l)f(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{I(n, \lambda, l)f(z) \cdot I(n+2, \lambda, l)f(z)}{[I(n+1, \lambda, l)f(z)]^2} \right\}$$

is univalent in  $U$ , and the following superordination condition

$$q(z) + \gamma z q'(z) \prec \frac{I(n, \lambda, l)f(z)}{I(n+1, \lambda, l)f(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{I(n, \lambda, l)f(z) \cdot I(n+2, \lambda, l)f(z)}{[I(n+1, \lambda, l)f(z)]^2} \right\}$$

holds, then

$$q(z) \prec \frac{I(n, \lambda, l)f(z)}{I(n+1, \lambda, l)f(z)}$$

and  $q(z)$  is the best subordinated.

Taking  $g(z)$  of the form (1.9) in Theorem 2, we obtain the following superordination result for the operator  $I_{q,s,\lambda}^{n,l}(a_1; b_1)$ .

**Corollary 7.** Let  $q(z)$  be convex univalent in  $U$  with  $q(0) = 1$ . Let  $\gamma \in \mathbb{C}$  with  $\Re(\bar{\gamma}) > 0$ . If  $f, g \in \mathcal{A}$  such that  $\frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)}{I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)} \in H[q(0), 1] \cap Q$ ,

$$\frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)}{I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z) I_{q,s,\lambda}^{n+2,l}(a_1, b_1)f(z)}{[I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)]^2} \right\}$$

is univalent in  $U$ , and the following superordination condition

$$q(z) + \gamma z q'(z) \prec \frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)}{I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)I_{q,s,\lambda}^{n+2,l}(a_1, b_1)f(z)}{[I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)]^2} \right\}$$

holds, then

$$q(z) \prec \frac{I_{q,s,\lambda}^{n,l}(a_1, b_1)f(z)}{I_{q,s,\lambda}^{n+1,l}(a_1, b_1)f(z)}$$

and  $q(z)$  is the best subordinant.

Taking  $l = 0, \lambda = 1$  and  $g(z) = \frac{z}{1-z}$  in Theorem 2, we obtain the following superordination result for Sălăgean operator which improves the result of Shanmugam et al. [24, Theorem 5.2] and obtained by Nechita [18].

**Corollary 8** [18, Corollary 12]. Let  $q(z)$  be convex univalent in  $U$  with  $q(0) = 1$ . Let  $\gamma \in \mathbb{C}$  with  $\Re(\bar{\gamma}) > 0$ . If  $f \in \mathcal{A}$  such that  $\frac{D^n f(z)}{D^{n+1} f(z)} \in H[q(0), 1] \cap Q$ ,

$$\frac{D^n f(z)}{D^{n+1} f(z)} + \gamma \left\{ 1 - \frac{D^n f(z).D^{n+2} f(z)}{[D^{n+1} f(z)]^2} \right\}$$

is univalent in  $U$ , and the following superordination condition

$$q(z) + \gamma z q'(z) \prec \frac{D^n f(z)}{D^{n+1} f(z)} + \gamma \left\{ 1 - \frac{D^n f(z).D^{n+2} f(z)}{[D^{n+1} f(z)]^2} \right\}$$

holds, then

$$q(z) \prec \frac{D^n f(z)}{D^{n+1} f(z)}$$

and  $q(z)$  is the best subordinant.

Combining Theorem 1 and Theorem 2, we get the following sandwich theorem for the linear operator  $D_{\lambda,l}^n(f * g)$ .

**Theorem 3.** Let  $q_1(z)$  be convex univalent in  $U$  with  $q_1(0) = 1$ ,  $\gamma \in \mathbb{C}$  with  $\Re(\bar{\gamma}) > 0$ ,  $q_2(z)$  be univalent in  $U$  with  $q_2(0) = 1$ , and satisfies (3.1). If  $f, g \in \mathcal{A}$  such that  $\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} \in H[q_2(0), 1] \cap Q$ ,

$$\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z).D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\}$$

is univalent in  $U$ , and

$$\begin{aligned} q_1(z) + \gamma z q_1'(z) &< \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z) \cdot D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\} \\ &< q_2(z) + \gamma z q_2'(z) \end{aligned}$$

holds, then

$$q_1(z) < \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} < q_2(z)$$

and  $q_1(z)$  and  $q_2(z)$  are, respectively, the best subordinant and the best dominant.

Taking  $q_i(z) = \frac{1+A_i z}{1+B_i z}$  ( $i = 1, 2; -1 \leq B_2 \leq B_1 < A_1 \leq A_2 \leq 1$ ) in Theorem 3, we obtain the following corollary.

**Corollary 9.** Let  $\gamma \in \mathbb{C}$  with  $\Re(\bar{\gamma}) > 0$ . If  $f, g \in \mathcal{A}$  such that  $\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} \in H[q(0), 1] \cap Q$ ,

$$\frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} + \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z) \cdot D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\}$$

is univalent in  $U$ , and

$$\begin{aligned} \frac{1+A_1 z}{1+B_1 z} + \gamma \frac{(A_1 - B_1)z}{(1+B_1 z)^2} &< \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} \\ &+ \frac{\gamma(l+1)}{\lambda} \left\{ 1 - \frac{D_{\lambda,l}^n(f * g)(z) \cdot D_{\lambda,l}^{n+2}(f * g)(z)}{[D_{\lambda,l}^{n+1}(f * g)(z)]^2} \right\} \\ &< \frac{1+A_2 z}{1+B_2 z} + \gamma \frac{(A_2 - B_2)z}{(1+B_2 z)^2} \end{aligned}$$

holds, then

$$\frac{1+A_1 z}{1+B_1 z} < \frac{D_{\lambda,l}^n(f * g)(z)}{D_{\lambda,l}^{n+1}(f * g)(z)} < \frac{1+A_2 z}{1+B_2 z}$$

and  $\frac{1+A_1 z}{1+B_1 z}$  and  $\frac{1+A_2 z}{1+B_2 z}$  are, respectively, the best subordinant and the best dominant.

Taking  $l = 0, \lambda = 1$  and  $g(z) = \frac{z}{1-z}$  in Theorem 3, we obtain the following sandwich result for Sălăgean operator which improves the result of Shanmugam et al. [24, Theorem 5.3].

**Corollary 10.** Let  $q_1(z)$  be convex univalent in  $U$  with  $q_1(0) = 1$ ,  $\gamma \in \mathbb{C}$  with  $\Re(\bar{\gamma}) > 0$ ,  $q_2(z)$  be univalent in  $U$  with  $q_2(0) = 1$ , and satisfies (3.1). If  $f \in \mathcal{A}$  such



that  $\frac{D^n f(z)}{D^{n+1} f(z)} \in H[q_2(0), 1] \cap Q$ ,

$$\frac{D^n f(z)}{D^{n+1} f(z)} + \gamma \left\{ 1 - \frac{D^n f(z) \cdot D^{n+2} f(z)}{[D^{n+1} f(z)]^2} \right\}$$

is univalent in  $U$ , and

$$q_1(z) + \gamma z q_1'(z) \prec \frac{D^n f(z)}{D^{n+1} f(z)} + \gamma \left\{ 1 - \frac{D^n f(z) \cdot D^{n+2} f(z)}{[D^{n+1} f(z)]^2} \right\} \prec q_2(z) + \gamma z q_2'(z)$$

holds, then

$$q_1(z) \prec \frac{D^n f(z)}{D^{n+1} f(z)} \prec q_2(z)$$

and  $q_1(z)$  and  $q_2(z)$  are, respectively, the best subordinant and the best dominant.

**Remarks** (i) Combining Corollary 2 and Corollary 6, we obtain similar sandwich theorem for the generalized multiplier transformation  $I(n, \lambda, l)$ ;

(ii) Combining Corollary 3 and Corollary 7, we obtain similar sandwich theorems for the operator  $I_{q,s,\lambda}^{n,l}(a_1, b_1)$ ;

(iii) Taking  $l = 0$  and  $g(z) = \frac{z}{1-z}$  in Theorems 1, 2 and 3, respectively, we obtain the results obtained by Nechita [18, Theorems 5, 10 and Corollary 13, respectively];

(iv) Taking  $n = l = 0, \lambda = 1$  and  $g(z) = \frac{z}{1-z}$  in Theorems 1, 2 and 3, respectively, we obtain the results obtained by Shanmugam et al. [24, Theorems 3.1, 3.2 and Corollary 3.3, respectively].

## References

- [1] R. M. Ali, V. Ravichandran and K. G. Subramanian, Differential sandwich theorems for certain analytic functions, *Far East J. Math. Sci.* 15 (2004), no. 1, 87-94.
- [2] F. M. Al-Oboudi, On univalent functions defined by a generalized Sălăgean operator, *Internat. J. Math. Math. Sci.*, 27 (2004), 1429-1436.
- [3] M. K. Aouf and A. O. Mostafa, Sandwich theorems for analytic functions defined by convolution, *Acta Univ. Apulensis*, 21(2010), 7-20.
- [4] S. D. Bernardi, Convex and starlike univalent functions, *Trans. Amer. Math. Soc.* 135 (1969), 429-446.
- [5] T. Bulboacă, Classes of first order differential subordinations, *Demonstratio Math.* 35 (2002), no. 2, 287-292.
- [6] T. Bulboaca, *Differential Subordinations and Superordinations, Recent Results*, House of Scientific Book Publ., Cluj-Napoca, 2005.

- [7] B. C. Carlson and D. B. Shaffer, Starlike and prestarlike hypergeometric functions, *SIAM J. Math. Anal.*, 15 (1984), 737-745.
- [8] A. Cătaş, G. I. Oros and G. Oros, Differential subordinations associated with multiplier transformations, *Abstract Appl. Anal.*, 2008 (2008), ID 845724, 1-11.
- [9] N. E. Cho and T. G. Kim, Multiplier transformations and strongly close-to-convex functions, *Bull. Korean Math. Soc.*, 40 (2003), no. 3, 399-410.
- [10] J. Dziok and H. M. Srivastava, Classes of analytic functions associated with the generalized hypergeometric function, *Appl. Math. Comput.*, 103 (1999), 1-13.
- [11] J. Dziok and H. M. Srivastava, Certain subclasses of analytic functions associated with the generalized hypergeometric function, *Integral Transform. Spec. Funct.*, 14 (2003), 7-18.
- [12] R. M. El-Ashwah and M. K. Aouf, Differential subordination and superordination for certain subclasses of  $p$ -valent functions, *Math. Comput. Modelling*, 51(2010), 349-360.
- [13] Yu. E. Hohlov, Operators and operations in the univalent functions, *Izv. Vyssh. Učebn. Zaved. Mat.*, 10 (1978), 83-89 ( in Russian).
- [14] R. J. Libera, Some classes of regular univalent functions, *Proc. Amer. Math. Soc.*, 16 (1965), 755-658.
- [15] A. E. Livingston, On the radius of univalence of certain analytic functions, *Proc. Amer. Math. Soc.*, 17 (1966), 352-357.
- [16] S. S. Miller and P. T. Mocanu, *Differential Subordination: Theory and Applications*, Series on Monographs and Textbooks in Pure and Applied Mathematics, Vol. 225, Marcel Dekker Inc., New York and Basel, 2000.
- [17] S. S. Miller and P. T. Mocanu, Subordinates of differential superordinations, *Complex Variables*, 48 (2003), no. 10, 815-826.
- [18] V. O. Nechita, Differential subordinations and superordinations for analytic functions defined by the generalized Sălăgean derivative, *Acta Univ. Apulensis*, 16(2008), 143-156.
- [19] S. Owa and H. M. Srivastava, Univalent and starlike generalized hypergeometric functions, *Canad. J. Math.* 39 (1987), 1057-1077.
- [20] St. Ruscheweyh, New criteria for univalent functions, *Proc. Amer. Math. Soc.*, 49 (1975), 109-115.
- [21] H. Saitoh, A linear operator and its applications of first order differential subordinations, *Math. Japon.* 44 (1996), 31-38.
- [22] G. S. Sălăgean, Subclasses of univalent functions, *Lecture Notes in Math.* (Springer-Verlag) 1013, (1983), 362 - 372 .

- [23] C. Selvaraj and K. R. Karthikeyan, Differential subordination and superordination for certain subclasses of analytic functions, *Far East J. Math. Sci. (FJMS)*, 29 (2008), no. 2, 419-430.
- [24] T. N. Shanmugam, V. Ravichandran and S. Sivasubramanian, Differential sandwich theorems for some subclasses of analytic functions, *J. Austr.Math. Anal. Appl.*, 3 (2006), no. 1, Art. 8, 1-11.
- [25] N. Tuneski, On certain sufficient conditions for starlikeness, *Internat. J. Math. Math. Sci.*, 23 (2000), no. 8, 521-527.

**DOI: 10.7862/rf.2014.9**

**T. M. Seoudy - corresponding author**

email: [tms00@fayoum.edu.eg](mailto:tms00@fayoum.edu.eg)

Department of Mathematics,  
Faculty of Science,  
Fayoum University, Fayoum 63514, Egypt

**M. K. Aouf**

email: [mkaouf127@yahoo.com](mailto:mkaouf127@yahoo.com)

Department of Mathematics,  
Faculty of Science  
Mansoura 35516

*Received 01.08.2013, Accepted 22.01.2014*

## Certain subordination results on the convolution of analytic functions

Poonam Sharma, Rajesh K. Maurya

ABSTRACT: In this paper, certain subordination results on the convolution of finite number of analytic functions are derived. Our results include a sufficiency condition for convexity of the convolution of analytic functions  $f_i$  satisfying  $f'_i \in \mathcal{P}(\alpha_i)$  ( $\alpha_i \leq 1, i = 1, 2, \dots, n$ ).

AMS Subject Classification: 30C45

Keywords and Phrases: Analytic function, Convolution (Hadamard product), Subordination, Convex functions, Zeta function

### 1 Introduction

Let  $\mathcal{H}(\mathbb{U})$  denote a class of all analytic functions defined in the open unit disk  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ . For  $a \in \mathbb{C}, j \in \mathbb{N}$ , let

$$\mathcal{H}[a, j] = \{f \in \mathcal{H}(\mathbb{U}) : f(z) = a + a_j z^j + a_{j+1} z^{j+1} + \dots\}.$$

We denote the special class of  $\mathcal{H}[0, 1]$  by  $\mathcal{A}$  whose members are of the form:

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k, \quad z \in \mathbb{U}. \quad (1.1)$$

Let  $\mathcal{K}$  denote a subclass of  $\mathcal{A}$  whose members are convex (univalent) in  $\mathbb{U}$  and satisfying

$$\Re \left( 1 + \frac{z f''(z)}{f'(z)} \right) > 0, \quad z \in \mathbb{U}.$$

For two functions  $p, q \in \mathcal{H}(\mathbb{U})$ , we say  $p$  is *subordinate* to  $q$ , or  $q$  is *superordinate* to  $p$  in  $\mathbb{U}$  and write  $p(z) \prec q(z), z \in \mathbb{U}$ , if there exists a Schwarz function  $\omega$ , analytic in  $\mathbb{U}$  with  $\omega(0) = 0$ , and  $|\omega(z)| < 1, z \in \mathbb{U}$  such that  $p(z) = q(\omega(z)), z \in \mathbb{U}$ . Furthermore, if the function  $q$  is univalent in  $\mathbb{U}$ , then we have following equivalence:

$$p(z) \prec q(z) \Leftrightarrow p(0) = q(0) \text{ and } p(\mathbb{U}) \subset q(\mathbb{U}). \quad (1.2)$$

Let  $\mathcal{P}_\alpha$  denote a class of functions  $p \in \mathcal{H}(\mathbb{U})$  satisfying  $p(0) = 1$  and

$$p(z) \prec q_\alpha(z) := \frac{1 + (1 - 2\alpha)z}{1 - z}, \alpha \leq 1, z \in \mathbb{U}. \quad (1.3)$$

Convolution (or Hadamard product)  $*$  of the functions  $g_1(z)$  and  $g_2(z)$  of the form:

$$g_1(z) = \sum_{k=0}^{\infty} a_k z^k \text{ and } g_2(z) = \sum_{k=0}^{\infty} b_k z^k, \quad (1.4)$$

is defined by

$$g_1(z) * g_2(z) = (g_1 * g_2)(z) = \sum_{k=0}^{\infty} a_k b_k z^k = (g_2 * g_1)(z). \quad (1.5)$$

In 1973, Rusheweyh and Sheil-Small [3] proved the Pòlya-Schoenberg conjecture which shows that the convolution of two convex functions is again a convex function. Due to this convexity preserving property, attempts are made to involve and study convolutions in the Geometric Function Theory.

In this paper, we derive certain subordination results on the convolution of any finite number of analytic functions. Mainly, by applying the subordination principle, a sufficiency condition for convexity of  $\phi(z) := (f_1 * f_2 * \dots * f_n)(z)$  which is a convolution of analytic functions  $f_i \in \mathcal{A}$  ( $i = 1, 2, \dots, n$ ) such that  $f'_i \in \mathcal{P}(\alpha_i)$  ( $\alpha_i \leq 1, i = 1, 2, \dots, n$ ) is derived.

In order to obtain our results, we use following theorem of Rusheweyh and Stankiewicz [4]:

**Theorem 1** *Let  $F, G \in \mathcal{H}(\mathbb{U})$  be any convex univalent functions in  $\mathbb{U}$ . If  $f \prec F$  and  $g \prec G$ , then*

$$f * g \prec F * G \text{ in } \mathbb{U}.$$

Also, we use a result of Stankiewicz and Stankiewicz [6] which is as follows:

**Theorem 2** *If  $\alpha \leq 1$  and  $\beta \leq 1$ , then*

$$P_\alpha * P_\beta = P_\delta$$

where  $\delta = 1 - 2(1 - \alpha)(1 - \beta)$ .

## 2 Main Results

We may easily generalize Theorem 2 for the classes  $\mathcal{P}(\alpha_i)$  ( $i = 1, 2, \dots, n$ ) and get the following lemma:

**Lemma 1** *If  $\alpha_i \leq 1$  ( $i = 1, 2, \dots, n$ ), then*

$$P_{\alpha_1} * P_{\alpha_2} * \dots * P_{\alpha_n} = P_\delta$$

where

$$\delta = 1 - 2^{n-1}(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_n). \quad (2.1)$$

**Theorem 3** Let for each  $i = 1, 2, \dots, n$ ,  $f_i \in \mathcal{A}$  and  $\alpha_i \leq 1$ . If  $f'_i \in \mathcal{P}(\alpha_i)$  for each  $i = 1, 2, \dots, n$ , and

$$\phi(z) = (f_1 * f_2 * \dots * f_n)(z)$$

then

$$\phi'(z) \prec h(z), \quad z \in \mathbb{U}$$

where

$$h(z) = 1 + 2^n (1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_n) \left[ \frac{z}{2^{n-1}} + \frac{z^2}{3^{n-1}} + \dots \right] \quad (2.2)$$

is convex univalent in  $\mathbb{U}$ .

**Proof.** Let  $f'_i \in \mathcal{P}(\alpha_i)$  for each  $i = 1, 2, \dots, n$ . Then, in view of (1.3), we have for  $\alpha_i \leq 1, i = 1, 2, \dots, n, z \in \mathbb{U}$ ,

$$f'_i(z) \prec q_{\alpha_i}(z) := \frac{1 + (1 - 2\alpha_i)z}{1 - z} = 1 + 2(1 - \alpha_i) \sum_{k=1}^{\infty} z^k \quad (2.3)$$

where the superordinate functions  $q_{\alpha_i}(z)$  for each  $i = 1, 2, \dots, n$  map the disk  $\mathbb{U}$  onto convex univalent regions in the positive half plane. By Theorem 1, we get that

$$f'_1(z) * f'_2(z) * \dots * f'_n(z) \prec q_{\alpha_1}(z) * q_{\alpha_2}(z) * \dots * q_{\alpha_n}(z), \quad z \in \mathbb{U} \quad (2.4)$$

where

$$\begin{aligned} q_{\alpha_1}(z) * q_{\alpha_2}(z) * \dots * q_{\alpha_n}(z) &= 1 + 2^n (1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_n) \sum_{k=1}^{\infty} z^k \\ &= : \frac{1 + (1 - 2\delta)z}{1 - z}, z \in \mathbb{U} \end{aligned}$$

is convex univalent in  $\mathbb{U}$  and in view of Lemma 1,  $\delta$  is given by (2.1).

We know that the function

$$h_1(z) = -\frac{2}{z} [z + \ln(1 - z)] = \sum_{k=1}^{\infty} \frac{2}{k+1} z^k, \quad z \in \mathbb{U}$$

belongs to the class  $\mathcal{K}$  and for  $f \in \mathcal{A}$

$$(f * h_1)(z) = \frac{2}{z} \int_0^z f(t) dt.$$

Therefore, the function

$$h_2(z) = 1 + h_1(z), \quad z \in \mathbb{U}$$

being a translation of  $h_1(z)$ , is convex univalent in  $\mathbb{U}$  and for  $p \in \mathcal{H}[1, 1]$

$$(p * h_2)(z) = -1 + \frac{2}{z} \int_0^z p(t) dt. \quad (2.5)$$

On applying, Theorem 1, to the subordination (2.4) sequentially,  $n - 1$  times with the usual subordination:  $h_2(z) \prec h_2(z)$ ,  $z \in \mathbb{U}$ , we get

$$f'_1 * f'_2 * \dots * f'_n * \underbrace{h_2 * h_2 * \dots * h_2}_{n-1 \text{ times}} \prec q_{\alpha_1} * q_{\alpha_2} * \dots * q_{\alpha_n} * \underbrace{h_2 * h_2 * \dots * h_2}_{n-1 \text{ times}}$$

in  $\mathbb{U}$ , which can also be written as

$$(f'_1 * h_2) * (f'_2 * h_2) * \dots * (f'_{n-1} * h_2) * f'_n \prec (q_{\alpha_1} * h_2) * (q_{\alpha_2} * h_2) * \dots * (q_{\alpha_{n-1}} * h_2) * q_{\alpha_n}. \tag{2.6}$$

On suitably choosing series expansions of  $f'_i$  's and  $q_{\alpha_i}$  's, in view of (2.5), we observe that the subordination (2.6) reduces to

$$\begin{aligned} & \frac{f_1(z)}{z} * \frac{f_2(z)}{z} * \dots * \frac{f_{n-1}(z)}{z} * f'_n(z) \tag{2.7} \\ & \prec \frac{1}{z} \int_0^z q_{\alpha_1}(t) dt * \frac{1}{z} \int_0^z q_{\alpha_2}(t) dt * \dots * \frac{1}{z} \int_0^z q_{\alpha_{n-1}}(t) dt * q_{\alpha_n}(z) \\ & = h(z), z \in \mathbb{U} \end{aligned}$$

where  $h(z)$  is convex univalent in  $\mathbb{U}$  and is of the form (2.2). The left hand side (2.7) of above subordination is

$$(f_1 * f_2 * \dots * f_n)'(z) = \phi'(z).$$

This proves Theorem 3. ■

As the function  $h(z)$  given by (2.2) is convex univalent with real coefficients, we may easily get following result from Theorem 3:

**Corollary 1** *Let for each  $i = 1, 2, \dots, n$ ,  $f_i \in \mathcal{A}$  and  $\alpha_i \leq 1$ . If  $f'_i \in \mathcal{P}(\alpha_i)$  for each  $i = 1, 2, \dots, n$ , and*

$$\phi(z) = (f_1 * f_2 * \dots * f_n)(z)$$

then

$$h(-1) \leq \operatorname{Re} \{ \phi'(z) \} \leq h(1), z \in \mathbb{U}$$

where  $h(z)$  is given by (2.2).

In terms of Zeta function [[7], Ex.5, p.201], we may also find following result from Theorem 3:

**Corollary 2** *Let for each  $i = 1, 2, \dots, n$ ,  $f_i \in \mathcal{A}$  and  $\alpha_i \leq 1$ . If  $f'_i \in \mathcal{P}(\alpha_i)$  for each  $i = 1, 2, \dots, n$ , and*

$$\phi(z) = (f_1 * f_2 * \dots * f_n)(z)$$

then for  $n > 2$ ,

$$\operatorname{Re} \{ \phi'(z) \} \geq 1 + 2^n (1 - \alpha_1) (1 - \alpha_2) \dots (1 - \alpha_n) [(1 - 2^{2-n}) \zeta(n - 1) - 1]$$

and

$$\operatorname{Re} \{ \phi'(z) \} \leq 1 + 2^n (1 - \alpha_1) (1 - \alpha_2) \dots (1 - \alpha_n) [\zeta(n - 1) - 1], \quad z \in \mathbb{U}$$

where  $\zeta$  is well known Zeta Function.

Taking  $n = 3$  and writing  $\zeta(2) = \frac{\pi^2}{6}$ , Corollary 2 provides following result of Sokól [5]:

**Corollary 3** *Let for each  $i = 1, 2, 3$ ,  $f_i \in \mathcal{A}$  and  $\alpha_i \leq 1$ . If  $f'_i \in \mathcal{P}(\alpha_i)$  for each  $i = 1, 2, 3$ , and*

$$\phi(z) = (f_1 * f_2 * f_3)(z)$$

then

$$\operatorname{Re} \{ \phi'(z) \} \geq 1 + 8(1 - \alpha_1)(1 - \alpha_2)(1 - \alpha_3) \left[ \frac{\pi^2}{12} - 1 \right], \quad z \in \mathbb{U}$$

and

$$\operatorname{Re} \{ \phi'(z) \} \leq 1 + 8(1 - \alpha_1)(1 - \alpha_2)(1 - \alpha_3) \left[ \frac{\pi^2}{6} - 1 \right], \quad z \in \mathbb{U}$$

To prove our next result, we prove first a lemma which is as follows:

**Lemma 2** *Let for each  $i = 1, 2, \dots, n$ ,  $f_i \in \mathcal{A}$  and  $\alpha_i \leq 1$ . If  $f'_i \in \mathcal{P}(\alpha_i)$  for each  $i = 1, 2, \dots, n$ , and*

$$\phi(z) = (f_1 * f_2 * \dots * f_n)(z)$$

then there exist some positive integers  $\lambda_1, \lambda_2, \dots, \lambda_{n-2}$  depending upon  $n$  such that

$$\begin{aligned} & \phi'(z) + \lambda_1 z \phi''(z) + \lambda_2 z^2 \phi'''(z) + \dots + \lambda_{n-2} z^{n-2} \phi^{(n-1)}(z) + z^{n-1} \phi^{(n)}(z) \\ &= (f'_1 * f'_2 * \dots * f'_n)(z). \end{aligned}$$

**Proof.** Let  $f_i \in \mathcal{A}$  be of the form

$$f_i(z) = z + \sum_{k=2}^{\infty} a_k^i z^k, \quad z \in \mathbb{U}. \tag{2.8}$$

Then

$$(f'_1 * f'_2 * \dots * f'_n)(z) = 1 + \sum_{k=2}^{\infty} k^n d_k z^{k-1} \tag{2.9}$$

where

$$d_k := a_k^1 a_k^2 \dots a_k^n, \quad k \geq 2. \tag{2.10}$$

We note that for the functions  $f_i(z)$ ,  $i = 1, 2, \dots, n$ , of the form (2.8), the  $r$ -th ( $r \in \mathbb{N}$ ) derivative of  $\phi(z)$  is given by

$$\phi^{(r)}(z) = \sum_{k=1}^{\infty} k(k-1)\dots(k-r+1) d_k z^{k-r},$$



where  $d_1 = 1$  and for  $k \geq 2$ ,  $d_k$  is given by (2.10).

For some positive integers  $\lambda_1, \lambda_2, \dots, \lambda_{n-2}$  depending only upon  $n$ , we have for  $k, n \in \mathbb{N}$ , an identity :

$$k^n \equiv k + \lambda_1 k(k-1) + \lambda_2 k(k-1)(k-2) + \dots \\ + \lambda_{n-2} k(k-1)\dots(k-n+2) + k(k-1)\dots(k-n+1). \quad (2.11)$$

For the positive integers  $\lambda_1, \lambda_2, \dots, \lambda_{n-2}$ , appear in the identity (2.11) and with the use of this identity, we get

$$\phi'(z) + \lambda_1 z \phi''(z) + \lambda_2 z^2 \phi'''(z) + \dots + \lambda_{n-2} z^{n-2} \phi^{(n-1)}(z) + z^{n-1} \phi^{(n)}(z) \\ = \sum_{k=1}^{\infty} k^n d_k z^{k-1}, \text{ where } d_1 = 1.$$

This is the right hand side of (2.9). This proves Lemma 2. ■

**Theorem 4** Let for each  $i = 1, 2, \dots, n$ ,  $f_i \in \mathcal{A}$  and  $\alpha_i \leq 1$ . If  $f'_i \in \mathcal{P}(\alpha_i)$  for each  $i = 1, 2, \dots, n$ , and

$$\phi(z) = (f_1 * f_2 * \dots * f_n)(z)$$

then  $\phi \in \mathcal{K}$  whenever for  $n > 2$ ,

$$(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_n) \leq \frac{3}{2^{n+2} [1 - (1 - 2^{2-n}) \zeta(n-1)]} \quad (2.12)$$

where  $\zeta$  is well known Zeta Function.

**Proof.** Let  $p(z) = \phi'(z)$ , then by Lemma 2 and by (2.4), we get

$$\psi \left( p(z), zp'(z), \dots, z^{n-1} p^{(n-1)}(z) \right) \quad (2.13) \\ = p(z) + \lambda_1 z p'(z) + \lambda_2 z^2 p''(z) + \dots + \lambda_{n-2} z^{n-2} p^{(n-2)}(z) + z^{n-1} p^{(n-1)}(z) \\ = (f'_1 * f'_2 * \dots * f'_n)(z) \prec \frac{1 + (1 - 2\delta)z}{1 - z}, z \in \mathbb{U}$$

where  $\delta$  is given by (2.1). From Theorem 3, we have a possible solution of the above  $n$ -th order Euler-type differential subordination (2.13), as follows

$$\phi'(z) = p(z) \prec h(z), z \in \mathbb{U} \quad (2.14)$$

where  $h(z)$  is given by (2.2).

The  $r$ -th ( $r \in \mathbb{N}$ ) derivative of  $h(z)$  is given by

$$h^{(r)}(z) = 2(1 - \delta) \sum_{k=1}^{\infty} \frac{k(k-1)\dots(k-r+1)}{(k+1)^{n-1}} z^{k-r}.$$

For the positive integers  $\lambda_1, \lambda_2, \dots, \lambda_{n-2}$ , appearing in the identity (2.11), we observe that

$$\begin{aligned} & \psi \left( h(z), zh'(z), \dots, z^{n-1}h^{(n-1)}(z) \right) \\ &= h(z) + \lambda_1 zh'(z) + \lambda_2 z^2 h''(z) + \dots + \lambda_{n-2} z^{n-2} h^{(n-2)}(z) + z^{n-1} h^{(n-1)}(z) \\ &= 1 + 2(1 - \delta) \sum_{k=1}^{\infty} z^k = \frac{1 + (1 - 2\delta)z}{1 - z}, \quad z \in \mathbb{U} \end{aligned}$$

where  $\delta$  is given by (2.1). This verifies the admissibility condition for  $p(z)$  in (2.14) to be a solution of the subordination (2.13).

Now, the function  $\phi \in \mathcal{K}$  if

$$1 + \frac{z\phi''(z)}{\phi'(z)} \prec \frac{1+z}{1-z}, \quad z \in \mathbb{U}$$

or

$$1 + \frac{zp'(z)}{p(z)} \prec \frac{1+z}{1-z}, \quad z \in \mathbb{U}. \tag{2.15}$$

By [[1], Theorem 2.6b, p.60] the condition (2.15) implies

$$p(z) \prec \frac{1}{(1-z)^2}, \quad z \in \mathbb{U}. \tag{2.16}$$

Thus, in view of (2.14), the function  $\phi \in \mathcal{K}$  if

$$h(z) \prec \frac{1}{(1-z)^2}, \quad z \in \mathbb{U}$$

that is if

$$\min_{z \in \mathbb{U}} \Re \{h(z)\} = h(-1) \geq \frac{1}{4}$$

which is the given condition (2.12) if we write the expression of  $h(-1)$  (as it is written in Corollary 2) in terms of Zeta Function [[7], Ex.5, p.201]. This proves the result of Theorem 4. ■

Taking  $n = 3$  in Theorem 4 and on writing  $\zeta(2) = \frac{\pi^2}{6}$ , we get following result.

**Corollary 4** *Let for each  $i = 1, 2, 3$ ,  $f_i \in \mathcal{A}$  and  $\alpha_i \leq 1$ . If  $f'_i \in \mathcal{P}(\alpha_i)$  for each  $i = 1, 2, 3$ , then  $(f_1 * f_2 * f_3)(z) \in \mathcal{K}$  whenever*

$$(1 - \alpha_1)(1 - \alpha_2)(1 - \alpha_3) \leq \frac{9}{8(12 - \pi^2)} \approx 0.53. \tag{2.17}$$

**Remark 1** *We remark that Corollary 4 improves the result of Sokół obtained in [[5], Theorem 2, 124].*

**Acknowledgements.**

This work has been supported by J.R.F of C.S.I.R. (New Delhi, India) awarded to the second author

Authors are thankful to the referee for his/her useful comments.

## References

- [1] S. S. Miller, P. T. Mocanu, *Differential Subordinations*, Marcel Dekker, Inc., 2000.
- [2] St. Ruscheweyh, *Convolution in Geometric Function Theory*, Les Presses de l'Univ. de Montreal, 1982.
- [3] St. Ruscheweyh, T. Sheil-Small, Hadamard product of schlicht functions and the Poyla-Schoenberg conjecture, *Comm. Math. Helv.*, 48 (1973), 119-135.
- [4] St. Ruscheweyh, J. Stankiewicz, Subordination under convex univalent function, *Bull. Pol. Acad. Sci. Math.*, 33 (1985), 499-502.
- [5] J. Sokół, The convexity of Hadamard product of three functions, *J. Math. Appl.*, 29 (2007), 121-125.
- [6] J. Stankiewicz, Z. Stankiewicz, Some Applications of the Hadamard Convolution in the Theory of Functions, *Ann. Univ. Mariae Curie-Sklodowska*, 40 (1986), 251-265.
- [7] E. M. Stein and R. Shakarchi, *Complex Analysis*, Princeton University Press, New Jersey, USA

**DOI: 10.7862/rf.2014.10**

**Poonam Sharma**

email: sharma\_poonam@lkouniv.ac.in

**Rajesh K. Maurya**

email: rajeshkrmaurya@gmail.com

Department of Mathematics

University of Lucknow, Lucknow 226007

*Received 3.10.2013, Accepted 30.11.2013*

## Some inequalities for the polar derivative of a polynomial with restricted zeros

Ahmad Zireh, S. Asie Hosseini

ABSTRACT: Let  $p(z)$  be a polynomial of degree  $n$  and for any complex number  $\alpha$ ,  $D_\alpha p(z) = np(z) + (\alpha - z)p'(z)$  denote the polar derivative of the polynomial  $p(z)$  with respect to  $\alpha$ . In this paper, we obtain new results concerning maximum modulus of the polar derivative of a polynomial with restricted zeros. Our result generalize certain well-known polynomial inequalities.

AMS Subject Classification: *Primary 30A10; Secondary 30C10, 30D15.*

Keywords and Phrases: *Polynomial, Inequality, Maximum modulus, Polar Derivative, Restricted Zeros.*

### 1 Introduction and statement of results

Let  $p(z)$  be a polynomial of degree  $n$ , then according to Bernstein's inequality on the derivative of a polynomial, we have

$$\max_{|z|=1} |p'(z)| \leq n \max_{|z|=1} |p(z)|, \quad (1.1)$$

equality holds in (1.1) if  $p(z)$  has all its zeros at the origin.

The inequality (1.1) can be sharpened, if we restrict ourselves to the class of polynomials having no zeros in  $|z| < 1$ , in fact, P. Erdős conjectured and later Lax [9] proved that if  $p(z) \neq 0$  in  $|z| < 1$ , then (1.1) can be replaced by

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{2} \max_{|z|=1} |p(z)|. \quad (1.2)$$

The result is best possible and equality holds in (1.2) for a polynomial which has all its zeros on  $|z| = 1$ .

If the polynomial  $p(z)$  has all its zeros in  $|z| \leq 1$ , then it was proved by Turan [12] that

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{2} \max_{|z|=1} |p(z)|, \quad (1.3)$$

with equality for those polynomials, which have all their zeros on  $|z| = 1$ . For a polynomial  $p(z)$  of degree at most  $n$  which having no zeros in  $|z| < k$ ,  $k \geq 1$ , inequality (1.2) was generalized by Malik [10] who proved that

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{1+k} \max_{|z|=1} |p(z)|. \quad (1.4)$$

The inequality (1.4) is sharp and equality holds for  $p(z) = (z+k)^n$ . If the polynomial  $p(z)$  has all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , then it was proved by Govil[7] that

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{1+k^n} \max_{|z|=1} |p(z)|. \quad (1.5)$$

The result is best possible and equality holds in (1.5) for  $p(z) = z^n + k^n$ . Let  $\alpha$  be a complex number. For a polynomial  $p(z)$  of degree  $n$ ,  $D_\alpha p(z)$ , the polar derivative of  $p(z)$  is defined as

$$D_\alpha p(z) = np(z) + (\alpha - z)p'(z).$$

It is easy to see that  $D_\alpha p(z)$  is a polynomial of degree at most  $n - 1$ , also  $D_\alpha p(z)$  generalizes the ordinary derivative in the sense that

$$\lim_{\alpha \rightarrow \infty} \left[ \frac{D_\alpha p(z)}{\alpha} \right] = p'(z). \quad (1.6)$$

In order to extend inequality (1.5) for the polar derivative, Aziz and Rather[1] proved that if  $p(z)$  is a polynomial of degree  $n$  having all its zeros in  $|z| \leq k$  where  $k \geq 1$ , then for every real or complex number  $\alpha$  with  $|\alpha| \geq k$ ,

$$\max_{|z|=1} |D_\alpha p(z)| \geq n \frac{|\alpha| - k}{1 + k^n} \max_{|z|=1} |p(z)|. \quad (1.7)$$

The bounds are obtained depends only on the zero of largest modulus and not on the other zeros even if some of them are close to the origin. Therefore, it would be interesting to obtain a bound, which depends on the location of all the zeros of a polynomial. In this connection we use some known ideas in the literature and obtain the following interesting results.

**Theorem 1.1** *Let*

$$p(z) = \sum_{\nu=0}^n a_\nu z^\nu = a_n \prod_{\nu=1}^n (z - z_\nu), \quad a_n \neq 0,$$

*be a polynomial of degree  $n$ ,  $|z_\nu| \leq k_\nu$ ,  $1 \leq \nu \leq n$ , and  $k = \max(k_1, k_2, \dots, k_n) \geq 1$ .*

Then for every real or complex number  $\alpha$  with  $|\alpha| \geq k$ , we have

$$\begin{aligned} \max_{|z|=1} |D_{\alpha}p(z)| &\geq \frac{|\alpha| - k}{1 + k^n} \sum_{\nu=1}^n \frac{k}{k + k_{\nu}} \left[ 2 \max_{|z|=1} |p(z)| + \frac{k^n - 1}{k^n} \min_{|z|=k} |p(z)| \right. \\ &\quad \left. + \frac{4|a_{n-1}|}{k(n+1)} \left( \frac{k^n - 1 - n(k-1)}{n} \right) \right. \\ &\quad \left. + \frac{4|a_{n-2}|}{k^2} \left( \left( \frac{(k^n - 1) - n(k-1)}{n(n-1)} \right) - \left( \frac{(k^{n-2} - 1) - (n-2)(k-1)}{(n-2)(n-3)} \right) \right) \right] \\ &+ \frac{2(k^{n-1} - 1)}{(n+1)k^{n-1}} |na_0 + \alpha a_1| + \frac{1}{k^{n-1}} \left[ \frac{k^{n-1} - 1}{n-1} - \frac{k^{n-3} - 1}{n-3} \right] |(n-1)a_1 + 2\alpha a_2|, \end{aligned} \quad (1.8)$$

for  $n > 3$

and

$$\begin{aligned} \max_{|z|=1} |D_{\alpha}p(z)| &\geq \frac{|\alpha| - k}{1 + k^n} \sum_{\nu=1}^n \frac{k}{k + k_{\nu}} \left[ 2 \max_{|z|=1} |p(z)| + \frac{k^n - 1}{k^n} \min_{|z|=k} |p(z)| \right. \\ &\quad \left. + \frac{4|a_{n-1}|}{k(n+1)} \left( \frac{(k^n - 1) - n(k-1)}{n} \right) + \frac{4|a_{n-2}|(k-1)^n}{k^2 n(n-1)} \right] \\ &\quad + \frac{(k^2 - 1)}{2k^{n-1}} |na_0 + \alpha a_1| + \frac{(k-1)^2}{2k^{n-1}} |(n-1)a_1 + 2\alpha a_2|, \end{aligned} \quad (1.9)$$

for  $n = 3$ .

Since  $\frac{k}{k + k_{\nu}} \geq \frac{1}{2}$  for  $1 \leq \nu \leq n$ , the above theorem gives the following result which is an improvement of the inequality (1.7).

**Corollary 1.2** If  $p(z) = \sum_{\nu=0}^n a_{\nu} z^{\nu}$  is a polynomial of degree  $n$  having all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , then for every real or complex number  $\alpha$  with  $|\alpha| \geq k$ , we have

$$\begin{aligned} \max_{|z|=1} |D_{\alpha}p(z)| &\geq \frac{n(|\alpha| - k)}{1 + k^n} \left[ \max_{|z|=1} |p(z)| + \frac{k^n - 1}{2k^n} \min_{|z|=k} |p(z)| \right. \\ &\quad \left. + \frac{2|a_{n-1}|}{(n+1)k} \left( \frac{k^n - 1 - n(k-1)}{n} \right) \right. \\ &\quad \left. + \frac{2|a_{n-2}|}{k^2} \left( \left( \frac{(k^n - 1) - n(k-1)}{n(n-1)} \right) - \left( \frac{(k^{n-2} - 1) - (n-2)(k-1)}{(n-2)(n-3)} \right) \right) \right] \\ &\quad + \frac{2(k^{n-1} - 1)}{(n+1)k^{n-1}} |na_0 + \alpha a_1| + \frac{1}{k^{n-1}} \left[ \frac{k^{n-1} - 1}{n-1} - \frac{k^{n-3} - 1}{n-3} \right] |(n-1)a_1 + 2\alpha a_2|, \end{aligned} \quad (1.10)$$

for  $n > 3$

and

$$\begin{aligned} \max_{|z|=1} |D_\alpha p(z)| &\geq \frac{n(|\alpha| - k)}{1 + k^n} \left[ \max_{|z|=1} |p(z)| + \frac{k^n - 1}{2k^n} \min_{|z|=k} |p(z)| \right] \\ &+ \frac{2|a_{n-1}|}{(n+1)k} \left( \frac{(k^n - 1) - n(k-1)}{n} \right) + \frac{2|a_{n-2}|(k-1)^n}{k^2 n(n-1)} \\ &+ \frac{(k^2 - 1)}{2k^{n-1}} |na_0 + \alpha a_1| + \frac{(k-1)^2}{2k^{n-1}} |(n-1)a_1 + 2\alpha a_2|, \end{aligned} \quad (1.11)$$

for  $n = 3$ .

Dividing both sides of the inequalities (1.10) and (1.11) by  $|\alpha|$  and letting  $|\alpha| \rightarrow \infty$ , we have the following refinement of the inequality (1.5).

**Corollary 1.3** *If  $p(z) = \sum_{\nu=0}^n a_\nu z^\nu$  is a polynomial of degree  $n$ , having all its zeros in  $|z| \leq k$ ,  $k \geq 1$ , then for every real or complex number  $\alpha$  with  $|\alpha| \geq k$ , we have*

$$\begin{aligned} \max_{|z|=1} |p'(z)| &\geq \frac{n}{1 + k^n} \left[ \max_{|z|=1} |p(z)| + \frac{k^n - 1}{2k^n} \min_{|z|=k} |p(z)| \right] \\ &+ \frac{2|a_{n-1}|}{(n+1)k} \left( \frac{k^n - 1 - n(k-1)}{n} \right) \\ &+ \frac{2|a_{n-2}|}{k^2} \left( \left( \frac{(k^n - 1) - n(k-1)}{n(n-1)} \right) - \left( \frac{(k^{n-2} - 1) - (n-2)(k-1)}{(n-2)(n-3)} \right) \right) \\ &+ \frac{2(k^{n-1} - 1)}{(n+1)k^{n-1}} |a_1| + \frac{1}{k^{n-1}} \left[ \frac{k^{n-1} - 1}{n-1} - \frac{k^{n-3} - 1}{n-3} \right] |2a_2|, \end{aligned} \quad (1.12)$$

for  $n > 3$

and

$$\begin{aligned} \max_{|z|=1} |p'(z)| &\geq \frac{n}{1 + k^n} \left[ \max_{|z|=1} |p(z)| + \frac{k^n - 1}{2k^n} \min_{|z|=k} |p(z)| \right] \\ &+ \frac{2|a_{n-1}|}{(n+1)k} \left( \frac{(k^n - 1) - n(k-1)}{n} \right) + \frac{2|a_{n-2}|(k-1)^n}{k^2 n(n-1)} \\ &+ \frac{(k^2 - 1)}{2k^{n-1}} |a_1| + \frac{(k-1)^2}{2k^{n-1}} |2a_2|, \end{aligned} \quad (1.13)$$

for  $n = 3$ .

As an application of Theorem 1.1 we prove the following result.

**Theorem 1.4** *Let*

$$p(z) = \sum_{\nu=0}^n a_\nu z^\nu = a_n \prod_{\nu=1}^n (z - z_\nu), \quad a_n \neq 0,$$

be a polynomial of degree  $n$ ,  $|z_\nu| \geq k_\nu$ ,  $1 \leq \nu \leq n$ , and  $k = \min(k_1, k_2, \dots, k_n) \leq 1$ . Then for every real or complex number  $\delta$  with  $|\delta| \leq k$ , we have

$$\begin{aligned} \max_{|z|=1} |D_\delta p(z)| &\geq \frac{(k-|\delta|)k^{n-1}}{1+k^n} \sum_{\nu=1}^n \frac{k_\nu}{k+k_\nu} [2 \max_{|z|=1} |p(z)| + \frac{1-k^n}{k^n} \min_{|z|=k} |p(z)|] \\ &+ \frac{4|a_1|k}{(n+1)} \left( \frac{(1-k^n) - n(k^{n-1} - k^n)}{nk^n} \right) + 4|a_2|k^2 \left( \frac{(1-k^n) - n(k^{n-1} - k^n)}{n(n-1)k^n} \right) - \\ &\left( \frac{(1-k^{n-2}) - (n-2)(k^{n-3} - k^{n-2})}{(n-2)(n-3)k^{n-2}} \right) \Big] \\ &+ \frac{2(1-k^{n-1})}{(n+1)} |na_n + \alpha a_{n-1}| + \\ &k^{n-1} \left[ \frac{(1-k^{n-1})}{(n-1)k^{n-1}} - \frac{(1-k^{n-3})}{(n-3)k^{n-3}} \right] |(n-1)a_{n-1} + 2\alpha a_{n-2}|, \end{aligned} \quad (1.14)$$

for  $n > 3$   
and

$$\begin{aligned} \max_{|z|=1} |D_\delta p(z)| &\geq \frac{(k-|\delta|)k^{n-1}}{1+k^n} \sum_{\nu=1}^n \frac{k_\nu}{k+k_\nu} [2 \max_{|z|=1} |p(z)| + \frac{1-k^n}{k^n} \min_{|z|=k} |p(z)|] \\ &+ \frac{4|a_1|k}{(n+1)} \left( \frac{(1-k^n) - n(k^{n-1} - k^n)}{nk^n} \right) + 4|a_2|k^2(1-k^n) \\ &+ \frac{1-k^2}{2} |na_n + \alpha a_{n-1}| + \frac{(1-k)^2}{2} |(n-1)a_{n-1} + 2\alpha a_{n-2}|, \end{aligned} \quad (1.15)$$

for  $n = 3$ .

Since  $\frac{k_\nu}{k+k_\nu} \geq \frac{1}{2}$  for  $1 \leq \nu \leq n$ , then Theorem 1.4 gives the following result.

**Corollary 1.5** Let  $p(z) = \sum_{\nu=0}^n a_\nu z^\nu$  be a polynomial of degree  $n$  does not vanish in  $|z| < k$  where  $k \leq 1$ . Then for every real or complex number  $\delta$  with  $|\delta| \leq k$ , we have

$$\begin{aligned} \max_{|z|=1} |D_\delta p(z)| &\geq \frac{n(k-|\delta|)k^{n-1}}{1+k^n} \left[ \max_{|z|=1} |p(z)| + \frac{1-k^n}{2k^n} \min_{|z|=k} |p(z)| \right] \\ &+ \frac{2|a_1|k}{(n+1)} \left( \frac{(1-k^n) - n(k^{n-1} - k^n)}{nk^n} \right) + 2|a_2|k^2 \left( \frac{(1-k^n) - n(k^{n-1} - k^n)}{n(n-1)k^n} \right) - \\ &\left( \frac{(1-k^{n-2}) - (n-2)(k^{n-3} - k^{n-2})}{(n-2)(n-3)k^{n-2}} \right) \Big] \\ &+ \frac{2(1-k^{n-1})}{(n+1)} |na_n + \alpha a_{n-1}| + \\ &k^{n-1} \left[ \frac{(1-k^{n-1})}{(n-1)k^{n-1}} - \frac{(1-k^{n-3})}{(n-3)k^{n-3}} \right] |(n-1)a_{n-1} + 2\alpha a_{n-2}|, \end{aligned} \quad (1.16)$$



for  $n > 3$   
and

$$\begin{aligned} \max_{|z|=1} |D_\delta p(z)| &\geq \frac{n(k-|\delta|)k^{n-1}}{1+k^n} \left[ \max_{|z|=1} |p(z)| + \frac{1-k^n}{2k^n} \min_{|z|=k} |p(z)| \right] \\ &+ \frac{2|a_1|k}{(n+1)} \left( \frac{(1-k^n) - n(k^{n-1} - k^n)}{nk^n} \right) + 2|a_2|k^2(1-k)^n \\ &+ \frac{1-k^2}{2} |na_n + \alpha a_{n-1}| + \frac{(1-k)^2}{2} |(n-1)a_{n-1} + 2\alpha a_{n-2}|, \end{aligned} \quad (1.17)$$

for  $n = 3$ .

## 2 Lemmas

For proof of the theorems, we need the following lemmas. The first lemma is due to Dewan, Kaur and Mir [4].

**Lemma 2.1** *If  $p(z)$  is a polynomial of degree  $n$ , then for  $R \geq 1$ ,*

$$\begin{aligned} \max_{|z|=R} |p(z)| &\leq R^n \max_{|z|=1} |p(z)| - \frac{2(R^n - 1)}{n+2} |p(0)| \\ &- \left[ \frac{R^n - 1}{n} - \frac{R^{n-2} - 1}{n-2} \right] |p'(0)| \end{aligned} \quad (2.1)$$

if  $n > 2$ , and

$$\max_{|z|=R} |p(z)| \leq R^n \max_{|z|=1} |p(z)| - \frac{R-1}{2} [(R+1)|p(0)| + (R-1)|p'(0)|] \quad (2.2)$$

if  $n = 2$ .

**Lemma 2.2** *If  $p(z)$  is a polynomial of degree  $n$ , having no zeros in  $|z| < 1$ , then for  $R \geq 1$ ,*

$$\begin{aligned} \max_{|z|=R} |p(z)| &\leq \frac{R^n + 1}{2} \max_{|z|=1} |p(z)| - \frac{R^n - 1}{2} \min_{|z|=1} |p(z)| - \\ &\frac{2}{n+1} \left[ \frac{(R^n - 1)}{n} - (R-1) \right] |p'(0)| - \\ &\left[ \frac{(R^n - 1) - n(R-1)}{n(n-1)} - \frac{(R^{n-2} - 1) - (n-2)(R-1)}{(n-2)(n-3)} \right] |p''(0)| \end{aligned} \quad (2.3)$$

if  $n > 3$ , and

$$\begin{aligned} \max_{|z|=R} |p(z)| &\leq \frac{R^n + 1}{2} \max_{|z|=1} |p(z)| - \frac{R^n - 1}{2} \min_{|z|=1} |p(z)| \\ &- \frac{2}{n+1} \left[ \frac{R^n - 1}{n} - (R-1) \right] |p'(0)| \\ &- \frac{(R-1)^n}{n(n-1)} |p''(0)| \end{aligned} \quad (2.4)$$

if  $n = 3$ .

This lemma is due to Dewan, Singh and Mir [5].

**Lemma 2.3** *If  $p(z) = a_n \prod_{\nu=1}^n (z - z_\nu)$ ,  $a_n \neq 0$ , is a polynomial of degree  $n$ , such that  $|z_\nu| \leq 1$ ,  $1 \leq \nu \leq n$ , then*

$$\max_{|z|=1} |p'(z)| \geq \sum_{\nu=1}^n \frac{1}{1 + |z_\nu|} \max_{|z|=1} |p(z)|. \quad (2.5)$$

This lemma is due to Giroux, Rahman and Schmeisser [8].

**Lemma 2.4** *If  $p(z)$  is a polynomial of degree  $n$  and  $\alpha$  is any real or complex number with  $|\alpha| \neq 0$ , then for  $|z| = 1$*

$$|D_\alpha q(z)| = |n\bar{\alpha}p(z) + (1 - \bar{\alpha}z)p'(z)|, \quad (2.6)$$

where  $q(z) = z^n \overline{p(\frac{1}{\bar{z}})}$ .

This lemma is due to Aziz [2].

### 3 Proofs of the theorems

**Proof of the Theorem 1.1.** Let  $G(z) = p(kz)$ . Since all the zeros of  $p(z)$  lie in  $|z| \leq k$ , then all the zeros of  $G(z)$  lie in  $|z| \leq 1$ . Now on applying Lemma 2.3 to the polynomial  $G(z)$ , we get

$$\max_{|z|=1} |G'(z)| \geq \sum_{\nu=1}^n \frac{1}{1 + \frac{|z_\nu|}{k}} \max_{|z|=1} |G(z)|. \quad (3.1)$$

Let  $H(z) = z^n \overline{G(1/\bar{z})}$ . Then it can be easily verified that for  $|z| = 1$

$$|H'(z)| = |nG(z) - zG'(z)|. \quad (3.2)$$

The polynomial  $G(z)$  has all its zeros in  $|z| \leq 1$  and  $|H(z)| = |G(z)|$  for  $|z| = 1$ , therefore, by Gauss-Lucas theorem for  $|z| = 1$ , we have

$$|H'(z)| \leq |G'(z)|. \quad (3.3)$$

Now for every real or complex number  $\alpha$  with  $|\alpha| \geq k$ , we have

$$|D_{\frac{\alpha}{k}} G(z)| = |nG(z) - zG'(z) + \frac{\alpha}{k} G'(z)| \geq \frac{|\alpha|}{k} |G'(z)| - |nG(z) - zG'(z)|. \quad (3.4)$$

This gives with the help of (3.2) and (3.3) that

$$\max_{|z|=1} |D_{\frac{\alpha}{k}} G(z)| \geq \frac{|\alpha| - k}{k} \max_{|z|=1} |G'(z)|. \quad (3.5)$$

Using (3.1) in (3.4), we get

$$\max_{|z|=1} |D_{\frac{\alpha}{k}} G(z)| \geq \frac{|\alpha| - k}{k} \sum_{\nu=1}^n \frac{k}{k + |z_\nu|} \max_{|z|=1} |G(z)|. \quad (3.6)$$

Replacing  $G(z)$  by  $p(kz)$ , we get

$$\max_{|z|=1} |D_{\frac{\alpha}{k}} p(kz)| \geq (|\alpha| - k) \sum_{\nu=1}^n \frac{1}{k + |z_\nu|} \max_{|z|=1} |p(kz)|. \quad (3.7)$$

which implies

$$\max_{|z|=1} |np(kz) + (\frac{\alpha}{k} - z)kp'(kz)| \geq (|\alpha| - k) \sum_{\nu=1}^n \frac{1}{k + |z_\nu|} \max_{|z|=1} |p(kz)|.$$

which gives

$$\max_{|z|=k} |D_\alpha p(z)| \geq (|\alpha| - k) \sum_{\nu=1}^n \frac{1}{k + |z_\nu|} \max_{|z|=k} |p(z)|. \quad (3.8)$$

The polynomial  $p(z)$  is of degree  $n > 3$  and so  $D_\alpha p(z)$  is the polynomial of degree  $n - 1$ , where  $n - 1 > 2$ , hence applying Lemma 2.1 to the polynomial  $D_\alpha p(z)$ , we get

$$\begin{aligned} \max_{|z|=k} |D_\alpha p(z)| &\leq k^{n-1} \max_{|z|=1} |D_\alpha p(z)| - \frac{2(k^{n-1} - 1)}{n + 1} |na_0 + \alpha a_1| \\ &\quad - \left[ \frac{k^{n-1} - 1}{n - 1} - \frac{k^{n-3} - 1}{n - 3} \right] |(n - 1)a_1 + 2\alpha a_2|. \end{aligned} \quad (3.9)$$

Combining (3.9) and (3.8), we get

$$\begin{aligned} \max_{|z|=1} |D_\alpha p(z)| &\geq \frac{|\alpha| - k}{k^{n-1}} \left[ \sum_{\nu=1}^n \frac{1}{k + |z_\nu|} \max_{|z|=k} |p(z)| \right] + \\ &\frac{2(k^{n-1} - 1)}{(n + 1)k^{n-1}} |na_0 + \alpha a_1| + \frac{1}{k^{n-1}} \left[ \frac{k^{n-1} - 1}{n - 1} - \frac{k^{n-3} - 1}{n - 3} \right] |(n - 1)a_1 + 2\alpha a_2|. \end{aligned} \quad (3.10)$$

Since the polynomial  $p(z)$  has all zeros in  $|z| \leq k$ ,  $k \geq 1$ , then  $q(z) = z^n p(1/z)$  has no zero in  $|z| < 1/k$ , hence the polynomial  $q(z/k)$  having no zeros in  $|z| < 1$ , therefore on applying Lemma 2.2 to the polynomial  $q(z/k)$ , we get

$$\begin{aligned} \max_{|z|=k} |q(z/k)| &\leq \\ &\left[ \frac{k^n + 1}{2} \right] \max_{|z|=1} |q(z/k) - (\frac{k^n - 1}{2}) \min_{|z|=1} |q(z/k)| - \frac{2|a_{n-1}|}{(n + 1)k} \left[ \frac{k^n - 1}{n} - (k - 1) \right] \\ &\quad - \frac{2|a_{n-2}|}{k^2} \left[ \left( \frac{k^n - 1}{n(n - 1)} \right) - \left( \frac{k^{n-2} - 1}{(n - 2)(n - 3)} \right) \right]. \end{aligned} \quad (3.11)$$

Since  $\max_{|z|=1} |q(z/k)| = (1/k^n) \max_{|z|=k} |p(z)|$ , and  $\min_{|z|=1} |q(z/k)| = (1/k^n) \min_{|z|=k} |p(z)|$ , then (3.11) is equivalent to

$$\begin{aligned} \max_{|z|=k} |p(z)| &\geq \left(\frac{2k^n}{k^n+1}\right) \max_{|z|=1} |p(z)| + \left(\frac{k^n-1}{k^n+1}\right) \min_{|z|=k} |p(z)| \\ &+ \frac{4k^{n-1}|a_{n-1}|}{(n+1)(k^n+1)} \left[\frac{k^n-1}{n} - (k-1)\right] \\ &+ \frac{4k^{n-2}|a_{n-2}|}{k^n+1} \left[\left(\frac{(k^n-1)-n(k-1)}{n(n-1)}\right) - \left(\frac{(k^{n-2}-1)-(n-2)(k-1)}{(n-2)(n-3)}\right)\right]. \end{aligned} \quad (3.12)$$

Combining (3.10) and (3.12), we get

$$\begin{aligned} \max_{|z|=1} |D_\alpha p(z)| &\geq \frac{|\alpha| - k}{k^{n-1}} \sum_{\nu=1}^n \frac{1}{k+k_\nu} \left[\left(\frac{2k^n}{k^n+1}\right) \max_{|z|=1} |p(z)|\right. \\ &+ \left(\frac{k^n-1}{k^n+1}\right) \min_{|z|=k} |p(z)| + \frac{4k^{n-1}|a_{n-1}|}{(n+1)(k^n+1)} \left(\frac{k^n-1}{n} - (k-1)\right) \\ &+ \left.\frac{4k^{n-2}|a_{n-2}|}{k^n+1} \left(\left(\frac{(k^n-1)-n(k-1)}{n(n-1)}\right) - \left(\frac{(k^{n-2}-1)-(n-2)(k-1)}{(n-2)(n-3)}\right)\right)\right] \\ &+ \frac{2(k^{n-1}-1)}{(n+1)k^{n-1}} |na_0 + \alpha a_1| + \frac{1}{k^{n-1}} \left[\frac{k^{n-1}-1}{n-1} - \frac{k^{n-3}-1}{n-3}\right] \\ &\times |(n-1)a_1 + 2\alpha a_2|. \end{aligned} \quad (3.13)$$

which completes the proof of (1.10). The proof of the Theorem 1.1 in the case  $n = 3$  follows along the same lines as the proof of (1.10) but instead of inequalities (2.1) and (2.3), we use inequalities (2.2) and (2.4), respectively.  $\square$

**Proof of the Theorem 1.4.** Let  $q(z) = z^n \overline{p(1/\bar{z})}$  then  $1/|z_\nu| \leq 1/k_\nu$  for  $1 \leq \nu \leq n$  such that  $1/k = \max(1/k_1, 1/k_2, \dots, 1/k_n) \geq 1$ . On applying Theorem 1.1 to the polynomial  $q(z)$ , we get

$$\begin{aligned} \max_{|z|=1} |D_\alpha q(z)| &\geq (|\alpha| - 1/k) k^{n-1} \sum_{\nu=1}^n \frac{1}{1/k + 1/k_\nu} \left[\left(\frac{2/k^n}{1/k^n+1}\right) \max_{|z|=1} |q(z)|\right. \\ &+ \left(\frac{1/k^n-1}{1/k^n+1}\right) \min_{|z|=1/k} |q(z)| + \frac{4|a_1|}{(1+1/k^n)k^{n-1}(n+1)} \left(\frac{(1/k^n-1)-n(1/k-1)}{n}\right) \\ &+ \left.\frac{4|a_2|}{k^{n-2}(1+1/k^n)} \left(\left(\frac{(1/k^n-1)-n(1/k-1)}{n(n-1)}\right) - \left(\frac{(1/k^{n-2}-1)-(n-2)(1/k-1)}{(n-2)(n-3)}\right)\right)\right] \\ &+ \frac{2k^{n-1}(1/k^{n-1}-1)}{(n+1)} |na_n + \alpha a_{n-1}| + k^{n-1} \left[\frac{(1/k^{n-1}-1)}{n-1} - \frac{(1/k^{n-3}-1)}{n-3}\right] \\ &\times |(n-1)a_{n-1} + 2\alpha a_{n-2}| \end{aligned} \quad (3.14)$$

Now from lemma 2.4 it follows that for  $|z| = 1$ ,  $|D_\alpha q(z)| = |\alpha| |D_{\frac{1}{\alpha}} p(z)|$  Using the above equality in (3.14), we get for  $|\alpha| \geq 1/k$ ,

$$\begin{aligned}
|\alpha| \max_{|z|=1} |D_{\frac{1}{\alpha}} p(z)| &\geq (|\alpha| - 1/k) k^{n-1} \sum_{\nu=1}^n \frac{k k_\nu}{k + k_\nu} \left[ \left( \frac{2}{1 + k^n} \right) \max_{|z|=1} |p(z)| \right. \\
&+ \frac{1}{k^n} \left( \frac{1 - k^n}{1 + k^n} \right) \min_{|z|=k} |p(z)| + \frac{4|a_1|k}{(n+1)(k^n+1)} \left( \frac{(1 - k^n) - n(k^{n-1} - k^n)}{nk^n} \right) \\
&+ \left. \frac{4|a_2|k^2}{(1+k^n)} \left( \left( \frac{(1 - k^n) - n(k^{n-1} - k^n)}{n(n-1)k^n} \right) - \left( \frac{(1 - k^{n-2}) - (n-2)(k^{n-3} - k^{n-2})}{(n-2)(n-3)k^{n-2}} \right) \right) \right] \\
&+ \frac{2(1 - k^{n-1})}{(n+1)} |na_n + \alpha a_{n-1}| + k^{n-1} \left[ \frac{(1 - k^{n-1})}{(n-1)k^{n-1}} - \frac{(1 - k^{n-3})}{(n-3)k^{n-3}} \right] \\
&\times |(n-1)a_{n-1} + 2\alpha a_{n-2}|,
\end{aligned} \tag{3.15}$$

Replacing  $\frac{1}{\alpha}$  by  $\delta$ , so that  $|\delta| \leq k$ , we get from (3.15)

$$\begin{aligned}
\left| \frac{1}{\delta} \right| \max_{|z|=1} |D_\delta p(z)| &\geq \left( \left| \frac{1}{\delta} \right| - 1/k \right) k^{n-1} \sum_{\nu=1}^n \frac{k k_\nu}{k + k_\nu} \left[ \left( \frac{2}{1 + k^n} \right) \max_{|z|=1} |p(z)| \right. \\
&+ \frac{1}{k^n} \left( \frac{1 - k^n}{1 + k^n} \right) \min_{|z|=k} |p(z)| + \frac{4|a_1|k}{(n+1)(k^n+1)} \left( \frac{(1 - k^n) - n(k^{n-1} - k^n)}{nk^n} \right) \\
&+ \left. \frac{4|a_2|k^2}{(1+k^n)} \left( \left( \frac{(1 - k^n) - n(k^{n-1} - k^n)}{n(n-1)k^n} \right) - \left( \frac{(1 - k^{n-2}) - (n-2)(k^{n-3} - k^{n-2})}{(n-2)(n-3)k^{n-2}} \right) \right) \right] \\
&+ \frac{2(1 - k^{n-1})}{(n+1)} |na_n + \alpha a_{n-1}| + k^{n-1} \left[ \frac{(1 - k^{n-1})}{(n-1)k^{n-1}} - \frac{(1 - k^{n-3})}{(n-3)k^{n-3}} \right] \\
&\times |(n-1)a_{n-1} + 2\alpha a_{n-2}|.
\end{aligned} \tag{3.16}$$

Or

$$\begin{aligned}
\max_{|z|=1} |D_\delta p(z)| &\geq (k - |\delta|) k^{n-1} \sum_{\nu=1}^n \frac{k_\nu}{k + k_\nu} \left[ \left( \frac{2}{1 + k^n} \right) \max_{|z|=1} |p(z)| \right. \\
&+ \frac{1}{k^n} \left( \frac{1 - k^n}{1 + k^n} \right) \min_{|z|=k} |p(z)| + \frac{4|a_1|k}{(n+1)(k^n+1)} \left( \frac{(1 - k^n) - n(k^{n-1} - k^n)}{nk^n} \right) \\
&+ \left. \frac{4|a_2|k^2}{(1+k^n)} \left( \left( \frac{(1 - k^n) - n(k^{n-1} - k^n)}{n(n-1)k^n} \right) - \left( \frac{(1 - k^{n-2}) - (n-2)(k^{n-3} - k^{n-2})}{(n-2)(n-3)k^{n-2}} \right) \right) \right] \\
&+ \frac{2(1 - k^{n-1})}{(n+1)} |na_n + \alpha a_{n-1}| + k^{n-1} \left[ \frac{(1 - k^{n-1})}{(n-1)k^{n-1}} - \frac{(1 - k^{n-3})}{(n-3)k^{n-3}} \right] \\
&\times |(n-1)a_{n-1} + 2\alpha a_{n-2}|.
\end{aligned} \tag{3.17}$$

Which is (1.14). The proof of the Theorem 1.4 in the case  $n = 3$  follows along the same lines as the proof of Theorem 1.1. Hence the proof of Theorem 1.4 is complete.  $\square$

## References

- [1] A. Aziz and N. A. Rather, *A refinement of a theorem of Paul Turan concerning polynomials*, J. Math. Ineq. Appl, **1** (1998), 231-238.
- [2] A. Aziz, *Inequalities for the polar derivative of a polynomial*, J. Approx. Theory, **55** (1988), 183-193.
- [3] S. Bernstein, *Leons sur les propriétés extrémales et la meilleure approximation des fonctions analytiques d'une variable réelle*. Gauthier Villars, Paris, 1926.
- [4] K. K. Dewan, J. Kaur and A. Mir, *Inequalities for the derivative of a polynomial*, J. Math. Anal. Appl, **269** (2002), 489-499.
- [5] K. K. Dewan, N. Singh and A. Mir, *Growth of polynomials not vanishing inside a circle*, Int. J. Math. Anal, **1** (2007), 529-538.
- [6] K. K. Dewn and C. M. Upadhye, *Inequalities for the polar derivative of a polynomial*, J. Inequal. Pure and Appl. Math, **9** (2008), 1-9.
- [7] N.K. Govil, *On the derivative of a polynomial*, Proc. Amer. Math. Soc, **41** (1973), 543-546.
- [8] A. Giroux, Q. I. Rahman and G. Schmeisser, *On Bernstein's inequality*, Can. J. Math, **31** (1979), 347-353.
- [9] P. D. Lax, *Proof of a conjecture of P. Erdős on the derivative of a polynomial*, Bull. Amer. Math. Soc, **50** (1944), 509-513.
- [10] M. A. Malik, *On the derivative of a polynomial*, J. London. Math. Soc, **1** (1969), 57-60.
- [11] V. K. Jain, *Generalization of certain well known inequalities for polynomial*, Glas. Mat, **32** (1997), 45-51.
- [12] P. Turan, *Über die Ableitung von Polynomen*, Compositio Math, **7** (1939), 89-95.

**DOI: 10.7862/rf.2014.11**

**Ahmad Zireh – Corresponding author**

email: [azireh@shahroodut.ac.ir](mailto:azireh@shahroodut.ac.ir)

**S. Asie Hosseini**

email: [sarah.panic@yahoo.com](mailto:sarah.panic@yahoo.com)

Department of Mathematics,  
Shahrood University of Technology,  
Shahrood, Iran

*Received 13.12.2013, Accepted 12.02.2014*

## **ADDITIONAL INFORMATION**

**The Journal of Mathematics and Applications publishes a list of reviewers on the website:**

<http://oficyna.portal.prz.edu.pl/pl/zeszyty-naukowe/journal-of-math/>

<http://oficyna.portal.prz.edu.pl/en/>

**The journal uses as described on its website the procedure for reviewing:**

<http://oficyna.portal.prz.edu.pl/pl/zasady-recenzowania/>

<http://oficyna.portal.prz.edu.pl/en/>

**Information for authors available at:**

<http://oficyna.portal.prz.edu.pl/informacje-dla-autorow/>

<http://oficyna.portal.prz.edu.pl/en/>

**Review's form available at:**

<http://oficyna.portal.prz.edu.pl/pl/zeszyty-naukowe/journal-of-math/>

<http://oficyna.portal.prz.edu.pl/en/>

**Instruction for Authors**

<http://oficyna.portal.prz.edu.pl/pl/instrukcja-dla-autorow/>

<http://oficyna.portal.prz.edu.pl/en/>

**Contact details to Editorial Office available at:**

<http://oficyna.portal.prz.edu.pl/pl/zeszyty-naukowe/journal-of-math/>

<http://oficyna.portal.prz.edu.pl/en/>

**Electronic version of the published articles available at:**

<http://oficyna.portal.prz.edu.pl/pl/zeszyty-naukowe/journal-of-math/>

<http://oficyna.portal.prz.edu.pl/en/>

Cooperating Reviewers

Mohammad W. Alomari (Jordan)  
Józef Banaś (Poland)  
Karol Baron (Poland)  
Urszula Bednarz (Poland)  
S.H. Bhambri (India)  
R. Castillo (Wenezuela)  
Fabrizio Catanese (Germany)  
Bharath Chandran (India)  
C.S. Chen (USA)  
Krzysztof Ciesielski (Poland)  
P. Dhena (India)  
Janusz Dronka (Poland)  
Urszula Dudziak (Poland)  
Jacek Dziok (Poland)  
Elmetwally Mohamed Elabbasy (Egypt)  
Richard Fournier (Canada)  
Roman J. Ger (Poland)  
Janusz Godula (Poland)  
N. K. Govil (India)  
Jarosław Górnicki (Poland)  
E.A. Haroutunian (Armenia)  
Ping He (China)  
Kostaq Hila (Albania)  
W.K. Jain (India)  
Jasmine James (India)  
Sunil Jacob John (India)  
Y.B. Jun (Korea)  
Henryk Hudzik (Poland)  
Andrzej Jan Kamiński (Poland)  
Sukhendu Kar (India)  
Leopold Koczan (Poland)  
V.F. Kosheler (Russia)  
Joanna Kowalczyk (Poland)  
Jan Kurek (Poland)  
Rency Kurian (India)  
Adam Lecko (Poland)  
Edyta Leś (Poland)  
Alexander Linkov (Poland)  
Marian Matłoka (Poland)  
Roman F. Maciej Mączyński (Poland)  
Genadij Miszuris (England, Poland)  
Gangadharun Murugusundaramoorthy (India)  
Jamal M.M. Mustafa (Jordan)  
Leszek Olszowy (Poland)  
M. Kemal Ozdemir (Turkey)  
Antoni Pardała (Poland)  
Hari Sing Parihar (India)  
Krzysztof Piejko (Poland)  
E. Anzari Piri (Iran)  
Kulpid Raj (India)  
Donal O'Regan (Ireland)  
Simeon Reich (Israel)  
Beata Rzepka (Poland)  
Maciej Sablik (Poland)  
Fatemeh E.K. Saraei (Iran)  
A. Boruand Sayed (Iran)  
T.M. Seudy (Egypt)  
Yilin Shang (USA)  
Mohamed el Sheikh (Egypt)  
Janusz Sokół (Poland)  
Hari Mohan Srivastava (Canada)  
Jan Stankiewicz (Poland)  
Zbigniew Suraj (Poland)  
Andrzej Swankowski (Poland)  
Anna Szpila (Poland)  
Jan Szynal (Poland)  
Lucyna Trojnar Spelina (Poland)  
Bronisław Wajnryb (Poland)  
Z.G. Wang (China)  
Katarzyna Wilczek (Poland)  
Agnieszka Wiśniowska Wajnryb (Poland)  
Andrzej Włoch (Poland)  
Iwona Włoch (Poland)  
Andrzej Wrzesień (Poland)  
Paweł Zaprawa (Poland)  
Jaroslav Zemanek (Poland)

The Journal of Mathematics and Applications publishes a list of reviewers  
in vol. 37 and on the website:  
<http://oficyna.portal.prz.edu.pl/pl/zeszyty-naukowe/journal-of-math/>